

SUPPLEMENTAL LABEL

PYRAFLUFEN-ETHYL	GROUP	14	HERBICIDE
------------------	-------	----	-----------



EPA Reg. No. 71711-25

Crops: Bearing and Nonbearing - Pome Fruit Group; Pomegranate; Small Fruit Vine Climbing Subgroup Except Fuzzy Kiwifruit; Stone Fruit Group; Tree Nut Group; Tropical and Subtropical, Small Fruit, Edible Peel Subgroup

This supplemental label expires September 30, 2022 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for VENUE® Herbicide before applying. Use of VENUE Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for VENUE Herbicide. New use directions appear on this supplemental label that may be different from those that appear on the container label.

CROP USE DIRECTIONS

BEARING AND NONBEARING

Pome Fruit Group (Crop Group 11-10) apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

Pomegranate

Small Fruit Vine Climbing Subgroup - Except Fuzzy Kiwifruit (Crop Group 13-07F) amur river grape; gooseberry; grape; kiwifruit, hardy; Maypop; schisandra berry; cultivars varieties, and/or hybrids of these

Stone Fruit Group (Crop Group 12-12) apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these

Tree Nut Group (Crop Group 14-12) African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (Filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these

Tropical and Subtropical, Small Fruit, Edible Peel Subgroup (Crop Subgroup 23A) acerola; African plum; agritos; almondette; appleberry; arbutus berry; bayberry, red; bignay; breadnut; cabeluda; carandas-plum; Ceylon iron wood; Ceylon olive; cherry-of-the-Rio-Grande; Chinese olive, black; Chinese olive, white; chirauli-nut; cocoplum; desert-date; false sandalwood; fragrant manjack; gooseberry, abyssinian; gooseberry, Ceylon; gooseberry, otaheite; governor's plum; grumichama; guabiroba; guava berry; guava, Brazilian; guava, Costa Rican; guayabillo; illawarra plum; Indian-plum; Jamaica-cherry; jambolan; kaffir-plum; kakadu plum; kapundung; karanda; lemon aspen; mombin, yellow; monos plum; mountain cherry; olive; persimmon, black; pitomba; plum-of-Martinique; rukam; rumberry; sea grape; sete-capotes; silver aspen; water apple; water pear; water berry; wax jambu; cultivars, varieties, and hybrids of these commodities

Application	Pest	Rate/Acre	Maximum Applications Per Year	Application Instructions
Postharvest Dormant Prebloom	Listed Broadleaf Weeds	3.0 to 4.0 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	Do not exceed 3 applications per year for this use.	<ul style="list-style-type: none"> Apply in a minimum of 20 gallons spray solution per acre by ground equipment to target weeds and sucker growth. Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions. COC adjuvants are advised, although other adjuvants may be used. Use the higher rate for hard-to-control weeds. For the management of undesirable sucker growth on the basal portion of trunks, root sprouts, and vine trunks, growth must be controlled when the tissue is young, immature, and/or not hardened off. Avoid contact with green, uncallused bark of young trees or vines established less than one year unless protected from spray contact by non-porous wraps, grow tubes, or waxed containers. <p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> Do not apply by air for this use. Do not allow spray to drift onto desirable fruit, foliage, vines, or trees as damage will occur. Do not exceed 4.0 fl oz (0.0053 lb ai) per acre in a single application for these uses. Allow a minimum of 30 days between applications for this use. Do not exceed 6.8 fl oz (0.0090 lb ai) per acre per year for all postharvest, dormant, and pre-bloom applications combined. Do not exceed 12.0 fl oz (0.0159 lb ai) per acre per year for all in-season applications combined. Preharvest Interval (PHI): 0 days
	Sucker Management	3.0 to 4.0 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	Do not exceed 2 applications per year for this use.	
In-Season	Listed Broadleaf Weeds	3.0 to 4.0 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	Do not exceed a combined total of 3 applications per year for these uses.	
	Sucker Management			

©2019 Nichino America, Inc. Venue and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA®

Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700