

# Water Guard WG 4500 (ANTIMICROBIAL SOLUTION)

WATER GUARD™ WG 4500 is a peroxyacetic acid-based sanitizer/disinfectant developed for the following uses:

Institutional/Industrial Sanitizer and Disinfectant for Previously Cleaned Hard Non-Porous

Food Contact Surfaces in: Dairies, Wineries, Breweries, Food and Beverage Plants, Poultry and Egg Facilities, and Animal Housing.

Hard, Non-Porous Surface Disinfection in: Hospitals, Schools, Industrial Facilities, Office Buildings, Veterinary Clinics.

Bacteria, Slime, Odor and Algae Control in: Recirculating Cooling Water and Evaporative Coolers, Reverse Osmosis, Nano and Ultra Filtration, and Agricultural Waters

### **ACTIVE INGREDIENTS:** Darassus anatio Anid

reitxyacelic	9010	5.0 %
Hydrogen Per	oxide	26.5%
INERT INGREDIENTS		. 67.9%
TOTAL		100.0%
EPA Reg. No. 63838-1-8917	EPA Est. No. 63838-CA-01: 63838-AR-001	

EPA Est. No. 63838-CA-01: 63838-AR-001

Before Using This Product, Please Read This Entire Label Carefully

## **KEEP OUT OF REACH OF CHILDREN** DANGER PELIGRO

E C0/

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand this label, find someone to explain it to you in detail.)

Note to Reviewer: In accordance with 40 CFR 156.68(d), all first ald statements, as prescribed, will appear on the front panel of the product label.

### FIRST AID

IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes.     Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.     Call a poison control center or doctor for treatment advice.	
IF ON SKIN OR CLOTHING	Take off contaminated clothing.     Rinse skin immediately with plenty of water for 15-20 minutes.     Call a poison control center or doctor for treatment advice.	
IF INHALED	Move person to fresh air.     If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.     Call a poison control center or doctor for treatment advice.	
IF SWALLOWED	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.	
QUESTIONS? 1-209-581-9576	Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN:	Probable mucosal damage may contraindicate the use of gastric lavage.	

### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER CORROSIVE: Do not enter an enclosed area without proper respiratory protection. Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes, on skin, or on clothing. Wear goggles and face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Do not enter an enclosed area without proper resoiratory protection. Wear a minimum of a NIOSH-approved elastomeric half mask respirator with organic vapor (OV) cartridges and combination N1. R. or P filters; or a NIOSH-approved gas mask with OV canisters; or a NIOSH-approved powered air purifying respirator with OV cartridges and combination HE filters when handling concentrate product

PHYSICAL OR CHEMICAL HAZARDS.

STRONG OXIDIZING AGENT. CORROSIVE: Mix only with water fand adjuvant if applicablel below 140° F. Product must be diluted in accordance with label directions prior to use. At temperatures exceeding 156° F, decomposition occurs releasing oxygen. The oxygen released could initiate combustion.

ENVIRONMENTAL HAZARDS This pesticide is toxic to hirds, fish and aquatic invertebrates. Caution must be used when applying indoors because pets may be at risk. Do not discharge effluent containing this product into lakes streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of the National Pollutant Discharge System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage plant authority.

PERSONAL PROTECTIVE FOLIPMENT Handlers who may be exposed to the undiluted product through mixing, loading, application, or other tasks must wear: coveralls over long-sleeved shirt and long pants, rubber gloves, chemical

resistant footwear plus socks, and protective eyewear (goggles or face shield).

Handlers who may be exposed to the diluted product through application or other tasks must wear: long-sleeved shirt and long pants, and shoes plus socks.
Follow manufacturer's instructions for cleaning/maintaining PPE. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Users should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gel inside. Then wash thoroughly and out on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thorough and change into clean clothing.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Note: All volumes given in ounces are fluid ounces.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

### AGRICUI TURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers or arms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contain specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and Restricted-Entry Interval (REI). The requirements in this box only apply to the uses of this product that are covered by the Workers Protection Standard. There is a restricted entry of zero (0) hours for this product.

### NON-AGRICUI TURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies whe this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses

Keep unprotected persons out of treated areas until sprays have dried.

This peroxyacetic acid sanitizer is recommended for use on precleaned surfaces such as equipment, pipelines, tanks, vats, filters, evaporators, pasteurizers, and aseptic equipment in dairies, breweries, wineries, beverage and food processing/packing plants, egg processing/packing equipment surfaces, and eating establishments. This product is effective as a sanitizer when solution is prepared in water of up to 400 ppm hardness as CaCO<sub>3</sub>. This product has demonstrated greater than 99.999% reduction of organisms after 60 seconds exposure period in the Association of Official

Analytical Chemists (AOAC) Germicidal and Detergent Sanitizing Action of Disinfectants study.

NOTE: FOR MECHANICAL OPERATIONS prepared use solution may not be reused for sanitizing but may be reused for other purposes such as cleaning.
FOR MANUAL OPERATIONS fresh sanitizing solutions must be prepared daily or more often if the solution becomes diluted or solied.

Foam Cleaning of Food and Non-Food Contact Surfaces: As an adjunct to learning and santizing procedures this santizer/disinfectant may be added to PERAFOAM™ and foamed on environmental or equipment surfaces using conventional foam-generating equipment. PERAFOAM™ is the only approved product that may be used. The resultant foam blend can be used on equipment, proof, walls, ceilings, drains, etc., and must be left on surface for a minimum of 1 minute or long.

Food Contact Surface Directions for Mixing: Manually or mechanically blend no more than 1-6.1 fl. oz. of this product and 6-12 fl. oz. of PERAFOAM™ (foam additive) per 6 qallons of water. The dilution water must not exceed 150° F. Higher concentrations of this product and/or PERAFOAM™ may be used on food contact surfaces, but a potable water rinse is required. When used in organic production, a notable water rinse is required

Non-Food Contact Surface Directions for Mixing: Manually or mechanically blend 1-12 fl. oz. of this product and 6-36 fl. oz. of PERAFOAM™ (foam additive) per 6 gallons of water. The dilution water must not exceed 150° F. When used in organic production, a potable water rinse is required. Note: When using a foam additive, PERAFOAM™ is the only approved Drain Cleaning and Sanitizing: For use in open or closed drains such as in food, beverage, dairy, pharma and health care industries. Manually or mechanically blend 4-12 fl. oz. of this product with 1-12 ft, oz. of PERAFOAM (foam additive) per gallon of water and foam surfaces thoroughly using conventional foam-generating equipment. The dilution water must not exceed 150° F. Allow product to contact the surface slime/ biofilm for at least 10 minutes or more. A water rinse is optional. When used in organic production, a potable water rinse is required.

Entryway Sanitizing Systems: To help prevent cross-contamination from treated area to treated area, apply (spray) a sanitizing foam to the entryway. The foam must cover the entire path of the doorway. For effective coverage of footwear and forklift tires, etc., apply a foam layer 0.5-2 inches in depth. Set the system to deliver 1-6.1 fl. oz. (82-500 ppm active PAA) of this product and 3-12 fl. oz. of PERAFOAM™ (foam additive) per 6 gallons of water. Adjust the PAA concentration by testing the collapsed foam solution using a peroxyacetic acid test ki Note: When using a foam additive, PERAFOAM™ is the only approved product that may be used.

Alkaline Detergent Cleaning Adjunct (Booster) to Clean Food Processing Equipment: This product is an effective cleaning booster (hypochlorite alternative) for use with alkaline detergents. It may be used as a cleaning additive for Clean-In-Place (CIP) operations involving the circulation cleaning of pipelines, tanks, vessels, evaporators, HTSTs, and other food processing equipment. For cleaning applications as a detergent booster, use 1-6 oz. per gallon of water, to assist in the removal of organic soils. All hard nonporous food contact surfaces treated with this boosted detergent

must be thoroughly rinsed with potable water followed by sanitizing with an approved food contact surface sanitizer (such as this product). NON FOOD CONTACT HARD SURFACE DISINFECTION

mbination Disinfection and Cleaning: This product disinfects as it cleans in one operation. This product can be used to disinfect floors, walls and other hard nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, bed frames, shelves, racks, carts, refrigerators, coolers, tile, linoleum, vinyl, glazed porcelain, and use sites on this label made of plastic, stainless steel, or glass. For areas of use in hospitals, use this product for surgical and obstetrical suites, housekeeping services, physical therapy departments, nursing services, autopsy facilities. Also use this product in nursing homes, other health-care facilities, schools, colleges, veterinary clinics, animal life science laboratories, industrial facilities, dietary areas, office buildings, recreational facilities, etail and wholesale establishments

This product is effective against Staphylococcus aureus, Salmonella enterica, Pseudomonas aeruginosa, Trichophyton mentagrophytes and Escherichia coli O157:H7 at 0.38%-3% v/v (2.5-20 oz. per 5 gal) in hard water (400 ppm as CaCO<sub>3</sub>) and 5% organic soil loading on hard nonporous surfaces. For heavily soiled areas a pre-cleaning step is required, followed by a potable water rinse. Apply solution whater two plant as oxocy, and or organisation and an apply solution whater two plants as oxocy, and or organisation and entrapped soil with a clean wet more, cloth, sponge, brush, etc... or by soaking or immersion so as to make a solution and or organisation and entrapped soil with a clean wet more, cloth, wet vacuum pickup, or by draining. Surfaces that may directly or indirectly contact food must be rinsed with potable water before operations resume. A rinse for non-food contact surfaces is optional. Prepare a fresh solution daily or when it becomes soiled or diluted.

DISINFECTION OF ANIMAL AND POULTRY PREMISES, TRUCKS, COOPS AND CRATES This product is designed for use in animal hospitals, animal laboratories, kennels, pet shops, zoos, pet animal quarters, poultry premises, poultry hatcheries, and livestock quarters. When used as directed, this product is specifically designed to disinfect, deodorize and clean inanimate, hard, surfaces such as wells, floors, sink toos, furniture, operating tables, kennel runs, cages and feeding equipment. In addition this product will deodorize those areas which are generally hard to keep smelling fresh, such as garbage storage areas, empty garbage bins and cans, and any other areas which are prone to odors caused by microorganisms.

Disinfection of Poultry Premises: For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use, Remove all poultry and feeds from premises, trucks, coops and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with a detergent and rinse with water. Saturate surfaces with a 0.38-1.25% v/v (2.5-8.0 fl oz. per 5gal) solution of this product for a period of 10 minutes. Thoroughly scrub treated feed racks troughs automatic feeders fountains and waters with a detergent and rinse with potable water before rause. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse. See your echnical representative for specific recommendations for all cleaning and rinsing requirements

Disinfection And Deodorizing Of Animal Housing Facilities (Barns, Kennels, Hutches, Etc.): Remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter from floors, walls and surfaces of barns, pens, stalts, chutes, and other facilities (Barns, Kennels, Hutches, Etc.): Remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter from floors, walls and surfaces of barns, pens, stalts, chutes, and other facilities and fixtures occupied for the read of the remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter from floors, walls and surfaces of barns, pens, stalts, chutes, and other feed goupment. Throughly clean all surfaces with scap or detergent and rinse with water. Start and the remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter from floors, walls and surfaces of barns, pens, stalt, chutes, and other feed goupment. Throughly clean all surfaces with scap or detergent and the remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter from floors, and Deodorization of the remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter from floors and Deodorization of the remove animals and the remove animals and feed from premises, vehicles, and enclosures. Remove floors are found to the remove animals and feed from premises, vehicles, and enclosures. Remove floors are found to the remove animals and feed from premises, vehicles, and enclosures. Remove floors are found to the remove animals and feed from premises, vehicles, and enclosures. Remove floors are found to the remove floors and feed from premises and feed surfaces and allow to remain visibly wet for 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure. Ventilate build fedges, and other types of equipment used for removing litter and manure. Ventilate build fedges, and other types of equipment used for expensive manural manural scrapes used for removing litter and manure. Ventilate build fedges, fourthings and other other with scap or detergent, or uping the property of the scrapes of the sc CONTROL OF SLIME FORMING BACTERIA IN RECIRCULATING AND COOLING WATER SYSTEMS (COOLING TOWERS, EVAPORATIVE CONDENSERS, PASTEURIZERS

everely fouled systems must be cleaned before adding this product. This product must be added in the system directly and not mixed with any other chemicals or additives

Discontinue the use of chlorine or bromine products prior to using this product. Contamination with other chemicals could result in product decomposition. Add this product at a point in the system where uniform mixing and even distribution will occur

For slug treatment add 20 oz, of product per 1000 gallons of process water. Repeat as necessary until microbiological control is evident. Thereafter, to maintain control, use 0.3 to 1.5 lbs. (4.0-17.5 fl. oz.) of this product per 1000 gallons of process water (2-9 ppm active peroxyacetic acid) as a continuous or intermittent slug treatment. Continuous dosing methods usually require 2-5 ppm active xyacetic acid (4.0-10.2 fl. oz. per 1000 gal of process water) to achieve adequate control.

CLEANING: To remove sessile bacteria from cooling systems it is necessary to clean slime and slime-forming bacteria from the surfaces of all areas of water contact. This can be accomplished by treating the recycled water with 7.5-22.4 lbs. (102-306 fl oz.) of this product per 1000 gal of water (50-150 ppm active peroxyacetic acid) for 4-8 hours during normal tower operating cycles. This procedure can be used for online or offline cleaning. When finished bleed down the system until the PAA level is <5.10 ppm, then normal chlorine or bromine or PAA treatments can begin. This reatment must be done at least once or twice each year depending on exposure conditions.

Air Washers: This product may be used to control bacteria and biofouling in industrial air washing/scrubbing systems. The air washer must have operational and effective mist elimination systems. Prior to use of this product, heavily fouled systems must be pre-cleaned using the appropriate cleaner. Continuous dosing methods will require 2-7 ppm and intermittent dosing methods require 7-14

ppm (as peroxyacetic acid), as described in the previous paragraph, depending on the type of system and the level of microbiological control desired.

Evaporated or Condensed Water: This product may be used to treat SWEET or COW water (e.g. condensate of whey) collected from evaporated or condensing water systems in food or dairy Typically, the dosing regime would be using intermittent or continuous methods at 2-14 ppm as peroxyacetic acid.

REVERSE OSMOSIS (RO), ULTRA FILTRATION (UF) AND OTHER MEMBRANE CLEANING-SANITIZING his product may be used in the sanitization of ultra filtration (UF) and reverse osmosis (RO) membranes and other similar type membranes and their associated piping systems. This product may be added continuously in food, beverage, and drinking water systems for RO (reverse osmosis) systems only and in accordance with the instructions below. This product is not for use in kidney dialysis equipment. This product may not totally eliminate all vegetative microorganisms in RO or NF or UF membranes and their associated piping systems due to their construction or assembly but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed. Prior to using this product check with membrane manufacturer to confirm compatibility membranes with various types or concentration of peroxyacetic acid solutions.

Batch Sanitation of NF, UF and RO Systems: Isolate incompatible equipment, such as carbon filters and ion exchangers. Clean system with an appropriate cleaner and follow with RO permeate water or notable water Remove mineral denosits if necessary with an acidic cleaner, and rinse as before. Fill entire system with water and add up to 1% of this product by volume (620 ppm peroxyacetic acid) for heavily fouled systems. The typical sanitation use solution dosing of this product is 1-2 oz. per 5 gallons of water (98-195 ppm peroxyacetic acid). Recirculate the sanitizing solution through the piping and membrane system at 20° C for 10 minutes minimum, or up to 4 hours, depending on the severity of cleaning to be done. Open and close process valves and solenoids to be sure all parts are in contact with the solution. For occasional intermittent feed, do not exceed 98 ppm active peroxyacetic acid, which equals 1 oz, of this product per 5 gallons of feed water. Do not use the intermittent feed method for on-line use for potable water or direct food contact systems. Rinse the system with RO permeate or potable water until residual peroxygen concentration

RO Continuous or Intermittent Addition: For continuous addition methods for RO systems, use 2-5 ppm active peroxyacetic acid (36-90 ppm as product), which equals 1.8-4.5 oz. of this product per 430 gallons of process water. For occasional intermittent feed, do not exceed 98 ppm active peroxyacetic acid, which equals 1 oz. of this product per 5 gallons of feed water. Do not use the

TREATMENT OF FRUIT AND VEGETABLE PROCESS WATER SYSTEMS This product can be used in water or ice that contacts raw or fresh, post-harvest or further processed fruits and vegetables (in accordance with FCN 1738) for the control of spoilage and decay

causing bacteria and fungi in commercial operations and packinghouses. Batch, Continuous or Spray System Processes: Fill vessel containing fruits and vegetables with known amount of water. Ensure that water is circulating in vessel if using the submersion method.

Add this product to no more than 500 ppm residual peroxyacetic acid to the use solution in accordance with Food Contact Notification #1738, effective March 28, 2017. This can be accomplished by initially adding 10.0 fl. oz. per 10 gallons of water. The recommended concentration is between 30-300 ppm as peroxyacetic acid (0.60-6.0 fl. oz. per 10 gallons of water). The final concentration necessary to accomplish the intended task will vary from plant-to-plant. The fruits and vegetables can be continuously sprayed or submerged (dipped) in the resulting solution. Periodic or continuous additions of this product to maintain the required concentration may be added as necessary. It is also recommended to apply this product during the washing, chilling, or physical cleaning processes including the roller-spreader, washer or brush washer manifold, dip tank, or sorting processes. Contact time of 60 seconds is recommended to insure efficacy. A potable water rinse is not required. POULTRY, SWINE, LIVESTOCK WATERING OPERATING SYSTEMS

After watering lines have been cleaned, use this product at 0.8-1.1 fl. oz. per 100 gallons of water (4-5.7 ppm as peroxyacetic acid) to control algae and bacteria in drinking water and to control mineral build up in watering lines. Stop the use of this product twenty-four (24) hours prior to vaccination via the water line.

CLEANING POULTRY AND LIVESTOCK DRINKING WATER LINES

For drinking water lines using holding tanks make a stock solution by one of the following methods:

• Drinking water lines 500 feet in length or less; mix 2.1 gallons (270 fl. oz.) of this product with 100 gallons of water.

Drinking water lines exceeding 500 feet in length: mix 4.2 gallons (540 fl. oz.) of this product with 200 gallons of water. Pump the stock solution, completely filling the drinking water lines.

If the drinking water lines are not supplied by water from holding tanks, prepare a stock solution by one of the following methods:

- Mix 1.0 gallon (132 fl. oz.) of this product with 49 gallons of water in a 50 gallon tank, pumping this solution into the water line, repeating the process as often as needed, until water line
- Fill the water line, using a proportioner, set to inject this product undiluted at a rate of 1:47.

After the waterline is filled with the stock solution, activate nipole drinkers to ensure contact with drinkers. Allow the stock solution to remain in the water lines for 24-48 hours. Flush lines with fresh

water until water is visibly clear. Always make a fresh stock solution before use. <u>AGRICULTURAL or HORTICULTURAL USES</u>

Upon soil contact, this product decomposes rapidly to oxygen, carbon dioxide and water. This product may be harmful to fish if exposed on a continuous basis at concentrations of 0.5 ppm or more of active peroxyacetic acid. Meter this product into pressurized pipes using a plastic or stainless steel injection/backflow device installed far enough upstream from the target equipment to ensure thorough mixing. If open pouring of this product is required pour product as close to the surface of the water as possible to reduce odor exposure. Spray lines, hoses and tank must be clean before using this product. Make sure no iron or yellow metals are in contact with the spray solution at any time. Only stainless steel or plastic contact materials may be used in your spray rig

Treatment of Irrigation Water Systems (sand filters, humidification systems, storage tanks, ponds, reservoirs, canals, drip and sprinkler systems) (Not for Use in New York): For the control of odor, sulfides, non-pathogenic bacteria, slime and algae in water systems, apply this product at 0.4-2 oz. per 100 gal of water (2-10 ppm peroxyacetic acid). This feed rate equals 0.31-1.6 gal per 10,000 gallons of water. Repeat dose as necessary to maintain control, which will vary with seasonal conditions. For prevention of algae some systems may require continuous low

level dosing during warm sunny periods. Drip Irrigation System Cleaning: To clean slime and algae from drip system tapes and emitters, meter this product upstream from pumps or filters at the rate of 1-2 oz. per 50 gallons of water (10-20 ppm peroxyacetic acid). This feed rate equals 1.5-3 gal per 10,000 gallons of dilution water. When required, during normal irrigation cycles, use this product at the recommended dose for a

minimum of 30 minutes. After an irrigation cycle do not flush the lines. Spray Tank Treatment for Agricultural Water: In accordance with the Food Safety Modernization Act (FSMA), agricultural water applied to a growing food crop must be treated to within the microbial water quality profile (MWQP). For the control of odor, sulfides, non-pathogenic bacteria, slime and algae add 2 to 10 ppm PAA (0.4-2 fl. oz per 100 gal) to each spray tank of agricultural water to achieve hygienic conditions. These waters include municipal water, ground, well water or surface waters [rivers, streams, canals, lakes, ponds].

houses: This product can be used to suppress/control algae and slime formations in and around greenhouses. For normal use in various process, irrigation or sprinkler watering systems, this product may be used at 1:15,000 to 1:1,900 dilutions (4-33 ppm as peroxyacetic Heavily fouled systems, such as evaporative coolers or irrigation/drip lines, may need shock doses of up to 100 ppm as peroxyacetic acid (1.630 dilution). This product at its use dilution is compatible with stainless steel and aluminum surfaces. If product is intended to be used on any other surface, it is recommended that you apply product to a

small test area to determine compatibility before proceeding with its use. Pre-Plant Dip Treatment: Use this product for the control of damping-off, root disease and stem rot disease caused by Pythium (root rot) – Phytophthora (blights, rots) – Rhizoctonia (blight, stem rot) Fusarium (root-rot, leaf spot, Pink Snow Mold) – Thielaviopsis (black root rot), on seeds, seedlings, bulbs, or cuttings. Remove dead or dying foliage prior to dipping.

Use 20 fl. oz. per 50 gallons of water.

Immerse plants or cuttings; remove and allow to drain. Do not rinse.

Excessive foaming or bubbling during the dipping process is an indication of high levels of disease contamination.

Seed Treatment: Use this product for the control of damping-off, root disease and stem rot disease caused by Pythium (root rot) - Phytophthora (blights, rots) - Rhizoctonia (blight, stem rot)

Fusarium (root-rot, leaf spot, Pink Snow Mold) - Thielaviopsis (black root rot), on seeds of seed sprout crops such as mung bean, red clover, soybeans and alfalfa, and on crops grown exclusively for seed for planting.

. Use 20 fl. oz. per 50 gallons of water.

Immerse seeds and let soak for two minutes; remove and allow to drain. Do not rinse. Plant seed according to seed package directions

Soil Applications: This product is effective in the control of the following soil-borne plant pathogens: Fusarium (root-rot, leaf spot, Pink Snow Mold) – Phytophthora (blights, rots) – Pythium (root Rhizoctonia (blight, stem rot) - Verticillium (wilt).

se this product as a direct soil treatment, as a pre-plant application, at seeding or transplanting, and as a periodic soil treatment throughout the plant's life up to the day of harvest. The performance of this product is not affected by furnigation. Use this product on furnigated and unfurnigated soil.

Soil Treatment Prior to Seeding or Transplanting: Cultivate the soil prior to treatment. Break-up compacted soil and clods to loosen soil completely. Mix 102 fl. oz. of this product per 100 callons of

water to yield approximately 500 ppm PAA. Make banded or broadcast applications of 25 to 100 gallons of solution per acre-row either prior to planting or at the time of planting. This product will not harm seedlings or plants when applied at labeled rates. In fields with a history of disease pressure, use the 100 gallons of mixed solution per acre-row rate

Soil Treatment with Established Plants or Seedlings: Apply this product at any stage of plant growth as a soil treatment up to the day of harvest. Make applications using soil drench, flood or drip irrigation. Ensure that soil moisture of the beds is at or near field capacity prior to application.

Soil Drench: Apply 20-41 fil. oz. of this product per 200 gallons of water per 1,000 square feet of soil to be treated to yield approximately 50-100 ppm PAA.

Flood Irrigation: Inject this product through a metered system using 100 fil. oz. of this product per 1,000 gallons of water used to yield approximately 50 ppm PAA.

Drip Irrigation: Apply this product through the drip tape at a rate of 10.7-24 fl. oz. per 1,000 feet of row. Inject this product through a metered system using 100 fl. oz. of this product per 1,000 gallons of water used to yield approximately 50 ppm PAA. Apply first treatment during the first drip irrigation cycle. Apply two additional treatments at 7-14 day intervals. Under severe disease conditions apply at 7-day intervals using the highest rate. Under severe disease conditions and during periods of rainy weather, apply this product immediately following rain to suppress the spread of disease and help oxygenate the soil. Combine lower rates of this product with other nonmetal based fungicides.

Row Center Spacing	Rate of This Product	Application Instructions
5.5-6 ft.	100-134 fl. oz. per acre	Apply through irrigation systems using a 45 to 90 minute run time     Repeat applications at 7 to 14 day intervals

Foliar Applications: This product can be applied to the following growing crops to control fungi. Crops: root vegetables, potatoes, berries, strawberries, citrus fruit, pome fruit, stone fruit, herbs, spices, peppers, tomatoes, eggplant, sweet potatoes, bulbs, onions, cucurbits, cucumbers, tropical fruits, avocadoes, bananas, mangoes, grapes, brassicas, peas, beans, soybeans, cereal crops, rice, wheat, peanuts, alfalfa, chinese vegetables, greens, lettuce, leafy greens, celery, apiaceaes, cranberries, legumes, corn (field, sweet, seed), wild rice, cole crops, garlic, leeks, green onions, mushrooms sugar beets tobacco hops grass for seed or sod asparagus nuts valnuts nistachios macadamia nuts almonds cotton and coffee

o suppress/control/prevent the following non-human plant pathogens: Alternaria, Angular leaf spot, Anthracnose, Bacterial blotch, Bacterial speck, Bacterial spot, Black rot, Blights, Blue mold, Botrytis, Brown rot, Citrus canker, Cladosporium, Crown rot, Downey mildew, Early blight, Fruit rot, Fusarium, Gray leaf spot, Gummy stem blight, Leaf blight, Leaf rust, Leaf spot, Mycogene, Necrotic spot, Phytophthora, Potato brown rot, Powdery mildew, Pythium, Rhizoctonia, Rust, Scab, Sclerotina, Shot hole, Sooty mold, Stem rot, Trichoderma, Verticillum, White mold,

Initial Curative Application:

I. Use 3.4-6.8 fl. oz. of this product per 5 gallons of clean water.

Do not reuse already mixed solution; make fresh daily. Spray or mist plants and trees.

Thoroughly wet all surfaces of plant, upper and lower foliage, including stems, branches, and stalks to ensure full contact with plant tissue 1. Based on the disease severity, apply for one to three consecutive days and then follow directions for preventative treatment after the initial application.

kly Preventative Treatment

. Use 0.66-1.1 fl. oz. of this product per 5 gallons of clean water.

Spray or mist plants and trees.

oroughly wet all surfaces of plant, upper and lower foliage, including stems, branches, and stalks to ensure full contact with plant tissue.

Based on the disease pressure, spray every five to seven days as a preventative treatment 5. At the first sign of disease, spray daily with 3.4-6.8 ft. oz. of this product per 5 gallons of water for three consecutive days and then resume weekly preventative treatment. Apply solution at 50-100 gallons per treated acre, depending on spray method used.

Note: 1 fl. oz. of this product per 5 gallons of water = 100 ppm PAA A nonionic spreader (surfactant) adjuvant should be used for better results. Contact your local supplier or farm supply.

# STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage or disposal.

Storage: Never return this product to the original container after it has been removed. Avoid all contaminants, especially dirt, caustic, reducing agents, and metals. Contamination and impurities will reduce shelf life and can induce decomposition. In case of a decomposition, isolate container, spray container with cool water and dilute this product with large volumes of water. Avoid damage to containers. Keep container closed at all times when not in use. Keep container out of direct sunlight. To maintain product quality, store at temperatures below 86°F. Procedure to Leak or Spill: Stop leak if this can be done without risk. Shut off ignition sources: no flames, smoking, flares, or spark producing tools. Keep combustible and organic materials away. Flus pilled material with large quantities of water. Undiluted material must not enter confined spaces.

esticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste representative at the nearest EPA Regional Office for guidance. I naterial has been spilled, an acceptable method of disposal is to dilute with at least 20 volumes of water followed by discharge into suitable treatment system in accordance with all local, state and Federal environmental laws rules regulations standards and other regulirements. Recause acceptable methods of disposal may vary by location, regulatory agencies must be contacted prior to disposal. This product which is to be discarded, must be disposed of as hazardous waste after contacting the appropriate local state of Federal agency to determine proper procedures Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Offer for recycling if available. Triple rinse as follows: Empty the emaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring a least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate or later use or disposal. Repeat this procedure two more times.

Container Handling: (Containers equal to or less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two nore times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances

Lot #:



Manufactured for: J.R. Simplot Company,

P.O. Box 27, Boise, ID 83707 • (208) 336-2110 IN CASE OF EMERGENCY CALL CHEMTREC 1-800-424-9300

Net Contents:

DOT: UN3098, Oxidizing liquid, corrosive, n.o.s. (contains hydrogen peroxide and peroxyacetic acid mixture, stabilized), 5.1(8), PG II

ITEM#80031 19323 20210218