



WEED & FEED

25-3-5

ACTIVE INGREDIENT:

2,4-dichlorophenoxyacetic acid.....0.69%
 (+)-(R)-2-(4-chloro-2 methylphenoxy)
 propionic acid0.15%
 Dicamba: 3,6-dichloro-2-methoxybenzoic
 acid0.06%

OTHER INGREDIENTS.....99.10%
TOTAL100.00%

THIS PRODUCT CONTAINS:

13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
 (CAS No. 94-75-7)
 3.0 lbs. (+)-(R)-2-(4-chloro-2 methylphenoxy) propionic acid
 equivalent per ton. (CAS No. 16484-77-8)
 1.2 lbs. 3,6-dichloro-2-methoxybenzoic acid equivalent per
 ton. (CAS No. 1918-00-9)

Isomer Specific by AOAC Methods.

GUARANTEED ANALYSIS:

Total Nitrogen (N) 25.00%
 9.6% Ammoniacal Nitrogen
 15.4% Urea Nitrogen
 Available Phosphate (P₂O₅) 3.00%
 Soluble Potash (K₂O) 5.00%

Derived From: Urea, Muriate of Potash,
 Ammonium Phosphate, and Ammonium Sulfate.

EPA Reg. No. 2217-579-2935
 EPA Est. No. 69759-OR-4

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes and socks, and chemical-resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect nontarget plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. Do not use on dichondra, carpetgrass, or St. Augustinegrass. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing. Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, because injury may result. Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

WEED & FEED 25-3-5 contains Trimec® Herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many other listed weeds commonly found in home lawns. In addition, a 25-3-5 fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made 30 days after the initial treatment. Treated areas may be seeded 3 to 4 weeks after application.

The maximum broadcast application rate is 4 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 pounds of product per 1,000 square feet (2.4 lb. 2,4-D ae, 0.54 lb. MCP-p ae and 0.22 lb. Dicamba ae per acre).

Mow lawn to normal height 1 to 2 days before application. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

WEEDS CONTROLLED			
Bedstraw, Catchweed	Dock, smooth	Plantain, bracted	Speedwell, snow
Bedstraw, smooth	Ground Ivy	Plantain, broadleaf	Speedwell, thymeleaf
Bedstraw, Northern	Healall	Plantain, buckhorn	Spurge, spotted
Bedstraw, pineywoods	Henbit	Plantain, hoary	Spurge, prostrate
Black medic	Knotweed, prostrate	Plantain, narrowleaf	Thistle, blessed
Burdock	Lambsquarters	Plantain, slender	Thistle, bull
Chickweed, common	Lespedeza	Plantain, woolly	Thistle, Canada
Chickweed, mousear	Mallow, alkali	Poison ivy	Thistle, flodman (prairie)
Chickweed, sticky	Mallow, bristly	Poison oak	Thistle, musk
Chicory	Mallow, common	Purslane, common	Thistle, platte
Clover, alyce	Mallow, dwarf	Ragweed, bur	Thistle, plumeless
Clover, crimson	Mallow, Venice	Ragweed, common	Thistle, Russian
Clover, hop	Morningglory, bigroot	Ragweed, lanceleaf	Thistle, scotch
Clover, rabbit foot	Morningglory, ivyleaf	Ragweed, western	Thistle, wavyleaf
Clover, red	Pepperweed, perennial	Sorrel, red (sheep)	Thistle, yellow (horrible)
Clover, strawberry	Pepperweed, Virginia (Peppergrass)	Shepherdspurse	Wild Carrot
Clover, yellow sweet	Pigweed, prostrate	Speedwell, bi-lobed	Wild garlic
Clover, white sweet	Pigweed, redroot	Speedwell, common	Wild lettuce
Clover, white	Pigweed, Russian	Speedwell, corn	Wild onion
Dandelion, common	Pigweed, smooth	Speedwell, ivyleaf	Yarrow, common
Dock, broadleaf	Pigweed, tumble	Speedwell, Persian	Yarrow, western
Dock, curly	Plantain, blackseed	Speedwell, purslane	Yarrow, yellow
		Speedwell, slender	

SPREADERS AND CALIBRATION:

1. Drop spreaders and rotary spreaders must be calibrated for accurate application of this product.
2. Close spreader opening (orifice).
3. Adjust spreader setting using the Spreader Suggestions chart.

**Application Rate = Apply 4 lb. of WEED & FEED 25-3-5 per 1,000 square feet or 175 lb. per acre.
(1 lb. of nitrogen per 1,000 square feet.)**

CENTRIFUGAL/ROTARY SPREADERS

Frank's Broadcast	2¼
Scotts® Speedy Green.....	5
Radio Flyer #0047.....	8¼
True Temper Rotary CB2000.....	3¾
Earthway Rotary #2050SU.....	14½

DROP SPREADERS

Frank's Drop.....	8¼
Greenview II.....	5½
True Temper CD20.....	4-4½
Scotts Accugreen AG1.....	5½
Earthway Drop #7300SU.....	13½

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

4. Calibration Instructions: For 4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
6. Apply the product to lawn.
7. Close the spreader openings when turning to change directions or when stopping.
8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

This 50-pound bag covers approximately 12,500 square feet (100 ft. x 125 ft.)

This 20-pound bag covers approximately 5,000 square feet (50 ft x 100 ft.)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If Empty:** Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS® logo and Ideas to Grow With® are registered trademarks of Wilbur-Ellis Company LLC.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

Scotts® is a registered trademark of The Scotts Company.

For chemical spills, leaks, fire or exposure, call CHEMTREC: (800) 424-9300.

NET CONTENTS: 50 POUNDS

F-012219

DISTRIBUTED BY:

Wilbur-Ellis Company LLC
PO Box 16458
Fresno, CA 93755

GUARANTEED BY:

Wilbur-Ellis Company LLC.
7 E. Washington Ave.,
Yakima, WA 98903

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

