



YaraVita™

Citrel™

NOTICE

Before using this product, read the entire label

Citrel™

3-0-0

A concentrated liquid fertilizer with micronutrient.

F1673

Guaranteed analysis:

Total Nitrogen (N)	3%
3% Urea Nitrogen	
Magnesium (Mg)	1.8%
0.001% water soluble Magnesium (Mg)	
Manganese (Mn)	12.3%
0.001% water soluble Manganese (Mn)	
Zinc (Zn)	15.3%
0.001% water soluble Zinc (Zn)	

Derived from Urea, magnesium hydroxide, manganese carbonate, zinc oxide.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

Contents made in Great Britain

Guaranteed & Manufactured by Yara UK Ltd., Manor Place, Wellington Road, The Industrial Estate, Pocklington, York YO42 1DN, U.K.


Manufactured for Yara North America, Inc. 100 North Tampa Street, Suite 3200, Tampa, FL 33602 Customer Service: 800-234-9376

Warning

H411: Toxic to aquatic life with long lasting effects.

P273: Avoid release to the environment.

P391: Collect spillage.



**2½ gallons/
9.4 litres**

**Net weight
34 lb/15.4kg**

UN 3082
ENVIRONMENTALLY
HAZARDOUS
SUBSTANCE, LIQUID,
N.O.S. (zinc oxide)

YaraVita Citrel™

APPLICATION RATES AND TIMINGS

Citrus: 1 to 2 qts/ac applied during spring flush and also during summer / autumn flush. In both instances a repeat application 10 to 14 days later may be required where moderate to severe deficiency exists.

Water rate: 50 to 100 gallons/acre.

Melon: Field Grown: 1 qt/ac from the 4 to 6 leaf stage, repeated at 10 to 14 day intervals if necessary. N.B. Final application to be made at least one month before harvest. Water rate: 20 to 50 gallons/acre.

Strawberry: (field grown): 1 pint to 1 quart/acre at spring regrowth and also at green bud.

Water rate: 3 to 20 gallons/acre.

Tomato: Field Grown: 1 qt/ac applied from the 4 to 6 leaf stage onwards. Repeat after a 10 to 14 day interval if required. N.B. Final application to be made at least one month before harvest.

Water rate: 20 to 50 gallons/acre.

Extended recommendations available at <https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and reproductive Harm - www.P65Warnings.ca.gov