




YaraVita™

Photrel™ Advance SC



2½ gallons/
9.4 litres

Net weight
31lb 13oz/
14,4 kg

NOTICE

Before using this product, read the entire label

Safety data sheet available for professional user on request

Photrel™ Advance SC

4-0-0

A formulated multinutrient product for foliar application

Guaranteed analysis:

Total Nitrogen (N)	4%
4% Urea Nitrogen	
Calcium (Ca)	5.8%
Magnesium (Mg)	4.6%
0.001% water soluble Magnesium (Mg)	
Boron (B)	3.9%
Manganese (Mn)	4.6%
0.001% water soluble Manganese (Mn)	
Molybdenum (Mo)	0.3%

Derived from Urea, Calcium Borate, Magnesium Hydroxide, Manganese Carbonate, Sodium Molybdate

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.com/metals.html>

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

F1673

Contents made in Great Britain

Guaranteed & Manufactured by Yara UK Ltd., Manor Place, Wellington Road, The Industrial Estate, Pocklington, York YO42 1DN, U.K.

Manufactured for Yara North America, Inc. 100 North Tampa Street, Suite 3200, Tampa, FL 33602 Customer Service: 800-234-9376

Warning

H361: Suspected of damaging fertility or the unborn child.

P202: Do not handle until all safety precautions have been read and understood.

P280: Wear protective gloves/clothing and eye/face protection.

P308+P313: IF exposed or concerned: Get medical attention.

P501: Dispose of contents and container according to local regulations.



YaraVita™ Photrel™ Advance SC

WARNING

This product contains boron which may be injurious to certain crops. Contact your county agent or field consultant for specific information.

This product is to be used only on crops which respond to molybdenum (Mo). Crops high in molybdenum (Mo) are toxic to grazing animals (ruminants).

APPLICATION RATES AND TIMINGS

Beans, Peas: 1-2 quarts per acre as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. N.B. Do not spray during flowering. Water rate: 20 gallons/acre.

Brassicas: 1-2 quarts/acre as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest.

Water rate: 20 gallon/acre.

Canola: 1 to 2 qts/acre at the 4 to 9 leaf stage and again at the onset of the rapid stem extension phase up to the start of flowering. Repeat at 10 to 14 day intervals within these growth stages as necessary. Water rate: 5 to 20 gallons/acre.

Carrot: Two applications of 1-2 pints/acre applied when the crop is 6 inches tall and again 2 weeks later. Water rate: 20 gallons acre.

Cereals, Wheat: 1.5 qts/acre applied at any time from the 2 leaf stage to 1st node detectable. Water rate: 20 gallons/acre.

Legumes/Pulses: 1-2 quarts/acre as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. N.B. Do not spray during flowering. Water rate: 20 gallons/acre.

Lettuce (Field Grown): 1-2 quarts/acre as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. Water rate: 20 gallons/acre.

Onion: 2 applications of 1-2 pints/acre applied as soon as there is adequate leaf cover (6 leaves) and repeated 2 weeks later. Water rate: 20 gallons/acre.

Potatoes: 1.5 qts./acre applied one week after 100 % emergence with a second application 10 to 14 days later. Water rate: 20 gallons/acre.

DRY FERTILIZER IMPREGNATION

This product may be impregnated onto dry fertilizer granules. N.B. Must not be used with a straight ammonium nitrate fertilizer, or any fertilizers containing ammonium nitrate which may have self-sustaining decomposition properties. Consult your local Yara North America technical representative for details.

Extended recommendations available at

<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

NOTICE: This product may contain a chemical (or chemicals) known to the State of California to cause cancer and birth defects or other reproductive harm.

DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.