




YaraVita™

Phoztrac™



2½ gallons/
9.4 litres

Net weight
32lb 10oz/
14.8 kg

NOTICE

Before using this product, read the entire label

Phoztrac™

4-15-0

A formulated product for the treatment of phosphorus, calcium and zinc deficiencies by foliar application

Guaranteed analysis:

Total Nitrogen (N).....4%
4% Urea Nitrogen

Available Phosphate (P₂O₅).....15%

Calcium (Ca).....13%

Zinc (Zn).....7.7%

Derived from Urea, Calcium Phosphate, Calcium Carbonate, Zinc Phosphate

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

F1673

Contents made in Great Britain

Guaranteed & Manufactured by Yara UK Ltd., Manor Place, Wellington Road, The Industrial Estate, Pocklington, York YO42 1DN, U.K.

Manufactured for Yara North America, Inc. 100 North Tampa Street, Suite 3200, Tampa, FL 33602 Customer Service: 800-234-9376

Warning

H400: Very toxic to aquatic life.

H410: Very toxic to aquatic life with long lasting effects.



P273: Avoid release to the environment.

P391: Collect spillage.

THIS PRODUCT IN NON-BULK PACKAGING IS A MARINE POLLUTANT WHEN TRANSPORTED BY VESSEL. HOWEVER, IT IS NOT DOT REGULATED WHEN TRANSPORTED BY HIGHWAY ONLY PER 49 CFR, 171.4(c)

UN 3082

ENVIRONMENTALLY
HAZARDOUS
SUBSTANCE,
LIQUID,
N.O.S.
(zinc bis(orthophosphate))



YaraVita™ Phoztrac™

APPLICATION RATES AND TIMINGS

Apples, Pears: 2 to 4 quarts/acre commencing at petal fall with repeat applications at 10 to 14 day intervals. On russet sensitive varieties delay applications until 6 weeks after petal fall.

Water rate: 50 gallons/acre.

Apricots, Cherries, Nectarines, Peaches, Plums: 2-4 quarts/acre applied 2 to 5 times. First application at pink bud. Thereafter at petal fall with repeat applications at 7 to 14 day intervals. Water rate: 50 to 100 gallons/acre

Cole Crops, Broccoli, Brussel Sprouts, Cabbage, Calabrese,

Cauliflower: 2 to 4 quarts/acre applied 2 or 3 times at stem extension/head development with 10 to 14 day intervals.

Water rate: 50 gallons per acre.

Canola: 2 to 4 quarts/acre applied 2 to 3 times - at 4 to 6 leaf stage, onset of stem extension, and 10 to 14 days later.

Water rate: 25 gallons/acre.

Carrots: 1 to 2 quarts/acre applied 2 or 3 times when crop is 6 inches tall and at 7 to 10 day intervals.

Water rate: 25 gallons/acre.

Citrus: 2 to 4 quarts/acre applied 2 to 3 times starting at fruit set with repeat applications at 10 to 14 day intervals.

Water rate: 50 to 100 gallons/acre.

Conifers: 1 gallon/acre at the start of new season leaf production, and again in early autumn.

Water rate: 3 to 50 gallons/acre.

Corn: 1 to 2 quarts/acre at 4 to 8 leaf stage.

Water rate: 25 gallons/acre.

Cucurbits (Field Grown): 1 to 2 quarts/acre at start of flowering.

Repeat if necessary at 10 to 14 day intervals.

Water rate: 50 gallons/acre.

Garlic: 1-2 quarts/acre at the 6 inches stage, and again 10 to 14 days later. Water rate: 3 to 20 gallons/acre.

Nuts (Deciduous): 2-4 quarts/acre 7 days after petal fall. Repeat if necessary 10 to 14 days later.

Water rate: 50 gallons/acre.

Onions: 1 to 2 quarts/acre applied once or twice during bulb filling with a 10 to 14 day interval between sprays.

Water rate: 25 gallons/acre.

Peppers (Field Grown): 1-2 quarts/acre applied 2 to 4 times commencing from flowering on second truss. Repeat at 10 to 14 day intervals. Water rate: 3 to 50 gallons/acre.

Potatoes: 2-4 quarts/acre 1 week after 100% crop emergence.

Repeat as necessary at 10 to 14 day intervals.

Water rate: 3 to 25 gallons/acre.

Raspberry: 2-4 quarts/acre to new leaf growth in the spring. Water rate: 50 gallons/acre.

Sweet Potatoes: 1-2 quarts/acre one week after 100% emergence or transplanting. Repeat applications during tuber bulking at 10 to 14 day intervals. Also, apply at the same rate following recommendation from analysis.

Water rate: 20 gallons/acre.

Tomatoes (Field Grown): 1-2 quarts/acre applied 2 to 4 times commencing from flowering on second truss. Repeat at 10 to 14 day intervals. Water rate: 3 to 50 gallons/acre.

Vines: 2 to 4 quarts/acre when new shoots are at the 4 to 6 leaf stage. Repeat if necessary at 10 to 14 day intervals.

Water rate: 50 gallons/acre.

Extended recommendations available at

<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov