

Supplemental Label



For weed control in dry bulb onions and leek

This supplemental label expires December 31, 2022, and must not be used or distributed after this date.

Active Ingredient:

pyroxasulfone: 3-[[[5-(difluoromethoxy)-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl]methyl]sulfonyl]-4,5-dihydro-5,5-dimethylisoxazole	41.46%
Other Ingredients:	58.54%
Total:	100.00%

Contains 4.17 pounds of pyroxasulfone per gallon formulated as a water-based suspension concentrate

EPA Reg. No. 7969-374
KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Zidua® SC herbicide** container label, EPA Reg. No. 7969-374, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Zidua SC** before applying.
- Use of **Zidua SC** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Zidua SC**.

Product Information

Zidua SC is a selective rate-dependent residual preemergence herbicide for controlling annual grasses, sedges, and annual broadleaf weeds. Refer to the **Zidua SC** container label for a complete list of weeds controlled.

Application Information

Zidua SC may be applied to dry bulb onions (direct seeded and transplanted) and leek (transplanted only) as a postemergence application for residual

preemergence weed control. Use **Zidua SC** as part of a weed control program in dry bulb onions and leek either in combination or sequentially with other herbicides for a broader spectrum of weed control and/or control of emerged weeds. **Zidua SC** does not control emerged weeds.

Application Rate

Apply **Zidua SC** in dry bulb onions and leek at the residual rates provided in **Table 1**.

Table 1. Residual Rates of Zidua SC in dry bulb onions and leek

Application Timing	Use Rate by Soil Texture (fl ozs/A)		
	Coarse	Medium to Fine	Muck (greater than 20% organic matter)
Postemergence	DO NOT USE	2.0 to 2.75	2.75

Application Method and Timing

Zidua® SC herbicide may only be applied in a single application.

Postemergence Application

Apply **Zidua SC** at the use rates specified in **Table 1** as a broadcast spray to dry bulb onions and leek that have 2 to 6 true leaves.

Crop-specific Restrictions

- **DO NOT** apply more than 2.75 ozs/A of **Zidua SC** (0.09 lb ai/A of pyroxasulfone) in a single yearly application.
- **DO NOT** apply **Zidua SC** within 60 days of harvest of dry bulb onions or leek.
- **DO NOT** apply **Zidua SC** to directed-seeded leek.

Crop-specific Precautions

- **Crop response** - The use of **Zidua SC** may result in temporary growth suppression, leaf burn, and/or other injury or stand reduction to dry bulb onions or leek under stressful conditions including inadequate or excessive moisture, extended periods of water-saturated soil occur during early transplant growth and development, cool and hot temperatures, compacted soils, injury from other pesticides, disease or other pest damage, mechanical injury, nutrient imbalances, or other conditions known to cause plant stress.

Tank Mixes

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Zidua SC may be tank mixed or applied sequentially with other herbicide products registered for use in dry bulb onions or leek for a broader spectrum of control and/or control of emerged weeds. Refer to the tank mix product labels to confirm that the respective tank mix products are registered for use on dry bulb onions or leek. Follow the adjuvant directions for the tank mix partner of **Zidua SC**.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.


TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Zidua is a registered trademark of BASF.

The  symbol is a registered trademark of Kumiai Chemical Industry Co., Ltd.

© 2019 BASF Corporation
All rights reserved.

007969-00374.20191010b.NVA 2019-04-489-0217
Based on: NVA 2019-04-489-0213

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry