

## FIFRA 2(ee) Recommendation

EPA Reg. No.: 279-3618

## FOR DISTRIBUTION AND USE IN ALL REGISTERED STATES (EXCEPT TEXAS)

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation is valid until December 31, 2027 or until withdrawn, canceled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Off-the-Seed Placement of Zironar® Biofungicide/Bionematicide at-plant in Corn (Field, Corn Grown for Seed, Popcorn, Sweet Corn)

## **Directions for Use**

- Zironar<sup>®</sup> Biofungicide/Bionematicide may be applied at planting time to the soil in-furrow with the seed, or placed sub-surface off the row to the side or below the seed (i.e., 2x2 inch, 2x2x2 dual-placement, or various placements) within 3-inches of the seed.
- Zironar<sup>®</sup> Biofungicide/Bionematicide can be tank mixed with other products approved for 2x2, a 2x2x2 or similar dual placement. Always perform a jar test to confirm physical mixture compatibility.
- Refer to row space conversion chart and application rates on the EPA registered Zironar<sup>®</sup> Biofungicide/Bionematicide label.

R-4869 010824 10-28-22

