

SUPPLEMENTAL LABELING

FLUOXASTROBIN	GROUP	11	FUNGICIDE
TETRACONAZOLE	GROUP	3	FUNGICIDE

ZOLERA[®] FX FUNGICIDE

Supplemental Label to Add DIRECTIONS FOR USE on RAPESEED (CANOLA)

This supplemental label expires May 2, 2021 and must not be used or distributed after this date.

ACTIVE INGREDIENTS:	% BY WT.
Fluoxastrobin: [(1 <i>E</i>)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] (5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime].....	17.76%
Tetraconazole: 1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy)propyl]1 <i>H</i> -1,2,4-triazole	17.76%
OTHER INGREDIENTS:	64.48%
TOTAL:	100.00%

This product contains 1.67 pounds of Fluoxastrobin per gallon (200 g per liter).

This product contains 1.67 pounds of Tetraconazole per gallon (200 g per liter).

KEEP OUT OF REACH OF CHILDREN

CAUTION / PRECAUCIÓN

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **ZOLERA[®] FX FUNGICIDE** BEFORE APPLYING. USE OF **ZOLERA FX FUNGICIDE** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS, RESTRICTIONS, AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **ZOLERA FX FUNGICIDE (EPA Reg. No. 66330-424)**.

121818V003Su

RAPESEED (CANOLA)

(Crop Subgroup 20A: Borage; Crambe; Cuphea; Flax Seed; Gold of Pleasure; Hare’s Ear Mustard; Lesquerella; Lunaria; Meadowfoam; Milkweed; Mustard Seed; Oil Radish; Poppy Seed; Rapeseed; Sesame; Sweet Rock. Cultivars, varieties and/or hybrids of these crops.)

Disease Control	Rate (fl oz product/A) (lb ai/A)	Application Timing and Information
Alternaria Blackspot (<i>Alternaria</i> spp.) Blackleg (<i>Leptosphaeria maculans</i>) Suppression Sclerotinia Stem Rot (<i>Sclerotinia sclerotiorum</i>)	5.0 – 7.7 (0.07 - 0.10 fluoxastrobin + 0.07 - 0.10 tetraconazole)	<p>For optimum results, begin applications preventively and continue as needed on a 7- to 14-day interval. Use the higher rate and shorter interval when disease pressure is high.</p> <p>Specifically for blackleg, applications should be made at the 2 to 4 leaf stage.</p> <p>For Alternaria or Sclerotinia, applications should be made at 10 to 25% flowering (3 to 7 days following first flower).</p> <p>Apply by ground in a minimum of 10 GPA or by air in a minimum of 2 GPA. Use sufficient water to obtain thorough coverage of plants.</p>

RESTRICTIONS:

- Do not apply more than 7.7 fl oz (0.1 lb ai tetraconazole and 0.1 lb ai fluoxastrobin) of **ZOLERA FX FUNGICIDE** per acre per single application.
- Do not apply more than 15.4 fl oz (0.2 lb ai tetraconazole and 0.2 lb ai fluoxastrobin) of **ZOLERA FX FUNGICIDE** per acre per year.
- Do not apply more than 0.3 lb ai of fluoxastrobin per acre per year (total of all applications of fluoxastrobin products and application methods).
- Do not apply more than 0.2 lb ai of tetraconazole per acre per year (total of all applications of tetraconazole products and application methods).
- Do not make more than 2 applications of **ZOLERA FX FUNGICIDE** per year.
- Do not make applications of **ZOLERA FX FUNGICIDE** less than 7 days apart.
- Pre-Harvest Interval (PHI): Do not apply **ZOLERA FX FUNGICIDE** within 21 days of harvest.

Produced For:
ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. ZOLERA and the ZOLERA FX logo are registered trademarks of an Arysta LifeScience Group Company.
 ©2019 Arysta LifeScience Group Company.