

SUPPLEMENTAL LABELING

FLUOXASTROBIN	GROUP	11	FUNGICIDE
TETRACONAZOLE	GROUP	3	FUNGICIDE

ZOLERA[®] FX FUNGICIDE

Supplemental Label to Add DIRECTIONS FOR USE
on DRY PEAS and DRY BEANS (EXCEPT SOYBEAN)

This supplemental label expires May 2, 2021 and
must not be used or distributed after this date.

ACTIVE INGREDIENTS:	% BY WT.
Fluoxastrobin: [(1 <i>E</i>)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] (5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime].....	17.76%
Tetraconazole: 1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy)propyl]1 <i>H</i> -1,2,4-triazole	17.76%
OTHER INGREDIENTS:	64.48%
TOTAL:	100.00%

This product contains 1.67 pounds of Fluoxastrobin per gallon (200 g per liter).

This product contains 1.67 pounds of Tetraconazole per gallon (200 g per liter).

KEEP OUT OF REACH OF CHILDREN

CAUTION / PRECAUCIÓN

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **ZOLERA[®] FX FUNGICIDE** BEFORE APPLYING. USE OF **ZOLERA FX FUNGICIDE** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS, RESTRICTIONS, AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **ZOLERA FX FUNGICIDE (EPA Reg. No. 66330-424)**.

121818V003Su

DRY PEAS and DRY BEANS (EXCEPT SOYBEAN)

(Subgroup 6C: **Bean (*Lupinus Spp.*)** Grain Lupin, Sweet Lupin, White Lupin, and White Sweet Lupin); **Bean (*Phaseolus Spp.*)** (Includes Field Bean, Kidney Bean, Lima Bean (Dry), Navy Bean, Pinto Bean, Runner Bean, Snap Bean, Tepary Bean, Wax Bean); **Bean (*Vigna Spp.*)** (Adzuki Bean, Asparagus Bean, Blackeyed Pea, Cowpea, Catjang, Chinese Longbean, Crowder Pea, Moth Bean, Mung Bean, Rice Bean, Southern Pea, Urd Bean, Yardlong Bean; Immature Seed (Edamame); **Broad Bean** (Fava Bean - Dry) (*Vicia faba*); **Chickpea**; **Guar** (*Cyamopsis tetragonoloba*); **Lablab Bean** (Hyacinth Bean) (*Lablab purpureus*); **Lentil** (*Lens esculenta*); **Pea** (*Pisum spp.*) (Dwarf Pea, Ediblepod Pea, English Pea, Garden Pea, Green Pea, Field Pea, Snow Pea, Sugar Snap Pea); **Pigeon Pea** (*Cajanus cajan*).

Disease Control	Rate (fl oz product/A) (lb ai/A)	Application Timing and Information
Alternaria Blight (<i>Alternaria spp.</i>)	5.5 - 7.7	For optimum results, begin applications preventively and continue as needed on a 7- to14-day interval.
Alternaria Leaf Spot (<i>Alternaria alternata</i>)	(0.07 - 0.10 fluoxastrobin	For management of Ascochyta use the highest specified rate.
Anthrachnose (<i>Colletotrichum lindemuthianum</i>)	+ 0.07 - 0.10 tetraconazole)	To limit the potential for development of disease resistance follow the guidelines outlined in the RESISTANCE MANAGEMENT section.
Ascochyta Blight (<i>Mycosphaerella pinodes</i>)		An adjuvant may be added to the tank.
Ascochyta Leaf and Pod Spot (<i>Ascochyta spp.</i>)		Apply by ground in a minimum of 10 GPA or by air in a minimum of 2 GPA. Use sufficient water to obtain thorough coverage of plants.
Ascochyta Leaf Spot (<i>Ascochyta phaseolomm</i>)		
Bean Rust (<i>Uromyces appendiculatus</i>)		
Rust (<i>Phakopsora spp.</i>)		
Southern Blight (<i>Sclerotium rolfsii</i>)		
Web Blight (<i>Rhizoctonia solani</i>)		
Powdery Mildew of pea (<i>Erysiphe pisi</i>)		
Suppression		
Sclerotinia White Mold/ Stem Rot (<i>Sclerotinia sclerotiorum</i>)		

RESTRICTIONS:

- Do not apply more than 7.7 fl oz (0.1 lb ai tetraconazole and 0.1 lb ai fluoxastrobin) of **ZOLERA FX FUNGICIDE** per acre per single application.
- Do not apply more than 15.4 fl oz (0.2 lb ai tetraconazole and 0.2 lb ai fluoxastrobin) of **ZOLERA FX FUNGICIDE** per acre per year.
- Do not apply more than 0.3 lb ai of fluoxastrobin per acre per year (total of all applications of fluoxastrobin products and application methods).
- Do not apply more than 0.2 lb ai of tetraconazole per acre per year (total of all applications of tetraconazole products and application methods).
- Do not make more than 2 applications of **ZOLERA FX FUNGICIDE** per year.
- Do not make applications of **ZOLERA FX FUNGICIDE** less than 7 days apart.
- To be grown for pea and bean, dry seed only. Do not feed or harvest field pea or cowpea forage and hay.
- Pre-Harvest Interval (PHI): Do not apply **ZOLERA FX FUNGICIDE** within 14 days of harvest.

Produced For:

ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. ZOLERA and the ZOLERA FX logo are registered trademarks of an Arysta LifeScience Group Company.
 ©2019 Arysta LifeScience Group Company.