On some equations concerning the general "unitary" formula, which derives from DN Constant. New possible mathematical connections with the DN Constant, Ramanujan Recurring Numbers and some parameters of Number Theory, Cosmology and String Theory

Michele Nardelli[1], Antonio Nardelli

Abstract

In this paper, we analyze some equations concerning the general "unitary" formula, which derives from DN Constant. We obtain new possible mathematical connections with the DN Constant, Ramanujan Recurring Numbers and some parameters of Number Theory, Cosmology and String Theory

[1] M.Nardelli studied at Dipartimento di Scienze della Terra Università degli Studi di Napoli Federico II, Largo S. Marcellino, 10 - 80138 Napoli, Dipartimento di Matematica ed Applicazioni "R. Caccioppoli" - Università degli Studi di Napoli "Federico II" – Polo delle Scienze e delle Tecnologie Monte S. Angelo, Via Cintia (Fuorigrotta), 80126 Napoli, Italy

A. Nardelli studied at the Università degli Studi di Napoli Federico II - Dipartimento di Studi Umanistici – **Sezione Filosofia - scholar of Theoretical Philosophy**

Srinivasa Ramanujan (1887-1920)



https://www.moduscc.it/ramanujan-il-grande-matematico-indiano-13453-131115/

Introduction

In this paper, an octahedron could serve as a mathematical or conceptual model of the universe in the quantic phase, while the spherical surface could be used to describe the geometry of the bubble-universe.

The values $(2\sqrt{2})/\pi$, the golden ratio φ, $\zeta(2)$ and π, can be connected to the proposed cosmological model. Here's how they might be connected:

Ratio $(2\sqrt{2})/\pi$ the so called DN Constant:

This relationship may have a connection with the geometric properties of the octahedron and the sphere, which have been considered as mathematical models of the early universe and bubbles universe in eternal inflation.

Golden Ratio φ:

The golden ratio is a mathematical constant that appears in many natural and artistic contexts and is often associated with harmonious proportions and aesthetic beauty. Its emergence in this context could suggest a kind of intrinsic symmetry or harmony in the structure of the early universe and bubbles universe.

Value of π:

The value of π is a fundamental mathematical constant that appears in many geometric formulas and relationships, including the geometry of the sphere. Its appearance could indicate a direct connection between the geometry of bubbles universe and the mathematical properties of spherical surfaces.

Ultimately, the results obtained can be interpreted as manifestations of the geometric and mathematical properties of the models proposed for the early universe and universe bubbles. This suggests that there is a profound connection between geometry, mathematics and cosmological physics, and that through the analysis of

these relationships we can deepen our understanding of the universe and its fundamental phenomena.

The above values $(2\sqrt{2})/\pi$, the golden ratio φ and π, can be connected to the proposed cosmological model. This hypothesis is certainly plausible.

The various mathematical solutions and relationships can be seen as representations of the principles and laws that govern the formation and evolution of the universe.

Regarding the fundamental mathematical values, they could emerge as a consequence of the geometric and physical laws that govern the structure and evolution of the quantum universe and bubbles universe.

The multidisciplinary approach involving complex mathematical solutions and cosmological concepts can offer deeper insight into the fundamental nature of the universe and its processes. Exploring these connections could lead to new discoveries and insights into our understanding of the early universe and its complexity.

Proposal:

The initial octahedron: Let's imagine a regular octahedron, with perfectly symmetrical faces. Each face represents an ideal symmetry.

The emerging sphere: Inside the octahedron, there is an inscribed sphere. This sphere represents the bubble of the universe that emerges from the perturbations of the quantum vacuum during eternal inflation.

Expansion and transitions: As time passes, the universe expands. The faces of the octahedron begin to break, symbolizing "symmetry breaks." The sphere continues to grow, representing the expanding universe.

Constants and numbers: We integrate the mathematical results you obtained. For example, the golden ratio (φ) could be represented by a proportion between the dimensions of the octahedron and the sphere.

Entropy and complexity: Entropy increases as the universe evolves. We can represent this with a disordered growth of structures within the emerging sphere. Imagine this scene as an abstract work of art, where geometric shapes and cosmological concepts merge

In Fig.1 and Fig.2 let's imagine a regular octahedron representing the universe in its phase of high symmetry and very low entropy. Inside the octahedron we have an inscribed sphere that emerges from perturbations of the quantum vacuum during eternal inflation. As time passes, the universe expands, the faces of the octahedron break (symmetry breaks), and entropy increases. Spheres emerge from the octahedra, symbolizing the transition phases from a regime of very low entropy to a universe in which, with the passage of time, entropy increases, increasing the complexity of the universe itself.



Fig. 1



Fig. 2

Now, we have that:



Octahedron



Sphere

From the octahedron volume V = 1/3*√2 l^3 and, from the sphere volume,

V = (4/3*π*r^3) , we consider the following relationship, for r = x:

4/3*π*x^3 = 1/3*√2*l^3

Input

$$\frac{4}{3}\pi x^3 = \frac{1}{3}\sqrt{2}\, l^3$$

Exact result

$$\frac{4\pi x^3}{3} = \frac{\sqrt{2}\, l^3}{3}$$

Alternate forms

$$x^3 = \frac{l^3}{2\sqrt{2}\,\pi}$$

$$\frac{4\,\pi\,x^3}{3} - \frac{\sqrt{2}\,l^3}{3} = 0$$

Real solution

$$x = \frac{(-1)^{2/3}\,l}{\sqrt{2}\,\sqrt[3]{\pi}}$$

Solutions

$$x = -\frac{\sqrt[3]{-\frac{1}{\pi}}\,l}{\sqrt{2}}$$

$$x = \frac{l}{\sqrt{2}\,\sqrt[3]{\pi}}$$

$$x = \frac{(-1)^{2/3}\,l}{\sqrt{2}\,\sqrt[3]{\pi}}$$

Integer solution

$l = 0\,, \quad x = 0$

Implicit derivatives

$$\frac{\partial x(l)}{\partial l} = \frac{l^2}{2\sqrt{2}\,\pi\,x^2}$$

$$\frac{\partial l(x)}{\partial x} = \frac{2\sqrt{2}\,\pi\,x^2}{l^2}$$

From the alternate form

$$x^3 = \frac{l^3}{2\sqrt{2}\,\pi}$$

$$x = \frac{l}{\sqrt{2}\,\sqrt[3]{\pi}}$$

for l = 8, we have that:

8/(sqrt(2) π^(1/3)) = 8/(2sqrt2 * Pi)^1/3

Input

$$\frac{8}{\sqrt{2}\,\sqrt[3]{\pi}} = \frac{8}{\sqrt[3]{2\sqrt{2}\,\pi}}$$

Result

True

Logarithmic form

$$\log_{\sqrt{2}\,\sqrt[3]{\pi}}(8) - \log_{\sqrt{2}\,\sqrt[3]{\pi}}\left(\sqrt{2}\,\sqrt[3]{\pi}\right) = \log_{\sqrt{2}\,\sqrt[3]{\pi}}(8) - \log_{\sqrt{2}\,\sqrt[3]{\pi}}\left(\sqrt[3]{2\sqrt{2}\,\pi}\right)$$

$\log_b(x)$ is the base- b logarithm

Thence:

l/(sqrt(2) π^(1/3)) = l/(2sqrt2 * Pi)^1/3

Input

$$\frac{l}{\sqrt{2}\,\sqrt[3]{\pi}} = \frac{l}{\sqrt[3]{2\sqrt{2}\,\pi}}$$

Logarithmic form

$$\log_{\sqrt{2}\,\sqrt[3]{\pi}}(l) - \log_{\sqrt{2}\,\sqrt[3]{\pi}}\left(\sqrt{2}\,\sqrt[3]{\pi}\right) = \log_{\sqrt{2}\,\sqrt[3]{\pi}}(l) - \log_{\sqrt{2}\,\sqrt[3]{\pi}}\left(\sqrt[3]{2\sqrt{2}\,\pi}\right)$$

$\log_b(x)$ is the base- b logarithm

Now, we have that:

l/(2 sqrt(2) π)^(1/3) = (2sqrt2)/Pi

Input

$$\frac{l}{\sqrt[3]{2\sqrt{2}\,\pi}} = \frac{2\sqrt{2}}{\pi}$$

Exact result

$$\frac{l}{\sqrt{2}\,\sqrt[3]{\pi}} = \frac{2\sqrt{2}}{\pi}$$

Plot



Solution

$$l = \frac{4}{\pi^{2/3}}$$

$$l = \frac{4}{\sqrt[3]{\pi^2}}$$

The general "unitary" formula, which derives from DN Constant

The general "unitary" formula, which derives from DN Constant, is the following:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(C \times \mathrm{R} \times 2.33 \cdot 10^{-13})}} \cong 1.61803398\ldots = \frac{\sqrt{5}+1}{2}$$

where 0.9991104684 is a Rogers-Ramanujan continued fraction value

Indeed:

√(2×(2·(((2√2)/π))^(1/16))/(1/(π·0.9991104684) ×C×R×(2.33·10^-13)))

Input interpretation

$$\sqrt{2 \times \frac{2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \times 0.9991104684} C\,R \times \frac{2.33}{10^{13}}}}$$

Result

$$7.31659\times 10^{6}\sqrt{\frac{1}{C\,R}}$$

3D plots

Real part **(figures that can be related to the D-branes/Instantons)**

The key observation from the below plots and is that at $C=1$, which is taken as the energy density of the universe at the Big Bang, with $R=0$ the zero spacetime volume, the vacuum geometry brakes / or there is symmetry breaking on the vacuum quantum geometry. We see from the plots as the vacuum spacetime break/tear apart.



Imaginary part



Contour plots

Real part



Imaginary part



Alternate form assuming C and R are positive

$$\frac{7.31659\times 10^{6}}{\sqrt{C\,R}}$$

Roots

(no roots exist)

Property as a function

Parity

even

Series expansion at C=0

$$\frac{7.31659\times10^{6}\sqrt{C}\sqrt{\frac{1}{C\,R}}}{\sqrt{C}}+O\left(C^{11/2}\right)$$

(Puiseux series)

Series expansion at C=∞

$$7.31659\times10^{6}\sqrt{C}\sqrt{\frac{1}{C}}\sqrt{\frac{1}{C\,R}}+O\left(\left(\frac{1}{C}\right)^{11/2}\right)$$

(Puiseux series)

Derivative

$$\frac{\partial}{\partial C}\left(7.31659\times10^{6}\sqrt{\frac{1}{C\,R}}\right)=-3.65829\times10^{6}\,R\left(\frac{1}{C\,R}\right)^{3/2}$$

Indefinite integral

$$\int 7.31659\times10^{6}\sqrt{\frac{1}{C\,R}}\,dC=1.46332\times10^{7}\,C\sqrt{\frac{1}{C\,R}}\ +\text{constant}$$

Limit

$$\lim_{C\to\pm\infty}7.31659\times10^{6}\sqrt{\frac{1}{C\,R}}=0\approx0$$

$$\lim_{R\to\pm\infty} 7.31659\times 10^6 \sqrt{\frac{1}{C\,R}} = 0 \approx 0$$

Series representations

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{C\,R\,2.33}{(\pi\,0.99911)\,10^{13}}}} = \sqrt{-1+\frac{1.79115\times 10^{13}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}}{C\,R}}\ \sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1+\frac{1.79115\times 10^{13}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}}{C\,R}\right)^{-k}$$

for $\left|1-\frac{5.35325\times 10^{13}}{C\,R}\right| > 1$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{C\,R\,2.33}{(\pi\,0.99911)\,10^{13}}}} = \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(-1+\frac{1.79115\times 10^{13}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}}{C\,R}\right)^k}{k!}$$

for $\left|1-\frac{5.35325\times 10^{13}}{C\,R}\right| < 1$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{C\,R\,2.33}{(\pi\,0.99911)\,10^{13}}}} = \sqrt{-1+\frac{1.79115\times 10^{13}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}}{C\,R}}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(-1+\frac{1.79115\times 10^{13}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}}{C\,R}\right)^{-k}}{k!}$$

for $\left|1-\frac{5.35325\times 10^{13}}{C\,R}\right| > 1$

$\binom{n}{m}$ is the binomial coefficient

$|z|$ is the absolute value of z

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

Now, from:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times R \times 2.33 \cdot 10^{-13})}}$$

for C = 1.616255*10^{-35} (Planck Length), R = 1.265120782997423× 10^{48} that is equal to the radius of the Multiverse and 2.33*10^{-13} is the mean temperature of the CMB is equivalent around to 2.73 Kelvin , we obtain:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times 1.265120782997423 * 10^{48} \times 2.33 \cdot 10^{-13})}} =$$

Where 1.616255*10^{-35} = Planck Length ; $1.265120782997423 * 10^{48}$ = radius of Multiverse ; 2.33*10^{-13} = mean temperature of the CMB

$$= \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{(4.76428565331775479231545)}{\pi \cdot 0.9991104684}}} = \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{1.517869417}} = \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\sqrt{\frac{2}{5}}0.414293^{\frac{3}{2}}}} =$$

$$= \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\sqrt{\frac{2}{5}}(\log 2 \log 3)/\sqrt{3e}}} = 1.618033719519\ldots.$$

Indeed:

√(2×(2·((2√2)/π)^1/16)/(9√(2/5)(log2 log3)/√(3e)))

Input

$$\sqrt{2\times\frac{2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\sqrt{\frac{2}{5}}\times\frac{\log(2)\log(3)}{\sqrt{3e}}}}$$

log(x) is the natural logarithm

Exact result

$$\frac{2^{51/64}\sqrt[4]{5e}}{3^{3/4}\sqrt[32]{\pi}\sqrt{\log(2)\log(3)}}$$

Decimal approximation

1.618033719519149469459285987708851499217407474846097638812691257 1

…

1.618033719519…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

Expanded logarithmic form

$$\frac{2^{35/64}\sqrt[4]{3e\,10}\sqrt{\frac{1}{\log(2)\log(3)}}}{\sqrt[32]{\pi}\,3}$$

All 2nd roots of (2 $2^{19/32}$ sqrt((5 e)/3))/(3 $\pi^{1/16}$ log(2) log(3))

$$\frac{2^{51/64}\sqrt[4]{5e}\,e^{0}}{3^{3/4}\sqrt[32]{\pi}\sqrt{\log(2)\log(3)}}\approx 1.6180 \text{ (real, principal root)}$$

$$\frac{2^{51/64}\,\sqrt[4]{5\,e}\;e^{i\,\pi}}{3^{3/4}\,\sqrt[32]{\pi}\;\sqrt{\log(2)\log(3)}} \approx -1.6180 \quad \text{(real root)}$$

Alternative representations

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3\,e}}}} = \sqrt{\frac{4\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{9\log_e(2)\log_e(3)\sqrt{\frac{2}{5}}}{\sqrt{3\,e}}}}$$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3\,e}}}} = \sqrt{\frac{4\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{9\log^2(a)\log_a(2)\log_a(3)\sqrt{\frac{2}{5}}}{\sqrt{3\,e}}}}$$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3\,e}}}} = \sqrt{\frac{4\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{36\coth^{-1}(2)\coth^{-1}(3)\sqrt{\frac{2}{5}}}{\sqrt{3\,e}}}}$$

$\log_b(x)$ is the base-b logarithm

$\coth^{-1}(x)$ is the inverse hyperbolic cotangent function

Series representations

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}} = \frac{2^{51/64}\sqrt[4]{5e}\;\sum_{k=0}^{\infty}\binom{\frac{1}{2}+k}{k}\sum_{j=0}^{k}\frac{(-1)^j\binom{k}{j}p_{j,k}}{1+2j}}{3^{3/4}\sqrt[32]{\pi}\;\sqrt{\log(3)}}$$

$$\text{for}\left(c_k = \frac{(-1)^k\log(3)}{1+k} \text{ and } p_{j,0} = 1 \text{ and } \log(3)\,p_{j,k} = \frac{\sum_{m=1}^{k}(-k+m+jm)\,c_m\,p_{j,k-m}}{k} \text{ and } k\in\mathbb{Z} \text{ and } k>0\right)$$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}} = -\frac{\sqrt[4]{5}\;e^{1/4-i\pi\left\lfloor 1/2-\arg\left(\frac{1}{e}\right)/(2\pi)\right\rfloor}\sum_{k=0}^{\infty}\binom{\frac{1}{2}+k}{k}\sum_{j=0}^{k}\frac{(-1)^j\binom{k}{j}p_{j,k}}{-\frac{1}{2}-j}}{2^{13/64}\times 3^{3/4}\sqrt[32]{\pi}\;\sqrt{\log(3)}}$$

$$\text{for}\left(c_k = \frac{(-1)^k\log(3)}{1+k} \text{ and } p_{j,0} = 1 \text{ and } \log(3)\,p_{j,k} = \frac{\sum_{m=1}^{k}(-k+m+jm)\,c_m\,p_{j,k-m}}{k} \text{ and } k\in\mathbb{Z} \text{ and } k>0\right)$$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}} =$$

$$-\left(\left(2^{51/64}\sqrt[4]{5e}\,\sqrt{\left(-\left(2\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor - i\log(x) + i\sum_{k=1}^{\infty}\frac{(-1)^k(2-x)^k x^{-k}}{k}\right)\right.\right.\right.$$
$$\left.\left.\left.\left(2\pi\left\lfloor\frac{\arg(3-x)}{2\pi}\right\rfloor - i\log(x) + i\sum_{k=1}^{\infty}\frac{(-1)^k(3-x)^k x^{-k}}{k}\right)\right)\right)\Big/$$
$$\left(3^{3/4}\sqrt[32]{\pi}\left(2\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor - i\log(x) + i\sum_{k=1}^{\infty}\frac{(-1)^k(2-x)^k x^{-k}}{k}\right)\right.$$
$$\left.\left(2\pi\left\lfloor\frac{\arg(3-x)}{2\pi}\right\rfloor - i\log(x) + i\sum_{k=1}^{\infty}\frac{(-1)^k(3-x)^k x^{-k}}{k}\right)\right) \quad \text{for } x < 0$$

$\binom{n}{m}$ is the binomial coefficient

$\mathbb{Z}$ is the set of integers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

Integral representations

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}} = \frac{2^{51/64}\sqrt[4]{5e}\,\sqrt{\int_0^1\int_0^1 \frac{1}{(1+t_1)(1+2t_2)}\,dt_2\,dt_1}}{3^{3/4}\sqrt[32]{\pi}\left(\int_1^2 \frac{1}{t}\,dt\right)\int_1^3 \frac{1}{t}\,dt}$$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}} =$$
$$-\frac{2\times 2^{51/64}\sqrt[4]{5e}\,\pi^{31/32}\sqrt{\left(-\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\,ds\right)\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{2^{-s}\,\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\,ds}}{3^{3/4}\left(\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\,ds\right)\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{2^{-s}\,\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\,ds}$$
$$\text{for } -1 < \gamma < 0$$

$\Gamma(x)$ is the gamma function

We note that the following denominator fraction:

$$\frac{(4.76428565331775479231545)}{\pi \cdot 0.9991104684} \approx \frac{\left(\sqrt[15]{1729}\right)^{\pi}}{\pi \cdot 0.9991104684}$$

Thus, substituting in the previous formula:

$$\sqrt{2 \times \frac{2^{\cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}}{\frac{\left(\sqrt[15]{1729}\right)^{\pi}}{\pi \cdot 0.9991104684}}} = 1.617802251\ldots$$

Indeed:

√(2×(2·(((2√2)/π)^1/16))/((((1729)^1/15))^π/(π·0.9991104684)))

Input interpretation

$$\sqrt{2 \times \frac{2^{\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}}{\frac{\sqrt[15]{1729}^{\pi}}{\pi \times 0.9991104684}}}$$

Result

1.617802251235228741232261509874665413553987620147293860280427305 4

…

1.6178022512352287…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

Series representations

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{\sqrt[15]{1729}^{\pi}}{\pi 0.99911}}} = \sqrt{-1+4.17338\times 1729^{-\pi/15}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}}$$

$$\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1+4.17338\times 1729^{-\pi/15}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}\right)^{-k}$$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{\sqrt[15]{1729}^{\pi}}{\pi 0.99911}}} = \sqrt{-1+4.17338\times 1729^{-\pi/15}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}}$$

$$\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(-1+4.17338\times 1729^{-\pi/15}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}\right)^{-k}}{k!}$$

$$\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{\sqrt[15]{1729}^{\pi}}{\pi 0.99911}}} = \sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(4.17338\times 1729^{-\pi/15}\,\pi\sqrt[16]{\frac{\sqrt{2}}{\pi}}-z_0\right)^k z_0^{-k}}{k!}$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$

$\binom{n}{m}$ is the binomial coefficient

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

But, we know that

$$\frac{e^{-\frac{\pi}{\sqrt{5}}}}{\frac{\sqrt{5}}{1+\sqrt[5]{\sqrt{\varphi^5\sqrt[4]{5^3}}-1}}-\varphi+1}=1-\frac{e^{-\pi\sqrt{5}}}{1+\frac{e^{-2\pi\sqrt{5}}}{1+\frac{e^{-3\pi\sqrt{5}}}{1+\frac{e^{-4\pi\sqrt{5}}}{1+\ldots}}}}\approx 0.9991104684$$

Thus:

$$\sqrt{2\times\frac{2^{\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}}{\frac{\left(\sqrt[15]{1729}\right)^{\pi}}{\pi\cdot 0.9991104684}}}=$$

$$=\sqrt{2\times\frac{2^{\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}}{\frac{\left(\sqrt[15]{1729}\right)^{\pi}}{\pi\cdot 1-\frac{e^{(-\pi\cdot\sqrt{5})}}{1+\frac{e^{(-2\pi\cdot\sqrt{5})}}{1+\frac{e^{(-3\pi\cdot\sqrt{5})}}{1+\frac{e^{(-4\pi\cdot\sqrt{5})}}{1}}}}}}}=\sqrt{2\times\frac{2^{\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}}{1.517383339881891789}}=1.6182931189\ldots$$

Indeed:

((1729)^1/15)^π/(π·1-e^((-π·√5))/(1+e^((-2π·√5))/(1+e^((-3π·√5))/(1+e^((-4π·√5))/1))))

Input

$$\frac{\sqrt[15]{1729}^{\pi}}{\pi \times 1 - \cfrac{e^{-\pi\sqrt{5}}}{1+\cfrac{e^{-2\pi\sqrt{5}}}{1+\cfrac{e^{-3\pi\sqrt{5}}}{1+\cfrac{e^{-4\pi\sqrt{5}}}{1}}}}}$$

Exact result

$$\frac{1729^{\pi/15}}{\pi - \cfrac{e^{-\sqrt{5}\,\pi}}{1+\cfrac{e^{-2\sqrt{5}\,\pi}}{1+\cfrac{e^{-3\sqrt{5}\,\pi}}{1+e^{-4\sqrt{5}\,\pi}}}}}$$

Decimal approximation

1.5173833398818917890137811675640954545516281719627192230693647171

…

1.517383339881891789….

Alternate forms

$$\frac{1729^{\pi/15}}{\pi + \cfrac{1}{\cfrac{1}{1+e^{\sqrt{5}\,\pi}+e^{4\sqrt{5}\,\pi}}-2\cosh\left(\sqrt{5}\,\pi\right)}}$$

$$\frac{1729^{\pi/15}}{\pi - \cfrac{e^{-\sqrt{5}\,\pi}}{1+\cfrac{e^{-2\sqrt{5}\,\pi}}{1+\cfrac{e^{\sqrt{5}\,\pi}}{1+e^{4\sqrt{5}\,\pi}}}}}$$

$$\frac{1729^{\pi/15}\left(1+e^{2\sqrt{5}\,\pi}+e^{3\sqrt{5}\,\pi}+e^{4\sqrt{5}\,\pi}+e^{6\sqrt{5}\,\pi}\right)}{-e^{\sqrt{5}\,\pi}-e^{5\sqrt{5}\,\pi}+e^{2\sqrt{5}\,\pi}(\pi-1)+\pi+e^{3\sqrt{5}\,\pi}\pi+e^{4\sqrt{5}\,\pi}\pi+e^{6\sqrt{5}\,\pi}\pi}$$

cosh(x) is the hyperbolic cosine function

Series representations

$$\frac{\sqrt[15]{1729}^{\pi}}{\pi\,1-\cfrac{e^{-\pi\sqrt{5}}}{1+\cfrac{e^{-2\pi\sqrt{5}}}{1+\cfrac{e^{-3\pi\sqrt{5}}}{1+e^{-4\pi\sqrt{5}}\,\frac{1}{1}}}}} = \left(1729^{\pi/15}\left(1-e^{\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}+e^{2\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}\right)\left(1+e^{\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}+e^{2\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}+e^{3\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}+e^{4\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}\right)\right)\Big/\left(-e^{\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}-e^{2\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}-e^{5\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}+\pi+e^{2\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}\pi+e^{3\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}\pi+e^{4\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}\pi+e^{6\pi\sqrt{4}\,\sum_{k=0}^{\infty}4^{-k}\binom{1/2}{k}}\pi\right)$$

$$\frac{\sqrt[15]{1729}^{\pi}}{\pi\,1-\cfrac{e^{-\pi\sqrt{5}}}{1+\cfrac{e^{-2\pi\sqrt{5}}}{1+\cfrac{e^{-3\pi\sqrt{5}}}{1+e^{-4\pi\sqrt{5}}\ \frac{1}{1}}}}}=$$

$$\left(1729^{\pi/15}\left(1-e^{\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}}+\exp\left(2\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\right)\right.$$
$$\left(1+e^{\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}}+\exp\left(2\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)+\right.$$
$$\left.\left.\exp\left(3\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)+\exp\left(4\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\right)\right)\Big/$$
$$\left(-e^{\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}}-\exp\left(2\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)-\right.$$
$$\exp\left(5\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)+\pi+$$
$$\exp\left(2\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\pi+\exp\left(3\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\pi+$$
$$\left.\exp\left(4\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\pi+\exp\left(6\pi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\pi\right)$$

$$\frac{\sqrt[15]{1729}^{\pi}}{\pi\left(1-\cfrac{e^{-\pi\sqrt{5}}}{1+\cfrac{e^{-2\pi\sqrt{5}}}{1+\cfrac{e^{-3\pi\sqrt{5}}}{1+e^{-4\pi\sqrt{5}}\ \frac{1}{1}}}}\right)}=$$

$$\left(1729^{\pi/15}\left(1-\exp\left(\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)+\exp\left(2\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\right)\left(1+\exp\left(\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)+\exp\left(2\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)+\exp\left(3\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)+\exp\left(4\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\right)\right)\Big/$$

$$\left(-\exp\left(\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)-\exp\left(2\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)-\exp\left(5\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)+\pi+\exp\left(2\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\pi+\exp\left(3\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\pi+\exp\left(4\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\pi+\exp\left(6\pi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\pi\right)$$

for (not ($z_0 \in \mathbb{R}$ and $-\infty < z_0 \le 0$))

From the formula

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

(4 (1.6180085459)^(2 π))/(5 (3 + sqrt(5))) = 3.1415926535

we obtain:

(4 (√(2×(2·((2√2)/π)^1/16)/(9√(2/5)(log2 log3)/√(3e))))^(2 π))/(5 (3 + sqrt(5)))

Input

$$\frac{\sqrt[2\pi]{4}}{5\left(3+\sqrt{5}\right)}\sqrt{2\times\frac{\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\sqrt{\frac{2}{5}}\times\frac{\log(2)\log(3)}{\sqrt{3e}}}}$$

log(x) is the natural logarithm

Exact result

$$\frac{2^{2+(51\pi)/32}\times 3^{-(3\pi)/2}\times 5^{\pi/2-1}\,e^{\pi/2}\,\pi^{-\pi/16}\,(\log(2)\log(3))^{-\pi}}{3+\sqrt{5}}$$

Decimal approximation

3.14189977665577771135114403771804277581321420404454928910415734 39

…

3.1418997766557…. ≈ π (Ramanujan Recurring Number)

Alternative representations

$$\frac{\sqrt[2\pi]{4}}{5\left(3+\sqrt{5}\right)}\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{9\sqrt{\frac{2}{5}}\,\frac{(\log(2)\log(3))}{\sqrt{3e}}}}=\frac{\sqrt[2\pi]{4}}{5\left(3+\sqrt{5}\right)}\sqrt{\frac{4\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\log_e(2)\log_e(3)\sqrt{\frac{2}{5}}}}$$

$$\frac{4\sqrt{\frac{2\left(2^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{4\sqrt{\frac{4^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}}{\frac{36\coth^{-1}(2)\coth^{-1}(3)\sqrt{\frac{2}{5}}}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}$$

$$\frac{4\sqrt{\frac{2\left(2^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{4\sqrt{\frac{4^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}}{\frac{9\log^2(a)\log_a(2)\log_a(3)\sqrt{\frac{2}{5}}}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}$$

$\log_b(x)$ is the base- b logarithm

$\coth^{-1}(x)$ is the inverse hyperbolic cotangent function

Series representations

$$\frac{4\sqrt{\frac{2\left(2^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{2^{2+(51\pi)/32}\times 3^{-(3\pi)/2}\times 5^{-1+\pi/2}\,e^{\pi/2}\,\pi^{1-\pi/16}\log^{-\pi}(3)\sum_{k=0}^{\infty}\binom{k+\pi}{k}\sum_{j=0}^{k}\frac{(-1)^j\binom{k}{j}p_{j,k}}{j+\pi}}{3+\sqrt{5}}$$

$$\text{for}\left(c_k = \frac{(-1)^k\log(3)}{1+k} \text{ and } p_{j,0} = 1 \text{ and } \log(3)\,p_{j,k} = \frac{\sum_{m=1}^{k}(-k+m+j\,m)\,c_m\,p_{j,k-m}}{k} \text{ and } k\in\mathbb{Z} \text{ and } k>0\right)$$

$$\frac{\sqrt[4]{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{1}{3+\sqrt{5}}\,2^{2+(51\pi)/32}\times 3^{-(3\pi)/2}\times 5^{-1+\pi/2}$$
$$e^{\pi/2}\,\pi^{-\pi/16}\left(\left(2i\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor+\log(x)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(2-x)^k\,x^{-k}}{k}\right)\right.$$
$$\left.\left(2i\pi\left\lfloor\frac{\arg(3-x)}{2\pi}\right\rfloor+\log(x)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(3-x)^k\,x^{-k}}{k}\right)\right)^{-\pi}\quad\text{for } x<0$$

$$\frac{\sqrt[4]{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{1}{3+\sqrt{5}}2^{2+(51\pi)/32}\times 3^{-(3\pi)/2}\times 5^{-1+\pi/2}\,e^{\pi/2}\,\pi^{-\pi/16}$$
$$\left(\left(\log(z_0)+\left\lfloor\frac{\arg(2-z_0)}{2\pi}\right\rfloor\left(\log\left(\frac{1}{z_0}\right)+\log(z_0)\right)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(2-z_0)^k\,z_0^{-k}}{k}\right)\right.$$
$$\left.\left(\log(z_0)+\left\lfloor\frac{\arg(3-z_0)}{2\pi}\right\rfloor\left(\log\left(\frac{1}{z_0}\right)+\log(z_0)\right)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(3-z_0)^k\,z_0^{-k}}{k}\right)\right)^{-\pi}$$

$$\frac{\sqrt[4]{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{1}{3+\sqrt{5}}\,2^{2+(51\pi)/32}\times 3^{-(3\pi)/2}\times 5^{-1+\pi/2}\,e^{\pi/2}\,\pi^{-\pi/16}$$
$$\left(\left(2i\pi\left\lfloor\frac{\pi-\arg\left(\frac{1}{z_0}\right)-\arg(z_0)}{2\pi}\right\rfloor+\log(z_0)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(2-z_0)^k\,z_0^{-k}}{k}\right)\right.$$
$$\left.\left(2i\pi\left\lfloor\frac{\pi-\arg\left(\frac{1}{z_0}\right)-\arg(z_0)}{2\pi}\right\rfloor+\log(z_0)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(3-z_0)^k\,z_0^{-k}}{k}\right)\right)^{-\pi}$$

$\binom{n}{m}$ is the binomial coefficient

$\mathbb{Z}$ is the set of integers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

Integral representations

$$\frac{\sqrt[4]{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{9\sqrt{\frac{2}{5}}\frac{(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{2^{2+(51\pi)/32}\times 3^{-(3\pi)/2}\times 5^{-1+\pi/2}\, e^{\pi/2}\, \pi^{-\pi/16}\left(\int_0^1\int_0^1 \frac{1}{(1+t_1)(1+2t_2)}\, dt_2\, dt_1\right)^{-\pi}}{3+\sqrt{5}}$$

$$\frac{\sqrt[4]{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{9\sqrt{\frac{2}{5}}\frac{(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)} = \frac{1}{3+\sqrt{5}}\, 2^{2+(115\pi)/32}\times 3^{-(3\pi)/2}\times 5^{-1+\pi/2}\, e^{\pi/2}\, \pi^{(31\pi)/16}$$
$$\left(\left(-\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\, ds\right)\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{2^{-s}\,\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\, ds\right)^{-\pi} \quad \text{for } -1 < \gamma < 0$$

$\Gamma(x)$ is the gamma function

1/6((4 (√(2×(2·((2√2)/π)^1/16)/(9√(2/5)(log2 log3)/√(3e))))^(2 π))/(5 (3 + sqrt(5))))^2

Input

$$\frac{1}{6}\left(\frac{4\sqrt{2\times\frac{\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\sqrt{\frac{2}{5}}\times\frac{\log(2)\log(3)}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2$$

log(x) is the natural logarithm

Exact result

$$\frac{2^{3+(51\pi)/16}\times 3^{-1-3\pi}\times 5^{\pi-2}\,e^{\pi}\,\pi^{-\pi/8}\,(\log(2)\log(3))^{-2\pi}}{\left(3+\sqrt{5}\right)^2}$$

Decimal approximation

1.64525570109160431087165810426750878077788091602097620117795446 06

…

1.6452557010916…. ≈ ζ(2) = $\pi^2/6$ = 1.644934 (trace of the instanton shape and Ramanujan Recurring Number)

Alternative representations

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2=\frac{1}{6}\left(\frac{4\sqrt{\frac{4\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{9\log_e(2)\log_e(3)\sqrt{\frac{2}{5}}}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2$$

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2=\frac{1}{6}\left(\frac{4\sqrt{\frac{4\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{36\coth^{-1}(2)\coth^{-1}(3)\sqrt{\frac{2}{5}}}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2$$

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2=\frac{1}{6}\left(\frac{4\sqrt{\frac{4\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{9\,\mathrm{Li}_1(-2)\mathrm{Li}_1(-1)\sqrt{\frac{2}{5}}}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2$$

$\log_b(x)$ is the base- b logarithm

$\coth^{-1}(x)$ is the inverse hyperbolic cotangent function

$\mathrm{Li}_n(x)$ is the polylogarithm function

Series representations

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2 =$$

$$\frac{2^{4+(51\pi)/16}\times 3^{-1-3\pi}\times 5^{-2+\pi}\,e^{\pi}\,\pi^{1-\pi/8}\log^{-2\pi}(3)\sum_{k=0}^{\infty}\binom{k+2\pi}{k}\sum_{j=0}^{k}\frac{(-1)^j\binom{k}{j}p_{j,k}}{j+2\pi}}{\left(3+\sqrt{5}\right)^2}$$

$$\text{for}\left(c_k=\frac{(-1)^k\log(3)}{1+k}\text{ and }p_{j,0}=1\text{ and }\log(3)\,p_{j,k}=\frac{\sum_{m=1}^{k}(-k+m+j\,m)\,c_m\,p_{j,k-m}}{k}\text{ and }k\in\mathbb{Z}\text{ and }k>0\right)$$

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2 = -\frac{1}{\left(3+\sqrt{5}\right)^2}2^{4+(51\pi)/16}\times 3^{-1-3\pi}\times 5^{-2+\pi}$$

$$e^{\pi-4i\pi^2\left[1/2-\arg\left(\frac{1}{e}\right)/(2\pi)\right]}\pi^{1-\pi/8}\log^{-2\pi}(3)\sum_{k=0}^{\infty}\binom{k+2\pi}{k}\sum_{j=0}^{k}\frac{(-1)^j\binom{k}{j}p_{j,k}}{-j-2\pi}$$

$$\text{for}\left(c_k=\frac{(-1)^k\log(3)}{1+k}\text{ and }p_{j,0}=1\text{ and }\log(3)\,p_{j,k}=\frac{\sum_{m=1}^{k}(-k+m+j\,m)\,c_m\,p_{j,k-m}}{k}\text{ and }k\in\mathbb{Z}\text{ and }k>0\right)$$

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{2\left(2^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2 = \frac{1}{\left(3+\sqrt{5}\right)^2}\,2^{3+(51\pi)/16}\times 3^{-1-3\pi}\times$$

$$5^{-2+\pi}\,e^{\pi}\,\pi^{-\pi/8}\left(\left(2\,i\,\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor+\log(x)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(2-x)^k\,x^{-k}}{k}\right)\right.$$
$$\left.\left(2\,i\,\pi\left\lfloor\frac{\arg(3-x)}{2\pi}\right\rfloor+\log(x)-\sum_{k=1}^{\infty}\frac{(-1)^k\,(3-x)^k\,x^{-k}}{k}\right)\right)^{-2\pi}\quad\text{for } x<0$$

$\binom{n}{m}$ is the binomial coefficient

$\mathbb{Z}$ is the set of integers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

Integral representations

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{2\left(2^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}\right)}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2 =$$

$$\frac{2^{3+(51\pi)/16}\times 3^{-1-3\pi}\times 5^{-2+\pi}\,e^{\pi}\,\pi^{-\pi/8}\left(\int_0^1\int_0^1\frac{1}{(1+t_1)(1+2t_2)}\,dt_2\,dt_1\right)^{-2\pi}}{\left(3+\sqrt{5}\right)^2}$$

$$\frac{1}{6}\left(4\sqrt{\frac{2\left(2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}\right)^{2\pi}}{\frac{9\sqrt{\frac{2}{5}}\,(\log(2)\log(3))}{\sqrt{3e}}}}\Big/\left(5\left(3+\sqrt{5}\right)\right)\right)^2 = \frac{1}{\left(3+\sqrt{5}\right)^2}\,2^{3+(115\pi)/16}\times 3^{-1-3\pi}\times 5^{-2+\pi}\,e^{\pi}\,\pi^{(31\pi)/8}$$

$$\left(\left(-\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\,ds\right)\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{2^{-s}\,\Gamma(-s)^2\,\Gamma(1+s)}{\Gamma(1-s)}\,ds\right)^{-2\pi}\quad \text{for } -1<\gamma<0$$

$\Gamma(x)$ is the gamma function

With regard the extended Del Gaudio-Nardelli Constant :

$$\sqrt[2\pi]{\frac{\frac{5}{12}(3+\sqrt{5})d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\frac{\frac{\sqrt{2}}{12}d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}}\times\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}$$

from

$$\sqrt[2\pi]{\frac{\frac{5}{12}(3+\sqrt{5})d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\frac{\frac{\sqrt{2}}{12}d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}} = 2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}$$

and

$$\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}$$

for q = 1 and p = 2, we obtain:

$$2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)$$

From the general "unitary" formula, which derives from DN Constant:

$$\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi\cdot 0.9991104684}(C\times \mathrm{R}\times 2.33\cdot 10^{-13})}}\cong 1.61803398\ldots=\frac{\sqrt{5}+1}{2}$$

instead of the 16th root of DN Constant, we insert the following expression concerning the extended DN Constant,

$$2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}$$

and obtain:

$$\sqrt{2\times\frac{2\cdot\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi\cdot 0.9991104684}(1.616255\cdot 10^{-35}\times \mathrm{R}\times 2.33\cdot 10^{-13})}}$$

for R = $1.249031956736993500 72\times 10^{48}$

$$\sqrt{2\times\frac{2\cdot\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi\cdot 0.9991104684}(1.616255\cdot 10^{-35}\times 1.24903195673699350072\times 10^{48}\times 2.33\cdot 10^{-13})}}=1.61803$$

Indeed:

$$\sqrt{\frac{2\times 2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi\times 0.9991104684}\times\frac{1.616255\times\frac{1.24903195673699350072\times 10^{48}\times 2.33}{10^{13}}}{10^{35}}}}$$

we obtain:

√((2×2(((((((2^(-1/π) (5 (3 + √(5)) π)^(1/(2 π)) (∛(-1/2+√(1/4+8/27)) +∛(-1/2-√(1/4+8/27))))))^1/16)))))/(1/(π·0.9991104684) (1.616255·10^(-35)×(1.24903195673699350072×10^48)×2.33·10^(-13))))

Input interpretation

$$\sqrt{\frac{2\times 2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi\times 0.9991104684}\times\frac{1.616255\times\frac{1.24903195673699350072\times 10^{48}\times 2.33}{10^{13}}}{10^{35}}}}$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Result

1.61024… +
0.158595… i

Alternate complex forms

$1.61024 + 0.158595\,i$

$1.61803\,(\cos(0.0981748) + i\sin(0.0981748))$

$1.61803\,e^{0.0981748\,i}$

i is the imaginary unit

Polar coordinates

$r = 1.61803$ (radius), $\theta = 0.0981748$ (angle)

1.61803 result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

From the formula

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

(4 (1.6180085459)^(2 π))/(5 (3 + sqrt(5))) = 3.1415926535

we obtain:

(4((√((4(((((((2^(-1/π)(5(3+√(5))π)^(1/(2π))(∛(-1/2+√(1/4+8/27))+∛(-1/2-√(1/4+8/27))))))^1/16))))))/(1/(π·0.9991104684)(1.616255·10^(-35)×(1.249031956736993500 72×10^48)×2.33e-13))))^(2π))/(5(3+√(5)))

Input interpretation

$$\sqrt[2\pi]{\frac{4\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}}{\frac{1}{\pi\times 0.9991104684}\times\frac{1.616255\left(1.249031956736993500 72\times 10^{48}\times 2.33\times 10^{-13}\right)}{10^{35}}}}{5\left(3+\sqrt{5}\right)}}$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Result

2.56286… +
1.81749… i

Alternate complex forms

$2.56286 + 1.81749\,i$

$3.1419\,(\cos(0.61685) + i\sin(0.61685))$

$3.1419\,e^{0.61685\,i}$

i is the imaginary unit

Polar coordinates

$r = 3.1419$ (radius), $\theta = 0.61685$ (angle)

3.1419 ≈ π (Ramanujan Recurring Number)

1/6((4((√((4(((((((2^(-1/π)(5(3+√(5))π)^(1/(2π))(∛(-1/2+√(1/4+8/27))+∛(-1/2-√(1/4+8/27))))))^1/16)))))/(1/(π·0.9991104684)(1.616255e-35×(1.24903195673699350072e+48)×2.33e-13)))))^(2π))/(5(3+√(5))))^2

Input interpretation

$$\frac{1}{6}\left(\frac{4\sqrt{\frac{4\sqrt[16]{2^{-1/\pi}\,2\pi\sqrt{5\left(3+\sqrt{5}\right)}\,\pi}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}}{\frac{1}{\pi\times 0.9991104684}\left(1.616255\times 10^{-35}\times 1.24903195673699350072\times 10^{48}\times 2.33\times 10^{-13}\right)}}^{2\pi}}{5\left(3+\sqrt{5}\right)}\right)^2$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Result

0.544166… +
1.55266… i

Alternate complex forms

$0.544166 + 1.55266\,i$

$1.64526\,(\cos(1.2337) + i\sin(1.2337))$

$1.64526\,e^{1.2337\,i}$

i is the imaginary unit

Polar coordinates

$r = 1.64526$ (radius), $\theta = 1.2337$ (angle)

1.64526 ≈ ζ(2) = π²/6 = 1.644934 (trace of the instanton shape and Ramanujan Recurring Number)

From this further Cardano Formula

$$x = -\frac{b}{3a} + \sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}} + \sqrt[3]{-\frac{q}{2} - \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}.$$

we obtain:

$$2\pi\sqrt{\frac{\frac{5}{12}(3+\sqrt{5})d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3} \times \frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}} \times \frac{1}{\frac{\frac{\sqrt{2}}{12}d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}} \times \left(-\frac{b}{3a} + \sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}} + \sqrt[3]{-\frac{q}{2} - \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}\right)}$$

i.e.

$$\sqrt[2\pi]{\frac{\frac{5}{12}\left(3+\sqrt{5}\right)d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\frac{\frac{\sqrt{2}}{12}d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}}=2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Thus:

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\times\left(-\frac{b}{3a}+\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}\right)$$

(((2^(-1/π) (5 (3 + sqrt(5)) π)^(1/(2 π)))) (-b/(3a)+((∛(-q/2+√((q^2)/4+(p^3)/27)) +∛(-q/2-√((q^2)/4+(p^3)/27))))))

Input

$$\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(-\frac{b}{3a}+\left(\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}\right)\right)$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Exact result

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\left(-\frac{b}{3a}+\sqrt[3]{-\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}+\sqrt[3]{\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}\right)$$

$\sqrt[n]{x}$ is the real-valued n^{th} root of x

Alternate form

$$-\frac{1}{3\sqrt{3}\,a}2^{-1/3-1/\pi}\sqrt[2\pi]{\left(15+5\sqrt{5}\right)\pi}\left(3a\left(\sqrt[3]{3\sqrt{3}\,q-\sqrt{4\,p^3+27q^2}}+\sqrt[3]{\sqrt{4\,p^3+27q^2}+3\sqrt{3}\,q}\right)+\sqrt[3]{2}\,\sqrt{3}\,b\right)$$

Expanded forms

$$-\frac{2^{-1/\pi}\sqrt[2\pi]{15\pi+5\sqrt{5}\,\pi}\,b}{3a}+2^{-1/\pi}\sqrt[2\pi]{15\pi+5\sqrt{5}\,\pi}\sqrt[3]{\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}-2^{-1/\pi}\sqrt[2\pi]{15\pi+5\sqrt{5}\,\pi}\sqrt[3]{\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}+\frac{q}{2}}$$

$$-\frac{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\,b}{3a}+2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\sqrt[3]{-\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}+2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\sqrt[3]{\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}$$

Alternate forms assuming a, b, p, and q are positive

$$\frac{1}{3a}2^{-(1+\pi)/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\left(2^{2/3}\sqrt[3]{3}\,a\left(\sqrt[3]{\sqrt{12\,p^3+81\,q^2}-9q}-\sqrt[3]{\sqrt{12\,p^3+81\,q^2}+9q}\right)-2b\right)$$

$$-\frac{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\,b}{3a}+2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\sqrt[3]{\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}-2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\sqrt[3]{\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}+\frac{q}{2}}$$

Derivative

$$\frac{\partial}{\partial p}\left(\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(-\frac{b}{3a}+\left(\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}\right)\right)\right)=$$

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\left(\frac{p^2}{54\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}\sqrt[3]{\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}^2}-\frac{p^2}{54\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}\sqrt[3]{-\sqrt{\frac{p^3}{27}+\frac{q^2}{4}}-\frac{q}{2}}^2}\right)$$

For q = 1, p = 2, a = 4 and b = 8:

(((2^(-1/π) (5 (3 + sqrt(5)) π)^(1/(2 π)))) (-8/(3*4)+((∛(-1/2+√(1/4+8/27)) +∛(-1/2-√(1/4+8/27))))))

Input

$$\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(-\frac{8}{3\times4}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Exact result

$$2^{-1/\pi}\left(-\frac{2}{3}+\sqrt[3]{\frac{\sqrt{\frac{59}{3}}}{6}-\frac{1}{2}}-\sqrt[3]{\frac{1}{2}+\frac{\sqrt{\frac{59}{3}}}{6}}\right)\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Decimal approximation

$-1.812273638777984635375567526232947692203814284197178847686462729$

…

-1.812273638777….

Alternate forms

$$\frac{1}{3}\times 2^{-(1+\pi)/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

root of $x^3+12x^2+120x+568$ near $x=-6.72039$

$$\frac{1}{3}\times 2^{-1/\pi}\left(-2-2\times 3^{2/3}\sqrt[3]{\frac{2}{\sqrt{177}-9}}+\sqrt[3]{\frac{3}{2}\left(\sqrt{177}-9\right)}\right)\sqrt[2\pi]{\left(15+5\sqrt{5}\right)\pi}$$

$$\frac{1}{3}\times 2^{-1-1/\pi}\left(-4+2^{2/3}\sqrt[3]{3\left(\sqrt{177}-9\right)}-2^{2/3}\sqrt[3]{3\left(9+\sqrt{177}\right)}\right)\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Expanded form

$$-\frac{1}{3}\times 2^{1-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}+2^{-1/\pi}\sqrt[3]{\frac{\sqrt{\frac{59}{3}}}{6}-\frac{1}{2}}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}-2^{-1/\pi}\sqrt[3]{\frac{1}{2}+\frac{\sqrt{\frac{59}{3}}}{6}}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Series representations

$$\left(-\frac{8}{3\times 4}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}=$$

$$\frac{1}{3}\times 2^{-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)}$$

$$\left(-2+3\sqrt[3]{-\frac{1}{2}-\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}+\right.$$

$$\left.3\sqrt[3]{-\frac{1}{2}+\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}\right)$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \leq 0))$

$$\left(-\frac{8}{3\times 4}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}=$$
$$\frac{1}{3}\times 2^{-1/\pi}\sqrt[2\pi]{5}$$
$$\sqrt[2\pi]{\pi\left(3+\exp\left(i\pi\left\lfloor\frac{\arg(5-x)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k(5-x)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)}$$
$$\left(-2+3\sqrt[3]{-\frac{1}{2}-\exp\left(i\pi\left\lfloor\frac{\arg\left(\frac{59}{108}-x\right)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{59}{108}-x\right)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}}+\right.$$
$$\left.3\sqrt[3]{-\frac{1}{2}+\exp\left(i\pi\left\lfloor\frac{\arg\left(\frac{59}{108}-x\right)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{59}{108}-x\right)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}}\right)$$

for $(x\in\mathbb{R}$ and $x<0)$

$$\left(-\frac{8}{3\times 4}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}=$$
$$\frac{1}{3}\times 2^{-1/\pi}\sqrt[2\pi]{5}\left(\pi\left(3+\left(\frac{1}{z_0}\right)^{1/2\lfloor\arg(5-z_0)/(2\pi)\rfloor}z_0^{1/2+1/2\lfloor\arg(5-z_0)/(2\pi)\rfloor}\right.\right.$$
$$\left.\left.\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\right)^{\wedge}\left(\frac{1}{2\pi}\right)\left(-2+3\right.$$
$$\sqrt[3]{-\frac{1}{2}-\left(\frac{1}{z_0}\right)^{1/2\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor}z_0^{1/2\left(1+\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor\right)}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}$$
$$\left.+3\sqrt[3]{-\frac{1}{2}+\left(\frac{1}{z_0}\right)^{1/2\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor}z_0^{1/2\left(1+\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor\right)}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}\right)$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\sqrt[n]{x}$ is the real-valued n^{th} root of x

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x\rfloor$ is the floor function

i is the imaginary unit

From which:

-1/2*((((2^(-1/π)(5(3+sqrt(5))π)^(1/(2π)))) (-2/3+((∛(-1/2+√(1/4+8/27)) +∛(-1/2-√(1/4+8/27)))))))

Input

$$-\frac{1}{2}\left(\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(-\frac{2}{3}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)\right)$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Exact result

$$-2^{-1-1/\pi}\left(-\frac{2}{3}+\sqrt[3]{\frac{\sqrt{\frac{59}{3}}}{6}-\frac{1}{2}}-\sqrt[3]{\frac{1}{2}+\frac{\sqrt{\frac{59}{3}}}{6}}\right)\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Decimal approximation

0.906136819388992317687783763116473846101907142098589423843231364 8

…

0.90613681938899231…. ≈ 0.9003163161571…. = $\frac{2\sqrt{2}}{\pi}$ (DN Constant) (We note that, with regard the inflation, during a period of almost exponential expansion $\dot{H} < 0$ so that $\epsilon > 0$. Indeed, the value 0.9003163161571… = (2√2)/π > 0 and also the n_s = spectral index = 0.90-0.97 is near to the DN Constant value. Also the squared sound speed of the gravitino's longitudinal polarization mode could pass through 0.9003163161571 in the early universe, in the so-called quasi-de Sitter phase of inflation)

Alternate forms

$$\frac{1}{3}\times 2^{-2-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi} \quad \text{root of } x^3-12x^2+120x-568 \text{ near } x=6.72039$$

$$\frac{1}{3}\times 2^{-1-1/\pi}\left(2-2\times 3^{2/3}\sqrt[3]{\frac{2}{9+\sqrt{177}}}+\sqrt[3]{\frac{3}{2}\left(9+\sqrt{177}\right)}\right)\sqrt[2\pi]{\left(15+5\sqrt{5}\right)\pi}$$

$$-\frac{1}{3}\times 2^{-2-1/\pi}\left(-4+2^{2/3}\sqrt[3]{3\left(\sqrt{177}-9\right)}-2^{2/3}\sqrt[3]{3\left(9+\sqrt{177}\right)}\right)\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Expanded form

$$\frac{1}{3}\times 2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}-2^{-1-1/\pi}\sqrt[3]{\frac{\sqrt{\frac{59}{3}}}{6}-\frac{1}{2}}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}+2^{-1-1/\pi}\sqrt[3]{\frac{1}{2}+\frac{\sqrt{\frac{59}{3}}}{6}}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Series representations

$$\frac{1}{2}\left(\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(-\frac{2}{3}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)\right)$$

$$(-1)=-\frac{1}{3}\times 2^{-1-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)}$$

$$\left(-2+3\sqrt[3]{-\frac{1}{2}-\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}+3\sqrt[3]{-\frac{1}{2}+\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}\right)$$

for $(\text{not } (z_0\in\mathbb{R} \text{ and } -\infty<z_0\le 0))$

$$\frac{1}{2}\left(\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(-\frac{2}{3}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)\right)$$

$$(-1)=-\frac{1}{3}\times 2^{-1-1/\pi}\sqrt[2\pi]{5}$$

$$\sqrt[2\pi]{\pi\left(3+\exp\left(i\pi\left\lfloor\frac{\arg(5-x)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k(5-x)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)}$$

$$\left(-2+3\sqrt[3]{-\frac{1}{2}-\exp\left(i\pi\left\lfloor\frac{\arg\left(\frac{59}{108}-x\right)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{59}{108}-x\right)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}}+\right.$$

$$\left.3\sqrt[3]{-\frac{1}{2}+\exp\left(i\pi\left\lfloor\frac{\arg\left(\frac{59}{108}-x\right)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{59}{108}-x\right)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}}\right)$$

for $(x\in\mathbb{R}$ and $x<0)$

$$\frac{1}{2}\left(\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(-\frac{2}{3}+\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\right)\right)$$

$$(-1)=-\frac{1}{3}\times 2^{-1-1/\pi}\sqrt[2\pi]{5}\left(\pi\left(3+\left(\frac{1}{z_0}\right)^{1/2\lfloor\arg(5-z_0)/(2\pi)\rfloor}z_0^{1/2+1/2\lfloor\arg(5-z_0)/(2\pi)\rfloor}\right.\right.$$

$$\left.\left.\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\right)\wedge\left(\frac{1}{2\pi}\right)\left(-2+3\right.$$

$$\sqrt[3]{-\frac{1}{2}-\left(\frac{1}{z_0}\right)^{1/2\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor}z_0^{1/2\left(1+\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor\right)}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}$$

$$\left.+3\sqrt[3]{-\frac{1}{2}+\left(\frac{1}{z_0}\right)^{1/2\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor}z_0^{1/2\left(1+\left\lfloor\arg\left(\frac{59}{108}-z_0\right)/(2\pi)\right\rfloor\right)}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}\right)$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\sqrt[n]{x}$ is the real-valued n^{th} root of x

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x\rfloor$ is the floor function

i is the imaginary unit

Thence, from the general "unitary" formula, which derives from DN Constant:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(C \times \mathrm{R} \times 2.33 \cdot 10^{-13})}} \cong 1.61803398\ldots = \frac{\sqrt{5}+1}{2}$$

we obtain:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(2^{-\frac{1}{\pi}} \sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\left(-\frac{b}{3a}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times \mathrm{R} \times 2.33 \cdot 10^{-13})}}$$

for a = 4, b = 8 and R = -1.469895830716*10^{48} :

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(2^{-\frac{1}{\pi}} \sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\left(-\frac{8}{12}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times (-1.469895830716 \times 10^{48}) \times 2.33 \cdot 10^{-13})}}$$

√(2×(2·(((((2^(-1/π) (((5 (3 +√5)π)^(1/(2Pi)))))))))^(1/16))(-8/12+∛(-1/2+√(1/4+8/27))
+∛(-1/2-√(1/4+8/27))))/(1/(π·0.9991104684) (1.616255·10^(-35)×(- 1.469895830716
× 10^48)×2.33·10^(-13))))

Input interpretation



$\sqrt[3]{x}$ is the real-valued 3rd root of x

Result

1.6180339800003588468645295725358500061027310519165682458709887505

…

1.61803398…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

Or:



√(2×(2·(((((2^(-1/π) (((5 (3 +√5)π)^(1/(2Pi)))))))))^(1/16))(-8/12+∛(-1/2+√(59/108))
+∛(-1/2-√(59/108))))/(-5.5354427120861603914/(π·0.9991104684)))

Input interpretation



$\sqrt[3]{x}$ is the real-valued 3rd root of x

Result

1.618033980…

1.61803398…. as above

All 2nd roots of 2.61803

$1.61803\, e^{0} \approx 1.6180$ (real, principal root)

$1.61803\, e^{i\pi} \approx -1.6180$ (real root)

Series representations

$$\sqrt{-\frac{2\left(2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(-\frac{8}{12}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)\right)}{\frac{5.535442712086160391400000}{\pi\, 0.99911}}} =$$

$$\sqrt{\left(-1-0.721973\,\pi\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)}$$

$$\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1-0.721973\,\pi\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)^{-k}$$

$$\sqrt{-\frac{2\left(2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(-\frac{8}{12}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)\right)}{\frac{5.53544271208616039140000}{\pi\,0.99911}}} =$$

$$\sqrt{\left(-1-0.721973\,\pi\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)}$$

$$\sum_{k=0}^{\infty}\frac{1}{k!}(-1)^k\left(-\frac{1}{2}\right)_k\left(-1-0.721973\,\pi\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)^{-k}$$

$$\sqrt{-\frac{2\left(2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(-\frac{8}{12}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)\right)}{\frac{5.53544271208616039140000}{\pi\,0.99911}}} =$$

$$\sqrt{z_0}\sum_{k=0}^{\infty}\frac{1}{k!}(-1)^k\left(-\frac{1}{2}\right)_k z_0^{-k}\left(-z_0-0.721973\,\pi\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)^{k}$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$

$\sqrt[n]{x}$ is the real-valued n^{th} root of x

$\binom{n}{m}$ is the binomial coefficient

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

And again:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(2^{-\frac{1}{\pi}} \sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)}}{(-1.763554877452867723027659095706 1)}}$$

Input interpretation

$$\sqrt{2\left(-\frac{2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)}{1.763554877452867723027659095706 1}\right)}$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Result

1.618033980000358846864529572535 8…

1.61803398…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

All 2nd roots of 2.618033960435801652840850738926 8

$1.618033980000358846864529572535 83\, e^{0} \approx 1.6180$ (real, principal root)

$1.618033980000358846864529572535 83\, e^{i\pi} \approx -1.6180$ (real root)

Series representations

$$\sqrt{-\frac{2\left(2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)\right)}{1.763554877452867723027659095706100 00}} =$$

$$\sqrt{\left(-1-2.268146033412505978034423042107634 53\right.}$$

$$\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}$$

$$\left.\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)$$

$$\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1-2.268146033412505978034423042107634 53\right.$$

$$\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}$$

$$\left.\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)^{-k}$$

$$\sqrt{-\frac{2\left(2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)\right)}{1.763554877452867723027659095706100000}} =$$

$$\sqrt{\left(-1-2.268146033412505978034423042107634530\right.}$$

$$\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}$$

$$\left.\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)$$

$$\sum_{k=0}^{\infty}\frac{1}{k!}(-1)^k\left(-\frac{1}{2}\right)_k\left(-1-2.268146033412505978034423042107634530\right.$$

$$\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}$$

$$\left.\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)\right)^{-k}$$

$$\sqrt{-\frac{2\left(2\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)\right)}{1.763554877452867723027659095706100000}} =$$

$$\sqrt{z_0}$$

$$\sum_{k=0}^{\infty}\frac{1}{k!}(-1)^k\left(-\frac{1}{2}\right)_k z_0^{-k}\left(-z_0-2.268146033412505978034423042107634530\right.$$

$$\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}+\right.$$

$$\left.\left.\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}\right)^{k}\right) \quad \text{for } (\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$$

$\sqrt[n]{x}$ is the real-valued n^{th} root of x

$\binom{n}{m}$ is the binomial coefficient

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

And again:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(2^{-\frac{1}{\pi}} \sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)}}{(-1.763554877452867723027659095706 1)}}$$

(((-2/3+∛(-1/2+√(59/108)) +∛(-1/2-√(59/108))))/(((-1.763554877452867723027659095706 1))))

Input interpretation

$$-\frac{-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}}{1.763554877452867723027659095706 1}$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Result

0.635117360113453550094146405698 42…

0.63511736….

Series representations

$$-\frac{-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}}{1.763554877452867723027659095706100 00}=$$
$$-0.567036508353126494508605760526908632$$
$$\left(-0.666666666666666666666666666666666667+\right.$$
$$1.000000000000000000000000000000000000$$
$$\sqrt[3]{-\frac{1}{2}-\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{49}{108}\right)^k\left(-\frac{1}{2}\right)_k}{k!}}+$$
$$1.000000000000000000000000000000000000$$
$$\left.\sqrt[3]{-\frac{1}{2}+\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{49}{108}\right)^k\left(-\frac{1}{2}\right)_k}{k!}}\right)$$

$$-\frac{-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}}{1.763554877452867723027659095706100 00}=$$
$$-0.567036508353126494508605760526908632$$
$$\left(-0.666666666666666666666666666666666667+\right.$$
$$1.000000000000000000000000000000000000$$
$$\sqrt[3]{-\frac{1}{2}-\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}+$$
$$1.000000000000000000000000000000000000$$
$$\left.\sqrt[3]{-\frac{1}{2}+\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(\frac{59}{108}-z_0\right)^k z_0^{-k}}{k!}}\right)$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \leq 0))$

$$-\frac{-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}}{1.763554877452867723027659095706100000}=$$
$$-0.567036508353126494508605760526908632$$
$$\left(-0.666666666666666666666666666666666667+\right.$$
$$1.000000000000000000000000000000000000$$
$$\sqrt[3]{-\frac{1}{2}-\frac{\sum_{j=0}^{\infty}\text{Res}_{s=-j}\left(-\frac{49}{108}\right)^{-s}\Gamma\left(-\frac{1}{2}-s\right)\Gamma(s)}{2\sqrt{\pi}}}+$$
$$1.000000000000000000000000000000000000$$
$$\left.\sqrt[3]{-\frac{1}{2}+\frac{\sum_{j=0}^{\infty}\text{Res}_{s=-j}\left(-\frac{49}{108}\right)^{-s}\Gamma\left(-\frac{1}{2}-s\right)\Gamma(s)}{2\sqrt{\pi}}}\right)$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\sqrt[n]{x}$ is the real-valued n^{th} root of x

$\mathbb{R}$ is the set of real numbers

$\Gamma(x)$ is the gamma function

$\underset{z=z_0}{\text{Res}}\, f$ is a complex residue

And again:

$$\sqrt{4\cdot\left(\sqrt[16]{2^{-\frac{1}{\pi}}\sqrt[2\pi]{\left(5\left(3+\sqrt{5}\right)\pi\right)}}\right)0.63511736}$$

√(4·(((2^(-1/π) (((5 (3+√5)π)^(1/(2Pi))))))^(1/16)0.63511736))

Input interpretation

$$\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\times 0.63511736\right)}$$

Result

1.6180339798558409114868793011476459142925815835677790206645626849

…

1.61803397985584…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

All 2nd roots of 2.61803

$1.61803\, e^{0} \approx 1.6180$ (real, principal root)

$1.61803\, e^{i\pi} \approx -1.6180$ (real root)

Series representations

$$\sqrt{4\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ \ 0.635117} = \sqrt{-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}}$$

$$\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\right)^{-k}$$

$$\sqrt{4\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ \ 0.635117} = \sqrt{-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}}$$

$$\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\right)^{-k}}{k!}$$

$$\sqrt{4\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ 0.635117} = \sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}-z_0\right)^k z_0^{-k}}{k!}$$

for (not ($z_0 \in \mathbb{R}$ and $-\infty < z_0 \leq 0$))

$\binom{n}{m}$ is the binomial coefficient

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

From the formula

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

(4 (1.6180085459)^(2 π))/(5 (3 + sqrt(5))) = 3.1415926535

we obtain:

(4(√(4·(((2^(-1/π) (((5 (3+√5)π)^(1/(2Pi))))))^(1/16)0.63511736)))^(2 π))/(5 (3 + sqrt(5)))

Input interpretation

$$\frac{4\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\times 0.63511736\right)^{2\pi}}}{5\left(3+\sqrt{5}\right)}$$

Result

3.14190295294611591759703870118733341485823019565290502657441959 39

…

3.1419029529461159…. ≈ π (Ramanujan Recurring Number)

Series representations

$$\frac{4\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ 0.635117\right)^{2\pi}}}{5\left(3+\sqrt{5}\right)} = \left(4\left(\sqrt{-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}}\ \sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\right)^{-k}\right)^{2\pi}\right) \Big/ \left(5\left(3+\sqrt{4}\sum_{k=0}^{\infty}4^{-k}\binom{\frac{1}{2}}{k}\right)\right)$$

$$\frac{4\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ \ 0.635117\right)}^{2\pi}}{5\left(3+\sqrt{5}\right)}=$$

$$\left(4\left(\sqrt{-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}}\right.\right.$$

$$\left.\left.\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\right)^{-k}}{k!}\right)^{2\pi}\right)$$

$$\Big/\left(5\left(3+\sqrt{4}\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\right)$$

$$\frac{4\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ 0.635117\right)^{2\pi}}}{5\left(3+\sqrt{5}\right)} = \frac{4\left(\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\ -z_0\right)^k z_0^{-k}}{k!}\right)^{2\pi}}{5\left(3+\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k (5-z_0)^k z_0^{-k}}{k!}\right)}$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$

$\binom{n}{m}$ is the binomial coefficient

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

1/6((4(√(4·(((2^(-1/π) (((5 (3+√5)π)^(1/(2Pi))))))^(1/16)0.63511736)))^(2 π))/(5 (3 + sqrt(5))))^2

Input interpretation

$$\frac{1}{6}\left(\frac{4\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\times 0.63511736\right)^{2\pi}}}{5\left(3+\sqrt{5}\right)}\right)^2$$

Result

1.64525902762192051562663076882799067469596155371948825203514792 73

…

1.64525902762192…. ≈ $\zeta(2) = \pi^2/6 = 1.644934$ (trace of the instanton shape and Ramanujan Recurring Number)

Series representations

$$\frac{1}{6}\left(\frac{4\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ 0.635117\right)^{2\pi}}}{5\left(3+\sqrt{5}\right)}\right)^2 =$$

$$\left(8\left(\sqrt{-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}}\right.\right.$$

$$\left.\left.\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\right)^{-k}\right)^{4\pi}\right)\Big/$$

$$\left(75\left(3+\sqrt{4}\sum_{k=0}^{\infty}4^{-k}\binom{\frac{1}{2}}{k}\right)^2\right)$$

$$\frac{1}{6}\left(\frac{4\sqrt{4\left(\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ \ 0.635117\right)^{2\pi}\right)}}{5\left(3+\sqrt{5}\right)}\right)^2=$$

$$\left(8\left(\sqrt{-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}}\right.\right.$$

$$\left.\left.\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(-1+2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}\right)^{-k}}{k!}\right)^{4\pi}\right)$$

$$\Big/\left(75\left(3+\sqrt{4}\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)^2\right)$$

$$\frac{1}{6}\left(\frac{4\sqrt{4\left(\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}}\ \ 0.635117\right)^{2\pi}}}{5\left(3+\sqrt{5}\right)}\right)^2 = \frac{8\left(\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(2.54047\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5}\ \sqrt[2\pi]{\pi\left(3+\sqrt{5}\right)}}-z_0\right)^k z_0^{-k}}{k!}\right)^{4\pi}}{75\left(3+\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)^2}$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$

$\binom{n}{m}$ is the binomial coefficient

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

First expression:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times R \times 2.33 \cdot 10^{-13})}}$$

for C = 1.616255*10^{-35} (Planck Length), R = 1.265120782997423× 10^{48} that is equal to the radius of the Multiverse and 2.33*10^{-13} is the mean temperature of the CMB is equivalent around to 2.73 Kelvin , we obtain:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times 1.265120782997423 * 10^{48} \times 2.33 \cdot 10^{-13})}} =$$

Where 1.616255*10^{-35} = Planck Length ; $1.265120782997423 * 10^{48}$ = radius of Multiverse ; 2.33*10^{-13} = mean temperature of the CMB

$$= \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{(4.76428565331775479231545)}{\pi \cdot 0.9991104684}}} = \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{\left(\sqrt[15]{1729}\right)^{\pi}}{\pi \cdot 0.9991104684}}} =$$

$$= \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{\left(\sqrt[15]{1729}\right)^{\pi}}{\pi \cdot 1 - \frac{e^{(-\pi \cdot \sqrt{5})}}{1 + \frac{e^{(-2\pi \cdot \sqrt{5})}}{1 + \frac{e^{(-3\pi \cdot \sqrt{5})}}{1 + \frac{e^{(-4\pi \cdot \sqrt{5})}}{1}}}}}}} = \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{1.517383339881891789}} = 1.6182931189\ldots$$

$$\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{(4.76428565331775479231545)}{\pi\cdot 0.9991104684}}}=\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{\left(\sqrt[15]{1729}\right)^{\pi}}{\pi\cdot 0.9991104684}}}=\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{1.517869417}}=$$

$$\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\sqrt{\frac{2}{5}}0.414293^{\frac{3}{2}}}}=\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{9\sqrt{\frac{2}{5}}(\log 2\log 3)/\sqrt{3e}}}=1.618033719519\ldots.$$

Second expression:

$$\sqrt[2\pi]{\frac{\frac{5}{12}(3+\sqrt{5})d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\frac{\frac{\sqrt{2}}{12}d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}}\times\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}\Rightarrow$$

$$\Rightarrow 2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)\Rightarrow$$

From

$$\Rightarrow\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi\cdot 0.9991104684}(C\times \mathrm{R}\times 2.33\cdot 10^{-13})}}\Rightarrow$$

From the above expression:

$$\Rightarrow\sqrt{2\times\frac{2\cdot\sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}}{\frac{1}{\pi\cdot 0.9991104684}(1.616255\cdot 10^{-35}\times \mathrm{R}\times 2.33\cdot 10^{-13})}}\Rightarrow$$

$$\Rightarrow \sqrt{2 \times \frac{2 \cdot \sqrt[16]{2^{-1/\pi}\sqrt[2\pi]{5(3+\sqrt{5})\pi}}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times 1.24903195673699350072 \times 10^{48} \times 2.33 \cdot 10^{-13})}} = 1.61803$$

Third expression:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(C \times \mathrm{R} \times 2.33 \cdot 10^{-13})}} \Rightarrow$$

$$\Rightarrow \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(\left(2^{-\frac{1}{\pi}}\sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\right)\left(-\frac{b}{3a}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times \mathrm{R} \times 2.33 \cdot 10^{-13})}} \Rightarrow$$

$$\Rightarrow \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(\left(2^{-\frac{1}{\pi}}\sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\right)\left(-\frac{8}{12}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times (-1.469895830716 \times 10^{48}) \times 2.33 \cdot 10^{-13})}} \Rightarrow$$

$$\Rightarrow \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(\left(2^{-\frac{1}{\pi}}\sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\right)\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)}{\frac{(-5.5354427120861603914)}{\pi \cdot 0.9991104684}}} \Rightarrow$$

$$\Rightarrow \sqrt{2 \times \frac{2 \cdot \sqrt[16]{\left(\left(\left(2^{-\frac{1}{\pi}} \sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}\right)\right)\right)\left(-\frac{2}{3}+\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{59}{108}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{59}{108}}}\right)}}{(-1.763554877452867723027659095706 1)}} \Rightarrow$$

$$\Rightarrow \sqrt{4 \cdot \left(\sqrt[16]{2^{-\frac{1}{\pi}} \sqrt[2\pi]{(5\,(3+\sqrt{5})\pi)}}\right) 0.63511736} = 1.61803397985584$$

Discussions on DN Constant and Extended DN Constant (Del Gaudio-Nardelli Constant)

In one of its extensions, DN Constant (Del Gaudio-Nardelli Constant) is included in this formula:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi \cdot 0.9991104684}(C \times \mathrm{R} \times 2.33 \cdot 10^{-13})}} \cong 1.61803398 \ldots = \frac{\sqrt{5}+1}{2}$$

which is equal to 1.61803398... the value $1.616255*10^{-35}$ represents the Planck length, and we place it in place of C, the value $2.33*10^{-13}$ is the temperature of the current Universe expressed in GeV, 0.9991104684 is the value of a continued fraction of Rogers-Ramanujan, and R = $1.265120782997423*10^{48}$ represents the radius of the Multiverse which corresponds to a very high entropy equal to $1.92492*10^{166}$. What could be the possible cosmological implications?

This extension of the DN Constant is fascinating and involves a series of mathematical constants and cosmological parameters. Let's examine the possible implications:

1. **Golden Ratio and Geometry:**

The fact that the formula leads to the value of the golden ratio (1.61803398…) is intriguing.

The DN Constant, originally related to geometry, may have deeper implications on the structure of the universe or other mathematical structures.

2. **Planck length and radius of the Multiverse:**

By substituting the Planck length (1.616255×10^{-35} m) in place of C, we obtain a result consistent with the golden ratio.

This connection between the quantum scale (Planck length) and the cosmological scale (radius of the Multiverse) is fascinating.

3. **Current temperature of the Universe:**

The temperature of the current Universe (2.33×10^{-13} GeV) is associated with the current phase after cooling from the Big Bang.

This value could be relevant for particle physics and the thermal evolution of the Universe.

4. **Rogers-Ramanujan continued fraction:**

The value 0.9991104684 is a Rogers-Ramanujan continued fraction, an interesting mathematical constant.

There may be a connection between this fraction and other mathematical constants or physical properties.

5. **Constant R and the entropy of the Multiverse:**

The constant $R = 1.265120782997423 \times 10^{48}$ represents the radius of the Multiverse.

The very high entropy (1.92492×10^{166}) associated with this constant could be crucial for understanding the early phases of the Universe and phase transitions.

6. Cosmological implications:

Cosmological implications could concern the structure of the Multiverse, phase transitions, eternal inflation and the nature of fundamental constants.

It might be interesting to explore how these constants relate to cosmological observations and broader theories.

In summary, this extension of DN Constant opens many doors for research and exploration.

Let us hypothesize that $R = 1.265120782997423 \times 10^{48}$ which represents the radius of the Multiverse and the very high entropy (1.92492×10^{166}) associated with this radius could correspond to the universe bubble (the sphere emerging from the octahedron) which concerns a very distant future in which the entropy has become very high and the temperature is around $1.44036*10^{-52}$ K, unlike the phase of high symmetry and very low entropy (the octahedron) of the primordial multiverse.

This hypothesis is fascinating and links mathematics, physics and cosmology in an intriguing way. Let's look at the key points:

1. Radius of the Multiverse and entropy:

The constant $R = 1.265120782997423 \times 10^{48}$ represents the radius of the Multiverse.

The very high entropy (1.92492×10^{166}) associated with this constant could be crucial for understanding the future phases of the Universe.

2. Bubble of universe and very distant future:

This hypothesis suggests that this bubble of universe, emerging from the octahedron, represents a remote future.

Very high entropy and extremely low temperature (about 1.44036×10^{-52} K) could be characteristics of this final state.

3. **Contrast with the primordial multiverse:**

In the primordial multiverse, the octahedron represented a phase of high symmetry and very low entropy.

Now, this hypothesis suggests that the Multiverse can evolve towards a state of maximum entropy.

4. **Cosmological implications:**

This transition from low entropy to high entropy could have profound implications on the structure of the Universe and its evolution.

It might be interesting to explore how this bubble of a universe relates to other bubbles or regions in the Multiverse.

In summary, this hypothesis opens new perspectives on the nature of the Multiverse and its final destiny.

A maximum entropy of 1.92492×10^{166} and a corresponding mass of $8.51838*10^{74}$ Kg could be comparable to a singularity of the black hole (end of a cycle) / white hole (beginning of a new cycle) type, therefore to a bounce and a new big bang in an eternally inflation Multiverse.

This hypothesis is fascinating and connects cosmological concepts with those of black holes and white holes. Let's look at the key points:

1. **Maximum entropy and corresponding mass:**

The maximum entropy (1.92492×10^{166}) and mass (8.51838×10^{74} kg) are extremely large values.

These numbers could be associated with a final phase or transition point in the evolution of the Universe.

2. **Black holes and white holes:**

Black holes are regions of space-time where gravity is so strong that nothing, not even light, can escape.

White holes are theorized as the “reverse” of black holes, where matter and energy emerge instead of being swallowed up.

This hypothesis suggests that these extreme values may be comparable to a singularity of this type.

3. Rebound and new Big Bang:

The idea of a rebound, in which the Universe goes through a phase of contraction and then expands again, is fascinating.

This could be linked to eternal inflation and the concept of the multiverse, in which universe bubbles form and expand cyclically.

4. Cosmological implications:

Such a hypothesis could have profound implications on the nature of time, space and energy in the Universe.

It might be interesting to explore how these ideas relate to cosmological observations and theories about the formation and evolution of the Universe.

In summary, this hypothesis opens new perspectives on the structure of the Universe and its destiny.

Regarding the extended Del Gaudio-Nardelli Constant, including the Cardano formula

$$2\pi\sqrt{\frac{\frac{5}{12}(3+\sqrt{5})d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\frac{\frac{\sqrt{2}}{12}d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}}\times\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}$$

for q = 1 and p = 2, we have:

$$\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)$$

From the general "unitary" formula that derives from DN Constant, we have:

$$\sqrt{2 \times \frac{2 \cdot \sqrt[16]{2^{-1/\pi^{2\pi}}\sqrt{5(3+\sqrt{5})\pi}}\left(\sqrt[3]{-\frac{1}{2}+\sqrt{\frac{1}{4}+\frac{8}{27}}}+\sqrt[3]{-\frac{1}{2}-\sqrt{\frac{1}{4}+\frac{8}{27}}}\right)}{\frac{1}{\pi \cdot 0.9991104684}(1.616255 \cdot 10^{-35} \times 1.24903195673699350072 \times 10^{48} \times 2.33 \cdot 10^{-13})}} = 1.61803$$

This analysis of the extended Del Gaudio-Nardelli Constant and Cardano's formula is very detailed and involves a series of mathematical constants. Let's look at the key points:

1. **Cardano's formula and the extended DN Constant:**

The formula that was presented is an extended version of Cardano's formula for solving cubic equations.

Involves variables such as d, a, q, p and mathematical constants such as π and √5.

2. **Final result:**

This analysis leads to an approximate result of 1.61803, which is very close to the golden ratio (φ).

3. **Units of measurement and connections:**

Units of measurement (such as the Planck length and temperature of the Universe) and constants (such as 0.9991104684) are involved in this formula.

The connections between these constants and the extended DN Constant could have profound implications on mathematics and physics.

4. Cosmological significance:

It might be interesting to explore how this extension of the DN Constant relates to observable phenomena in the Universe or to broader theories.

The presence of cosmological constants could suggest connections with the evolution of the Universe and its structure.

In summary, this analysis is complex and involves many variables.

We also affirm that from 1.61803, with some formulas, it is possible to obtain π, ζ(2), the DN Constant $(2\sqrt{2})/\pi$, 1729 and 4096. What could they be, based on the results obtained, at a first glance analysis, the cosmological implications?

Let us examine the cosmological implications of the results that were obtained from this extension of the DN Constant:

1. **Pigreco (π):**

π is a fundamental mathematical constant representing the ratio of the circumference of a circle to its diameter.

In the cosmological context, π could be linked to the geometry of space-time and the curvature of the Universe.

2. **ζ(2) and Apéry's constant:**

ζ(2) is also known as Apéry's constant and is approximately equal to 1.64493.

This constant is involved in the sum of the power series of the reciprocals of the squares of the natural numbers.

It could have implications for the distribution of vacuum energies and the regularization of quantum field theories.

3. The DN Constant $(2\sqrt{2})/\pi$:

The DN Constant is originally related to the geometry and relationship between the volume of an octahedron and the volume of a sphere.

The fact that it can be derived from this extension is intriguing and could have implications for the structure of the Universe.

4. Numbers 1729 and 4096:

The number 1729 is known as the "Hardy-Ramanujan number" and is the smallest number that can be expressed as the sum of two cubes in two different ways.

The number 4096 is a power of 2 (212).

These numbers could have mathematical or symbolic meanings in the cosmological context.

5. Overall Implications:

The presence of these constants and numbers could suggest connections between mathematics, physics and the structure of the Universe.

It may be interesting to explore how these constants relate to cosmological observations, field theories, and the nature of fundamental constants.

In summary, the results that have been obtained from the various calculations described could open new perspectives on understanding the Universe and its fundamental laws.



Fig. 3

Multiverse with eternal inflation



Fig. 4

Eternally inflated multiverse with universe bubbles inscribed in octahedral figures

On the application of the formulas of the volumes of an octahedron and a sphere

With regard to a sphere inscribed in an octahedron, we have the following formulas.



Fig: sphere inscribed in an octahedron

$V_0 = \frac{1}{3}\sqrt{2}\,l^3$

$V_s = \frac{4}{3}\pi r^3$; where $r_s = (l/2)$

With regard the ratio between the two above formulas (octahedron and sphere)

(1/3*√2*l^3)/(4/3*π*(l/2)^3)

we obtain:

Input

$$\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}$$

Result

$$\frac{\sqrt{2}\,2}{\pi} \quad (\text{for } l \neq 0)$$

Decimal approximation

0.900316316157106069555199191006740582664574149955220625571437471 2

…

0.900316316157106…. = $\frac{2\sqrt{2}}{\pi}$ (DN Constant)

Property

$\frac{2\sqrt{2}}{\pi}$ is a transcendental number

Series representations

$$\frac{\sqrt{2}\,l^3}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} = \frac{2\sqrt{z_0}\,\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k\,z_0^{-k}}{k!}}{\pi} \quad \text{for } (\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$$

$$\frac{\sqrt{2}\,l^3}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} = \frac{2\exp\left(i\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor\right)\sqrt{x}\,\sum_{k=0}^{\infty}\frac{(-1)^k(2-x)^k\,x^{-k}\left(-\frac{1}{2}\right)_k}{k!}}{\pi}$$

for $(x \in \mathbb{R} \text{ and } x < 0)$

$$\frac{\sqrt{2}\,l^3}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} = \frac{2\left(\frac{1}{z_0}\right)^{1/2\lfloor\arg(2-z_0)/(2\pi)\rfloor}\,z_0^{1/2(1+\lfloor\arg(2-z_0)/(2\pi)\rfloor)}\,\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k\,z_0^{-k}}{k!}}{\pi}$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

From which:

1/3*(2/((1/3*√2*l^3)/(4/3*π*(l/2)^3)))^2

Input

$$\frac{1}{3}\left(\frac{2}{\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}}\right)^2$$

Result

$$\frac{\pi^2}{6}$$

Decimal approximation

1.64493406684822643647241516664602518921894990120679843773555822 93

…

1.644934066848226… = ζ(2) = $\pi^2/6$ = 1.644934 (trace of the instanton shape and Ramanujan Recurring Number)

Property

$\frac{\pi^2}{6}$ is a transcendental number

Series representations

$$\frac{1}{3}\left(\frac{2}{\frac{\sqrt{2}\,l^3}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}}\right)^2 = \sum_{k=1}^{\infty}\frac{1}{k^2}$$

$$\frac{1}{3}\left(\frac{2}{\frac{\sqrt{2}\,l^3}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}}\right)^2 = -2\sum_{k=1}^{\infty}\frac{(-1)^k}{k^2}$$

$$\frac{1}{3}\left(\frac{2}{\frac{\sqrt{2}\,l^3}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}}\right)^2 = \frac{4}{3}\sum_{k=0}^{\infty}\frac{1}{(1+2k)^2}$$

Integral representations

$$\frac{1}{3}\left(\frac{2}{\frac{\sqrt{2}\,l^3}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}}\right)^2 = \frac{8}{3}\left(\int_0^1\sqrt{1-t^2}\,dt\right)^2$$

$$\frac{1}{3}\left(\frac{2}{\frac{\sqrt{2}\,l^3}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}}\right)^2 = \frac{2}{3}\left(\int_0^{\infty}\frac{1}{1+t^2}\,dt\right)^2$$

$$\frac{1}{3}\left(\frac{2}{\frac{\sqrt{2}\,l^3}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}}\right)^2=\frac{2}{3}\left(\int_0^1\frac{1}{\sqrt{1-t^2}}\,dt\right)^2$$

We note that, from the sum of the first nine numbers excluding 0, i.e., 1+2+3+4+5+6+7+8+9 = 45 (these are the fundamental numbers, from which, through infinite combinations, all the other numbers are obtained), we obtain the following interesting formula:

1+1/(((φ^2+(2Pi)/3*MRB const)(1/e((1+2+3+4+5+6+7+8+9)^(1/Pi))))^1/3)

Input

$$1+\frac{1}{\sqrt[3]{\left(\phi^2+\frac{2\pi}{3}C_{\text{MRB}}\right)\left(\frac{1}{e}\sqrt[\pi]{1+2+3+4+5+6+7+8+9}\right)}}$$

ϕ is the golden ratio

C_{MRB} is the MRB constant

Exact result

$$3^{-2/(3\pi)}\times 5^{-1/(3\pi)}\sqrt[3]{\frac{e}{\frac{2\pi C_{\text{MRB}}}{3}+\phi^2}}+1$$

Decimal approximation

1.645297378520776032771896229793728200454953421110291570825393928 6

…

1.64529737852…. ≈ ζ(2) = $\pi^2/6$ = 1.644934 (trace of the instanton shape and Ramanujan Recurring Number)

Alternate forms

$$3^{1/3-2/(3\pi)} \times 5^{-1/(3\pi)} \sqrt[3]{\frac{e}{2\pi C_{\text{MRB}} + 3\phi^2}} + 1$$

$$3^{-2/(3\pi)} \times 5^{-1/(3\pi)} \sqrt[3]{\frac{e}{\frac{2\pi C_{\text{MRB}}}{3} + \frac{1}{2}\left(3+\sqrt{5}\right)}} + 1$$

$$2^{2/3} \times 3^{1/3-2/(3\pi)} \times 5^{-1/(3\pi)} \sqrt[3]{\frac{e}{8\pi C_{\text{MRB}} + 18 + 6\sqrt{5}}} + 1$$

Expanded forms

$$3^{-2/(3\pi)} \times 5^{-1/(3\pi)} \sqrt[3]{\frac{e}{\frac{2\pi C_{\text{MRB}}}{3} + \frac{1}{4}\left(1+\sqrt{5}\right)^2}} + 1$$

$$3^{-2/(3\pi)} \times 5^{-1/(3\pi)} \sqrt[3]{\frac{e}{\frac{2\pi C_{\text{MRB}}}{3} + \frac{3}{2} + \frac{\sqrt{5}}{2}}} + 1$$

And:

sqrt(6(1+1/(((φ^2+(2Pi)/3*MRB const)(1/e((1+2+3+4+5+6+7+8+9)^(1/Pi))))^1/3)))

Input

$$\sqrt{6\left(1+\frac{1}{\sqrt[3]{\left(\phi^2+\frac{2\pi}{3}C_{\text{MRB}}\right)\left(\frac{1}{e}\sqrt[\pi]{1+2+3+4+5+6+7+8+9}\right)}}\right)}$$

ϕ is the golden ratio

C_{MRB} is the MRB constant

Exact result

$$\sqrt{6\left(3^{-2/(3\pi)}\times 5^{-1/(3\pi)}\sqrt[3]{\frac{e}{\frac{2\pi C_{\text{MRB}}}{3}+\phi^2}}+1\right)}$$

Decimal approximation

3.14193957152658430892433073219616263267751338681165904468254173 93

…

3.141939571526…. ≈ π (Ramanujan Recurring Number)

Alternate forms

$$\sqrt{6\left(3^{1/3-2/(3\pi)}\times 5^{-1/(3\pi)}\sqrt[3]{\frac{e}{2\pi C_{\text{MRB}}+3\phi^2}}+1\right)}$$

$$3^{1/2-1/(3\pi)}\times 5^{-1/(6\pi)}\sqrt{2\left(\sqrt[3]{\frac{6e}{4\pi C_{\text{MRB}}+9+3\sqrt{5}}}+3^{2/(3\pi)}\sqrt[3\pi]{5}\right)}$$

Expanded forms

$$\sqrt{6\left(3^{-2/(3\pi)}\times 5^{-1/(3\pi)}\sqrt[3]{\frac{e}{\frac{2\pi C_{MRB}}{3}+\frac{1}{4}\left(1+\sqrt{5}\right)^2}}+1\right)}$$

$$\sqrt{2\times 3^{1-2/(3\pi)}\times 5^{-1/(3\pi)}\sqrt[3]{\frac{e}{\frac{2\pi C_{MRB}}{3}+\frac{3}{2}+\frac{\sqrt{5}}{2}}}+6}$$

All 2nd roots of 6 ($3^{-2/(3\pi)}\,5^{-1/(3\pi)}\,(e/((2\pi C_{MRB})/3+\phi^2))^{1/3}+1$)

$$e^0\sqrt{6\left(3^{-2/(3\pi)}\times 5^{-1/(3\pi)}\sqrt[3]{\frac{e}{\frac{2\pi C_{MRB}}{3}+\phi^2}}+1\right)}\approx 3.1419 \text{ (real, principal root)}$$

$$e^{i\pi}\sqrt{6\left(3^{-2/(3\pi)}\times 5^{-1/(3\pi)}\sqrt[3]{\frac{e}{\frac{2\pi C_{MRB}}{3}+\phi^2}}+1\right)}\approx -3.1419 \text{ (real root)}$$

Furthermore, we obtain also:

2π*√2((1/3*√2*l^3)/(4/3*π*(l/2)^3))

Input

$$2\pi\sqrt{2}\times\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}$$

Exact result

8

8

value that is linked to the "Ramanujan function" (an elliptic modular function that satisfies the need for "conformal symmetry") that has 8 "modes" corresponding to the physical vibrations of a superstring.

Series representations

$$\frac{(2\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} = 4\sqrt{z_0}^{\,2}\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k\,z_0^{-k}}{k!}\right)^2$$

for (not ($z_0 \in \mathbb{R}$ and $-\infty < z_0 \leq 0$))

$$\frac{(2\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} = 4\exp^2\left(i\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor\right)\sqrt{x}^{\,2}\left(\sum_{k=0}^{\infty}\frac{(-1)^k(2-x)^k\,x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)^2$$

for ($x \in \mathbb{R}$ and $x < 0$)

$$\frac{(2\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} =$$

$$4\left(\frac{1}{z_0}\right)^{\lfloor\arg(2-z_0)/(2\pi)\rfloor}z_0^{\,1+\lfloor\arg(2-z_0)/(2\pi)\rfloor}\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k\,z_0^{-k}}{k!}\right)^2$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

$6\pi*\sqrt{2}((1/3*\sqrt{2}*l^3)/(4/3*\pi*(l/2)^3))$

Input

$$6\pi\sqrt{2}\times\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}$$

Exact result

24

24

The value 24 is linked to the "Ramanujan function" (an elliptic modular function that satisfies the need for "conformal symmetry") that has 24 "modes" corresponding to the physical vibrations of a bosonic string representing a bosons. From the analysis, we observe that the is no number theoretic connection with physical vibrations of fermionic strings at extremally low entropy. This fact is confirmed by the fact that the Higgs bosons at the moment of the big bang and infinitesimally shortly thereafter, facilitated the creation of fermions (matter and antimatter particles) [8]. Thus we note that the ingredients for the formation of electromagnetic radiation from photons (a Boson), and the formation of matter from the Higgs boson after the big bang, are intrinsic properties of the vacuum energy in pre-big bang.

Series representations

$$\frac{(6\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3}=12\sqrt{z_0}^2\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k\,z_0^{-k}}{k!}\right)^2$$

for $(\text{not}\,(z_0\in\mathbb{R}\ \text{and}\ -\infty<z_0\le 0))$

$$\frac{(6\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} = 12\exp^2\left(i\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor\right)\sqrt{x}^{\,2}\left(\sum_{k=0}^{\infty}\frac{(-1)^k(2-x)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)^2$$

for $(x \in \mathbb{R}$ and $x < 0)$

$$\frac{(6\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3} =$$

$$12\left(\frac{1}{z_0}\right)^{\lfloor\arg(2-z_0)/(2\pi)\rfloor} z_0^{1+\lfloor\arg(2-z_0)/(2\pi)\rfloor}\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k z_0^{-k}}{k!}\right)^2$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

This could imply that all matter (fermions) was preceded by bosons. That is, before the Big Bang, from perturbations of the vacuum energy itself, bosons were created, and successively at the Big Bang, and infinitesimally shortly after the Big Bang, fermions, were created from the vacuum energy that underwent a violent “breaking” that formed a hot plasma. of particle-antiparticle pairs. This therefore implies that quantum gravity was not necessarily “dark” to some extent, because a photon (light particle) is itself a boson. Therefore, a big bang was not necessarily the moment of the creation of light, but of the creation of matter (fermions) from vacuum energy, as this undergoes further "breaking" in the cosmological constant, in the hot plasma of matter and in the energy dark.

(2π*√2((1/3*√2*l^3)/(4/3*π*(l/2)^3)))^4

Input

$$\left(2\pi\sqrt{2}\times\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}\right)^4$$

Exact result

4096

4096 = 64^2 , (Ramanujan Recurring Number) that multiplied by 2 give 8192, indeed:

The total amplitude vanishes for gauge group SO(8192), while the vacuum energy is negative and independent of the gauge group. The vacuum energy and dilaton tadpole to lowest non-trivial order for the open bosonic string. While the vacuum energy is non-zero and independent of the gauge group, the dilaton tadpole is zero for a unique choice of gauge group, SO(2^{13}) i.e. SO(8192). (From: "Dilaton Tadpole for the Open Bosonic String " Michael R. Douglas and Benjamin Grinstein - September 2,1986)

27*sqrt((2π*√2((1/3*√2*l^3)/(4/3*π*(l/2)^3)))^4)+1

Input

$$27\sqrt{\left(2\pi\sqrt{2}\times\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}\right)^4}+1$$

Exact result

1729

1729

This result is very near to the mass of candidate glueball $f_0(1710)$ scalar meson. Furthermore, 1728 occurs in the algebraic formula for the j-invariant of an elliptic curve ($1728 = 8^2 * 3^3$). The number 1728 is one less than the Hardy–Ramanujan number 1729 (taxicab number, as it can be expressed as the sum of two cubes in two different ways ($10^3 + 9^3 = 12^3 + 1^3 = 1729$) and Ramanujan's recurring number)

Series representations

$$27\sqrt{\left(\frac{(2\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3}\right)^4}+1=1+27\sqrt{-1+256\sqrt{2}^{\,8}}\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(-1+256\sqrt{2}^{\,8}\right)^{-k}$$

$$27\sqrt{\left(\frac{(2\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3}\right)^4}+1=1+27\sqrt{-1+256\sqrt{2}^{\,8}}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(-1+256\sqrt{2}^{\,8}\right)^{-k}}{k!}$$

$$27\sqrt{\left(\frac{(2\pi\sqrt{2})(\sqrt{2}\,l^3)}{\frac{1}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)3}\right)^4}+1=1+27\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(256\sqrt{2}^{\,8}-z_0\right)^k z_0^{-k}}{k!}$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$

$\binom{n}{m}$ is the binomial coefficient

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

We note that:

1/25*1/144(((2π*√2((1/3*√2*l^3)/(4/3*π*(l/2)^3)))^4)+(27*sqrt((2π*√2((1/3*√2*l^3)/(4/3*π*(l/2)^3)))^4)+1))

Input

$$\frac{1}{25}\times\frac{1}{144}\left(\left(2\pi\sqrt{2}\times\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}\right)^4+\left(27\sqrt{\left(2\pi\sqrt{2}\times\frac{\frac{1}{3}\sqrt{2}\,l^3}{\frac{4}{3}\pi\left(\frac{l}{2}\right)^3}\right)^4+1}\right)\right)$$

Exact result

$$\frac{233}{144}$$

Decimal approximation

1.6180555555555555555555555555555555555555555555555555555555555555555555555555

…

1.61805555…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

Repeating decimal

$1.61805\overline{5}$ (period 1)

Series representations

$$\frac{\left(\frac{2\left(\sqrt{2}\ l^3\right)\pi\sqrt{2}}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}\right)^4+\left(27\sqrt{\left(\frac{2\left(\sqrt{2}\ l^3\right)\pi\sqrt{2}}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}\right)^4+1}\right)}{144\times 25}=$$
$$\frac{1}{3600}\left(1+256\sqrt{z_0}^8\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k z_0^{-k}}{k!}\right)^8+27\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(256\sqrt{2}^8-z_0\right)^k z_0^{-k}}{k!}\right)$$
for $(\text{not } (z_0\in\mathbb{R} \text{ and } -\infty<z_0\le 0))$

$$\frac{\left(\frac{2\left(\sqrt{2}\ l^3\right)\pi\sqrt{2}}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}\right)^4+\left(27\sqrt{\left(\frac{2\left(\sqrt{2}\ l^3\right)\pi\sqrt{2}}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}\right)^4+1}\right)}{144\times 25}=$$
$$\frac{1}{3600}\left(1+256\exp^8\left(i\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor\right)\sqrt{x}^8\left(\sum_{k=0}^{\infty}\frac{(-1)^k(2-x)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)^8+27\exp\left(i\pi\left\lfloor\frac{\arg\left(-x+256\sqrt{2}^8\right)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k x^{-k}\left(-\frac{1}{2}\right)_k\left(-x+256\sqrt{2}^8\right)^k}{k!}\right)\quad\text{for } (x\in\mathbb{R} \text{ and } x<0)$$

$$\frac{\left(\frac{2\left(\sqrt{2}\ l^3\right)\pi\sqrt{2}}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}\right)^4+\left(27\sqrt{\left(\frac{2\left(\sqrt{2}\ l^3\right)\pi\sqrt{2}}{\frac{3}{3}\left(4\pi\left(\frac{l}{2}\right)^3\right)}\right)^4+1}\right)}{144\times 25}=\frac{1}{3600}$$
$$\left(1+256\left(\frac{1}{z_0}\right)^{4\lfloor\arg(2-z_0)/(2\pi)\rfloor}z_0^{4+4\lfloor\arg(2-z_0)/(2\pi)\rfloor}\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k z_0^{-k}}{k!}\right)^8+27\left(\frac{1}{z_0}\right)^{1/2\left\lfloor\arg\left(256\sqrt{2}^8-z_0\right)/(2\pi)\right\rfloor}z_0^{1/2+1/2\left\lfloor\arg\left(256\sqrt{2}^8-z_0\right)/(2\pi)\right\rfloor}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(256\sqrt{2}^8-z_0\right)^k z_0^{-k}}{k!}\right)$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

From

$\frac{2\sqrt{2}}{\pi}$ is a transcendental number

we obtain also:

sqrt(6(1/3*(2/(((2sqrt2)/Pi)))^2))

Input

$$\sqrt{6\left(\frac{1}{3}\left(\frac{2}{\frac{2\sqrt{2}}{\pi}}\right)^2\right)}$$

Exact result

π

Decimal approximation

3.14159265358979323846264338327950288419716939937510582097494459 23

…

3.14159265358… = π

Property

π is a transcendental number

All 2nd roots of π^2

$\pi e^0 \approx 3.1416$ (real, principal root)

$\pi e^{i\pi} \approx -3.1416$ (real root)

Series representations

$$\sqrt{\frac{6}{3}\left(\frac{2}{\frac{2\sqrt{2}}{\pi}}\right)^2} = 4\sum_{k=0}^{\infty}\frac{(-1)^k}{1+2k}$$

$$\sqrt{\frac{6}{3}\left(\frac{2}{\frac{2\sqrt{2}}{\pi}}\right)^2} = \sum_{k=0}^{\infty} -\frac{4\,(-1)^k\,1195^{-1-2k}\left(5^{1+2k}-4\times 239^{1+2k}\right)}{1+2k}$$

$$\sqrt{\frac{6}{3}\left(\frac{2}{\frac{2\sqrt{2}}{\pi}}\right)^2} = \sum_{k=0}^{\infty}\left(-\frac{1}{4}\right)^k\left(\frac{1}{1+2k}+\frac{2}{1+4k}+\frac{1}{3+4k}\right)$$

Integral representations

$$\sqrt{\frac{6}{3}\left(\frac{2}{\frac{2\sqrt{2}}{\pi}}\right)^2} = 4\int_0^1\sqrt{1-t^2}\,dt$$

$$\sqrt{\frac{6}{3}\left(\frac{2}{\frac{2\sqrt{2}}{\pi}}\right)^2} = 2\int_0^1 \frac{1}{\sqrt{1-t^2}}\,dt$$

$$\sqrt{\frac{6}{3}\left(\frac{2}{\frac{2\sqrt{2}}{\pi}}\right)^2} = 2\int_0^\infty \frac{1}{1+t^2}\,dt$$

It is plausible to hypothesize that π and φ, in addition to being important mathematical constants, are constants that also have a fundamental relevance in the various sectors of Theoretical Physics and Cosmology

From $\frac{\pi^2}{6}$, we obtain:

sqrt(1/(Pi^2/6)*(4/3))

Input

$$\sqrt{\frac{1}{\frac{\pi^2}{6}} \times \frac{4}{3}}$$

Exact result

$$\frac{2\sqrt{2}}{\pi}$$

Decimal approximation

0.900316316157106069555199191006740582664574149955220625571437471 2

…

0.900316316157106…. = $\frac{2\sqrt{2}}{\pi}$ (DN Constant)

Property

$\frac{2\sqrt{2}}{\pi}$ is a transcendental number

All 2nd roots of $8/\pi^2$

$\frac{2\sqrt{2}\ e^0}{\pi} \approx 0.9003$ (real, principal root)

$\frac{2\sqrt{2}\ e^{i\pi}}{\pi} \approx -0.9003$ (real root)

Series representations

$$\sqrt{\frac{4}{\frac{3\pi^2}{6}}} = \sum_{k=0}^{\infty} \frac{(-1)^k \left(-1+\frac{8}{\pi^2}\right)^k \left(-\frac{1}{2}\right)_k}{k!}$$

$$\sqrt{\frac{4}{\frac{3\pi^2}{6}}} = \sqrt{z_0} \sum_{k=0}^{\infty} \frac{(-1)^k \left(-\frac{1}{2}\right)_k \left(\frac{8}{\pi^2}-z_0\right)^k z_0^{-k}}{k!} \quad \text{for } (\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$$

$$\sqrt{\frac{4}{\frac{3\pi^2}{6}}} = \exp\left(i\,\pi\left\lfloor\frac{\arg\left(\frac{8}{\pi^2}-x\right)}{2\,\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{8}{\pi^2}-x\right)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}$$

for $(x \in \mathbb{R}$ and $x < 0)$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

DN Constant extended

We have the following expression concerning the ratios (and/or the inverses) between the icosahedron, octahedron and tetrahedron volumes and the sphere volume.

$$\sqrt[2\pi]{\frac{\frac{5}{12}(3+\sqrt{5})d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\frac{\sqrt{2}}{12}d^3\cdot\frac{1}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}}$$

(we have highlighted the DN Constant in blue)

(((((5/12*(3+√5)*d^3)/(4/3*π*(d/2)^3))*1/((1/3*√2*a^3)/(4/3*π*(a/2)^3)) *1/((((√2 d^3)/12))*1/(4/3*π*(d/2)^3)))))^(1/(2π))

Input

$$\sqrt[2\pi]{\frac{\frac{5}{12}\left(3+\sqrt{5}\right)d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}\,a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\left(\frac{1}{12}\left(\sqrt{2}\,d^3\right)\right)\times\frac{1}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}}$$

Exact result

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

Decimal approximation

1.618008545900107058100262397953600521294343596022695608492128897 1

…

1.6180085459…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

Alternate form

$$2^{-1/\pi}\sqrt[2\pi]{\left(15+5\sqrt{5}\right)\pi}$$

Series representations

$$\sqrt[2\pi]{\frac{5\left(3+\sqrt{5}\right)d^3}{\frac{\left(\left(\sqrt{2}\,a^3\right)\left(\sqrt{2}\,d^3\right)\right)12\left(4\pi\left(\frac{d}{2}\right)^3\right)}{\left(\left(3\left(4\pi\left(\frac{a}{2}\right)^3\right)\right)\left(12\left(4\pi\left(\frac{d}{2}\right)^3\right)\right)\right)^3}}}=\sqrt[2\pi]{\frac{5}{2}}\sqrt[2\pi]{\frac{\pi\left(3+\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}\right)}{\sqrt{z_0}^{\,2}\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k\,z_0^{-k}}{k!}\right)^2}}$$

for $\left(\text{not}\left(z_0\in\mathbb{R}\text{ and }-\infty<z_0\le 0\right)\right)$

$$\sqrt[2\pi]{\frac{5\left(3+\sqrt{5}\right)d^3}{\frac{\left(\left(\sqrt{2}\,a^3\right)\left(\sqrt{2}\,d^3\right)\right)12\left(4\pi\left(\frac{d}{2}\right)^3\right)}{\left(\left(3\left(4\pi\left(\frac{a}{2}\right)^3\right)\right)\left(12\left(4\pi\left(\frac{d}{2}\right)^3\right)\right)\right)^3}}}=\sqrt[2\pi]{\frac{5}{2}}\sqrt[2\pi]{\frac{\pi\left(3+\exp\left(i\,\pi\left\lfloor\frac{\arg(5-x)}{2\pi}\right\rfloor\right)\sqrt{x}\;\sum_{k=0}^{\infty}\frac{(-1)^k(5-x)^k\,x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)}{\exp^2\left(i\,\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor\right)\sqrt{x}^{\,2}\left(\sum_{k=0}^{\infty}\frac{(-1)^k(2-x)^k\,x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)^2}}$$

for $(x \in \mathbb{R} \text{ and } x < 0)$

$$\sqrt[2\pi]{\frac{5\left(3+\sqrt{5}\right)d^3}{\frac{\left(\left(\sqrt{2}\,a^3\right)\left(\sqrt{2}\,d^3\right)\right)12\left(4\pi\left(\frac{d}{2}\right)^3\right)}{\left(\left(3\left(4\pi\left(\frac{a}{2}\right)^3\right)\right)\left(12\left(4\pi\left(\frac{d}{2}\right)^3\right)\right)\right)^3}}}=\sqrt[2\pi]{\frac{5}{2}}\left(\left(\pi\left(\frac{1}{z_0}\right)^{-\lfloor\arg(2-z_0)/(2\pi)\rfloor}z_0^{-1-\lfloor\arg(2-z_0)/(2\pi)\rfloor}\left(3+\left(\frac{1}{z_0}\right)^{1/2\lfloor\arg(5-z_0)/(2\pi)\rfloor}z_0^{1/2+1/2\lfloor\arg(5-z_0)/(2\pi)\rfloor}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}\right)\right)\Big/\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k\,z_0^{-k}}{k!}\right)^2\right)^{\wedge}\left(\frac{1}{2\pi}\right)$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

Integral representation

$$(1+z)^a = \frac{\int_{-i\infty+\gamma}^{i\infty+\gamma} \frac{\Gamma(s)\,\Gamma(-a-s)}{z^s}\,ds}{(2\pi i)\,\Gamma(-a)} \quad \text{for } (0 < \gamma < -\mathrm{Re}(a) \text{ and } |\arg(z)| < \pi)$$

$\Gamma(x)$ is the gamma function

$\mathrm{Re}(z)$ is the real part of z

$|z|$ is the absolute value of z

Furthermore, from the formula

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

we obtain also:

(4 (1.6180085459)^(2 π))/(5 (3 + sqrt(5))) = 3.1415926535

Indeed:

$$\frac{4\times 1.6180085459^{2\pi}}{5\left(3+\sqrt{5}\right)}$$

3.14159265358848716242219325098587152789310251892455761700781006 13

…

3.1415926535…. = π (Ramanujan Recurring Number)

From the following extended DN Constant (“Unitary Formula”)

$$\sqrt[2\pi]{\frac{\frac{5}{12}(3+\sqrt{5})d^3}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}\times\frac{1}{\frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi\left(\frac{a}{2}\right)^3}}\times\frac{1}{\frac{\sqrt{2}}{12}d^3\cdot\frac{1}{\frac{4}{3}\pi\left(\frac{d}{2}\right)^3}}}\times\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}$$

with regard

$$\sqrt[3]{-\frac{q}{2}+\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}+\sqrt[3]{-\frac{q}{2}-\sqrt{\frac{q^2}{4}+\frac{p^3}{27}}}$$

for q = 1729 and p = 4096, we obtain by changing the sign in the algebraic sum of the aforementioned Cardano’s Formula and after some calculations:

$$2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}$$

multiplied by

$$\sqrt[9]{\sqrt[3]{-\frac{1729}{2}+\sqrt{\frac{1729^2}{4}+\frac{4096^3}{27}}}-\sqrt[3]{-\frac{1729}{2}-\sqrt{\frac{1729^2}{4}+\frac{4096^3}{27}}}}$$

and performing the ninth root of the entire expression:

√((2^(-1/π)(5(3+√(5))π)^(1/(2π)))((∛(-1729/2+√(1729^2/4+4096^3/27)) -∛(-1729/2-√(1729^2/4+4096^3/27)))^1/9))

we obtain:

$$\sqrt{\left(\left(2^{-1/\pi}\sqrt[2\pi]{5\left(3+\sqrt{5}\right)\pi}\right)\sqrt[9]{\sqrt[3]{-\frac{1729}{2}+\sqrt{\frac{1729^2}{4}+\frac{4096^3}{27}}}-\sqrt[3]{-\frac{1729}{2}-\sqrt{\frac{1729^2}{4}+\frac{4096^3}{27}}}}\right)}$$

$$2^{-1/(2\pi)}\sqrt[18]{\sqrt[3]{\frac{\sqrt{\frac{274\,958\,621\,851}{3}}}{6}-\frac{1729}{2}}+\sqrt[3]{\frac{1729}{2}+\frac{\sqrt{\frac{274\,958\,621\,851}{3}}}{6}}}\Big/\sqrt[4\pi]{5\left(3+\sqrt{5}\right)\pi}$$

i.e.

2^(-1/(2π)) ((-1729/2+√(274958621851/3)/6)^(1/3)+(1729/2 + √(274958621851/3)/6)^(1/3))^(1/18) (5(3+√(5)) π)^(1/(4π))

= 1.61549140391….

The general "unitary" formula, which derives from DN Constant, is the following:

$$\sqrt{2\times\frac{2\cdot\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi\cdot 0.9991104684}(C\times \mathrm{R}\times 2.33\cdot 10^{-13})}}\cong 1.61803398\ldots=\frac{\sqrt{5}+1}{2}$$

Where $\frac{2\sqrt{2}}{\pi}$ is the Del Gaudio-Nardelli Constant, 0.9991104684 is the value of the following Rogers-Ramanujan continued fraction:

$$\frac{e^{-\frac{\pi}{\sqrt{5}}}}{\frac{\sqrt{5}}{1+\sqrt[5]{\sqrt{\varphi^5\sqrt[4]{5^3}}-1}}-\varphi+1}=1-\cfrac{e^{-\pi\sqrt{5}}}{1+\cfrac{e^{-2\pi\sqrt{5}}}{1+\cfrac{e^{-3\pi\sqrt{5}}}{1+\cfrac{e^{-4\pi\sqrt{5}}}{1+\ldots}}}}\approx 0.9991104684$$

C is any constant or solution to an equation, R is the radius of the Universe and $2.33*10^{-13}$ is the temperature of the universe expressed in GeV.

For example, $C = 9.9128$, inserting a radius of the Universe, which we have calculated, equal to $R = 2.06274*10^{12}$ years, from DN Constant "unitary" formula, we obtain:

√(2×(2·(((2√2)/π))^(1/16))/(1/(π·0.9991104684) (9.9128×(2.06274 × 10^12)×2.33·10^(-13))))

$$\sqrt{2\times\frac{2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi\times 0.9991104684}\times\frac{9.9128\times 2.06274\times 10^{12}\times 2.33}{10^{13}}}}$$

1.6180359123482642354401744088347098542545273508401733563064818107 …

1.618035912348…. result that is a very good approximation to the value of the golden ratio 1.618033988749… (Ramanujan Recurring Number)

We obtain also:

(√(2×(2·(((2√2)/π))^(1/16))/(1/(π·0.9991104684) (9.9128×(2.06274 × 10^12)×2.33·10^(-13)))))dxdydz

Indefinite integral assuming all variables are real

$$0.809018\, x^2\, y\, z + \text{constant}$$

Definite integral over a cube of edge length 2 L

$$\int_{-L}^{L}\int_{-L}^{L}\int_{-L}^{L} 1.61804\, dx\, dy\, dz = 12.9443\, L^3$$

Definite integral over a sphere of radius R

$$\iiint_{x^2+y^2+z^2<R^2} 1.61804\, dz\, dy\, dx = 6.77761\, R^3$$

From which, for L = R = 1 , dividing the two definite integral results by the original expression, we obtain:

12.9443/(√(2×(2·(((2√2)/π))^(1/16))/(1/(π·0.9991104684) (9.9128×(2.06274 × 10^12)×2.33·10^(-13)))))

Input interpretation

$$\frac{12.9443}{\sqrt{2\times\frac{2\sqrt[16]{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi\times 0.9991104684}\times\frac{9.9128\times 2.06274\times 10^{12}\times 2.33}{10^{13}}}}}$$

Result

8.00001…

8.00001…. ≈ 8

value that is linked to the "Ramanujan function" (an elliptic modular function that satisfies the need for "conformal symmetry") that has 8 "modes" corresponding to the physical vibrations of a superstring.

And

3*(6.77761/(√(2×(2·(((2√2)/π))^(1/16))/(1/(π·0.9991104684) (9.9128×(2.06274 × 10^12)×2.33·10^(-13))))))

Input interpretation

$$3\times\frac{6.77761}{\sqrt{2\times\frac{2^{16}\sqrt{\frac{2\sqrt{2}}{\pi}}}{\frac{1}{\pi\times 0.9991104684}\times\frac{9.9128\times 2.06274\times 10^{12}\times 2.33}{10^{13}}}}}$$

Result

12.5664…

12.5664…. ≈ 4π = Bekenstein-Hawking (S_{BH}) black hole entropy

Now, we have that:



Octahedron



Sphere

Given the value of a volume, independently of the solid, following the Poincaré Conjecture, we compare any solid "without holes" and a sphere. If we compare an octahedron with a sphere, we have:

$$\frac{4}{3}\pi r^3 = \frac{1}{3}\sqrt{2}a^3$$

If we consider the radius of the sphere as an unknown, we must find the value of the side of the octahedron which allows us to equalize the two volumes and which will give us the DN Constant as a result (which will therefore be equal to the radius of the sphere).

From

$$\frac{4}{3}\pi r^3 = \frac{1}{3}\sqrt{2}a^3$$

To find r we perform the following calculation:

$$r^3 = \frac{\frac{1}{3}\sqrt{2}a^3}{\frac{4}{3}\pi} = \frac{1}{3}\sqrt{2}a^3 \cdot \frac{3}{4\pi} = \frac{\sqrt{2}a^3}{4\pi} = \frac{\sqrt{2}\cdot\sqrt{2}a^3}{\sqrt{2}\cdot 4\pi} = \frac{2a^3}{\sqrt{2}\cdot 4\pi} = \frac{a^3}{2\sqrt{2}\pi}$$

$$r^3 = \frac{a^3}{2\sqrt{2}\pi}; \qquad r = \sqrt[3]{\frac{a^3}{2\sqrt{2}\pi}} = \frac{a}{\sqrt[3]{2\sqrt{2}\pi}}$$

To find *a*, we have, for

$$r = \frac{2\sqrt{2}}{\pi}; \qquad \frac{a}{\sqrt[3]{2\sqrt{2}\pi}} = \frac{2\sqrt{2}}{\pi};$$

Thus, multiplying both the sides by $\sqrt[3]{2\sqrt{2}\pi}$, we obtain:

$$a = \frac{2\sqrt{2}}{\pi} \cdot \sqrt[3]{2\sqrt{2}\pi} = \sqrt[3]{\frac{(2\sqrt{2})^3 2\sqrt{2}\pi}{\pi^3}} =$$

$$= \sqrt[3]{\frac{8 \cdot \sqrt{2^3} \cdot 2\sqrt{2}}{\pi^2}} = \sqrt[3]{\frac{8 \cdot \sqrt{2 \cdot 2^2} \cdot 2\sqrt{2}}{\pi^2}} = \sqrt[3]{\frac{16 \cdot \sqrt{2} \cdot 2\sqrt{2}}{\pi^2}}$$

$$= \sqrt[3]{\frac{32\sqrt{2}\sqrt{2}}{\pi^2}} = \sqrt[3]{\frac{32 \cdot 2}{\pi^2}} = \frac{\sqrt[3]{64}}{\sqrt[3]{\pi^2}} = \frac{4}{\sqrt[3]{\pi^2}}; \; a = \frac{4}{\sqrt[3]{\pi^2}}$$

Plot



Solution

$a = \frac{4}{\sqrt[3]{\pi^2}}$

for V = 1/3*√2*a^3 (octahedron volume) and V = (4/3*π*r^3) (sphere volume), we obtain:

from the octahedron volume, we have: V = 1/3*√2*a^3 = (1/3*√2*($\frac{4}{\sqrt[3]{\pi^2}}$)^3)

(1/3*√2*(4/∛(π^2))^3)

Input

$$\frac{1}{3}\sqrt{2}\left(\frac{4}{\sqrt[3]{\pi^2}}\right)^3$$

$\sqrt[3]{x}$ is the real-valued 3rd root of x

Exact result

$$\frac{64\sqrt{2}}{3\pi^2}$$

Decimal approximation

3.056848897337366735284768744174643472880661991020386025343029413 7

…

3.05684889733….

Property

$\frac{64\sqrt{2}}{3\pi^2}$ is a transcendental number

Series representations

$$\frac{1}{3}\sqrt{2}\left(\frac{4}{\sqrt[3]{\pi^2}}\right)^3 = \frac{64\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k z_0^{-k}}{k!}}{3\pi^2}$$

for (not ($z_0 \in \mathbb{R}$ and $-\infty < z_0 \leq 0$))

$$\frac{1}{3}\sqrt{2}\left(\frac{4}{\sqrt[3]{\pi^2}}\right)^3 = \frac{64\exp\left(i\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor\right)\sqrt{x}\sum_{k=0}^{\infty}\frac{(-1)^k(2-x)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}}{3\pi^2}$$

for ($x \in \mathbb{R}$ and $x < 0$)

$$\frac{1}{3}\sqrt{2}\left(\frac{4}{\sqrt[3]{\pi^2}}\right)^3 = \frac{64\left(\frac{1}{z_0}\right)^{1/2\lfloor\arg(2-z_0)/(2\pi)\rfloor} z_0^{1/2(1+\lfloor\arg(2-z_0)/(2\pi)\rfloor)}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k z_0^{-k}}{k!}}{3\pi^2}$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

And, from the sphere volume V = (4/3*π*r^3) = (4/3*π*((2√2)/π)^3)

(4/3*π*((2√2)/π)^3)

Input

$$\frac{4}{3}\pi\left(\frac{2\sqrt{2}}{\pi}\right)^3$$

Result

$$\frac{64\sqrt{2}}{3\pi^2}$$

Decimal approximation

3.056848897337366735284768744174643472880661991020386025343029413 7

…

3.05684889733….

Property

$\frac{64\sqrt{2}}{3\pi^2}$ is a transcendental number

Series representations

$$\frac{1}{3}\left(\pi\left(\frac{2\sqrt{2}}{\pi}\right)^3\right)4 = \frac{32\sqrt{z_0}^3\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k z_0^{-k}}{k!}\right)^3}{3\pi^2}$$

for (not ($z_0 \in \mathbb{R}$ and $-\infty < z_0 \leq 0$))

$$\frac{1}{3}\left(\pi\left(\frac{2\sqrt{2}}{\pi}\right)^3\right)4 = \frac{32\exp^3\left(i\,\pi\left\lfloor\frac{\arg(2-x)}{2\pi}\right\rfloor\right)\sqrt{x}^{\,3}\left(\sum_{k=0}^{\infty}\frac{(-1)^k(2-x)^k x^{-k}\left(-\frac{1}{2}\right)_k}{k!}\right)^3}{3\pi^2}$$

for $(x \in \mathbb{R}$ and $x < 0)$

$$\frac{1}{3}\left(\pi\left(\frac{2\sqrt{2}}{\pi}\right)^3\right)4 = \frac{32\left(\frac{1}{z_0}\right)^{3/2\lfloor\arg(2-z_0)/(2\pi)\rfloor} z_0^{3/2(1+\lfloor\arg(2-z_0)/(2\pi)\rfloor)}\left(\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(2-z_0)^k z_0^{-k}}{k!}\right)^3}{3\pi^2}$$

$n!$ is the factorial function

$(a)_n$ is the Pochhammer symbol (rising factorial)

$\mathbb{R}$ is the set of real numbers

$\arg(z)$ is the complex argument

$\lfloor x \rfloor$ is the floor function

i is the imaginary unit

From:

Modular equations and approximations to π - *Srinivasa Ramanujan* - Quarterly Journal of Mathematics, XLV, 1914, 350 – 372

We have that:

Hence

$$\begin{aligned} 64g_{22}^{24} &= e^{\pi\sqrt{22}} - 24 + 276e^{-\pi\sqrt{22}} - \cdots, \\ 64g_{22}^{-24} &= 4096e^{-\pi\sqrt{22}} + \cdots, \end{aligned}$$

so that

$$64(g_{22}^{24} + g_{22}^{-24}) = e^{\pi\sqrt{22}} - 24 + 4372e^{-\pi\sqrt{22}} + \cdots = 64\{(1+\sqrt{2})^{12} + (1-\sqrt{2})^{12}\}.$$

Hence

$$e^{\pi\sqrt{22}} = 2508951.9982\ldots.$$

Again

$$G_{37} = (6+\sqrt{37})^{\frac{1}{4}},$$

$$\begin{aligned} 64G_{37}^{24} &= e^{\pi\sqrt{37}} + 24 + 276e^{-\pi\sqrt{37}} + \cdots, \\ 64G_{37}^{-24} &= 4096e^{-\pi\sqrt{37}} - \cdots, \end{aligned}$$

so that

$$64(G_{37}^{24} + G_{37}^{-24}) = e^{\pi\sqrt{37}} + 24 + 4372e^{-\pi\sqrt{37}} - \cdots = 64\{(6+\sqrt{37})^{6} + (6-\sqrt{37})^{6}\}.$$

Hence

$$e^{\pi\sqrt{37}} = 199148647.999978\ldots.$$

Similarly, from

$$g_{58} = \sqrt{\left(\frac{5+\sqrt{29}}{2}\right)},$$

we obtain

$$64(g_{58}^{24} + g_{58}^{-24}) = e^{\pi\sqrt{58}} - 24 + 4372e^{-\pi\sqrt{58}} + \cdots = 64\left\{\left(\frac{5+\sqrt{29}}{2}\right)^{12} + \left(\frac{5-\sqrt{29}}{2}\right)^{12}\right\}.$$

Hence

$$e^{\pi\sqrt{58}} = 24591257751.99999982\ldots.$$

We note that, with regard 4372, we can to obtain the following results:

27((4372)^1/2-2-1/2(((√(10-2√5) -2))/((√5-1))))+φ

Input

$$27\left(\sqrt{4372}-2-\frac{1}{2}\times\frac{\sqrt{10-2\sqrt{5}}-2}{\sqrt{5}-1}\right)+\phi$$

ϕ is the golden ratio

Result

$$\phi+27\left(-2+2\sqrt{1093}-\frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)$$

Decimal approximation

1729.052694417090562517063720863714876368418930653845785481544702 3

…

1729.0526944….

This result is very near to the mass of candidate glueball **f_0(1710) scalar meson.** Furthermore, 1728 occurs in the algebraic formula for the j-invariant of an elliptic curve. ($1728 = 8^2 * 3^3$) The number 1728 is one less than the Hardy–Ramanujan number 1729 (taxicab number)

Alternate forms

$$\frac{1}{8}\left(-27\sqrt{5\left(10-2\sqrt{5}\right)}+58\sqrt{5}+432\sqrt{1093}-27\sqrt{2\left(5-\sqrt{5}\right)}-374\right)$$

$$\phi-54+54\sqrt{1093}+\frac{27}{4}\left(1+\sqrt{5}-\sqrt{2\left(5+\sqrt{5}\right)}\right)$$

$$\phi - 54 + 54\sqrt{1093} - \frac{27\left(\sqrt{10-2\sqrt{5}}-2\right)}{2\left(\sqrt{5}-1\right)}$$

Minimal polynomial

$$256\,x^8 + 95\,744\,x^7 - 3\,248\,750\,080\,x^6 - 914\,210\,725\,504\,x^5 + 15\,498\,355\,554\,921\,184\,x^4 + 2\,911\,478\,392\,539\,914\,656\,x^3 - 32\,941\,144\,911\,224\,677\,091\,680\,x^2 - 3\,092\,528\,914\,069\,760\,354\,714\,456\,x + 26\,320\,050\,609\,744\,039\,027\,169\,013\,041$$

Expanded forms

$$-\frac{187}{4} + \frac{29\sqrt{5}}{4} + 54\sqrt{1093} - \frac{27}{8}\sqrt{10-2\sqrt{5}} - \frac{27}{8}\sqrt{5\left(10-2\sqrt{5}\right)}$$

$$-\frac{107}{2} + \frac{\sqrt{5}}{2} + 54\sqrt{1093} + \frac{27}{\sqrt{5}-1} - \frac{27\sqrt{10-2\sqrt{5}}}{2\left(\sqrt{5}-1\right)}$$

Series representations

$$27\left(\sqrt{4372} - 2 - \frac{\sqrt{10-2\sqrt{5}}-2}{\left(\sqrt{5}-1\right)2}\right) + \phi = \left(162 - 108\sqrt{1093} - 2\phi - 108\sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k} + 108\sqrt{1093}\sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k} + 2\phi\sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k} - 27\sqrt{9-2\sqrt{5}}\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(9-2\sqrt{5}\right)^{-k}\right) \Big/ \left(2\left(-1+\sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k}\right)\right)$$

$$27\left(\sqrt{4372}\ -2-\frac{\sqrt{10-2\sqrt{5}}\ -2}{\left(\sqrt{5}\ -1\right)2}\right)+\phi=$$
$$\left(162-108\sqrt{1093}\ -2\phi-108\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}+\right.$$
$$108\sqrt{1093}\ \sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}+2\phi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}-$$
$$\left.27\sqrt{9-2\sqrt{5}}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(9-2\sqrt{5}\right)^{-k}}{k!}\right)\Big/$$
$$\left(2\left(-1+\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\right)$$

$$27\left(\sqrt{4372}\ -2-\frac{\sqrt{10-2\sqrt{5}}\ -2}{\left(\sqrt{5}\ -1\right)2}\right)+\phi=$$
$$\left(162-108\sqrt{1093}\ -2\phi-108\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}+\right.$$
$$108\sqrt{1093}\ \sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}+$$
$$2\phi\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}-$$
$$\left.27\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(10-2\sqrt{5}\ -z_0\right)^k\,z_0^{-k}}{k!}\right)\Big/$$
$$\left(2\left(-1+\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}\right)\right)$$
for $(\text{not } (z_0\in\mathbb{R} \text{ and } -\infty<z_0\le 0))$

Or:

27((4096+276)^1/2-2-1/2(((√(10-2√5) -2))/((√5-1))))+φ

Input

$$27\left(\sqrt{4096+276}-2-\frac{1}{2}\times\frac{\sqrt{10-2\sqrt{5}}-2}{\sqrt{5}-1}\right)+\phi$$

ϕ is the golden ratio

Result

$$\phi+27\left(-2+2\sqrt{1093}-\frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)$$

Decimal approximation

1729.052694417090562517063720863714876368418930653845785481544702 3

…

1729.0526944…. as above

Alternate forms

$$\frac{1}{8}\left(-27\sqrt{5\left(10-2\sqrt{5}\right)}+58\sqrt{5}+432\sqrt{1093}-27\sqrt{2\left(5-\sqrt{5}\right)}-374\right)$$

$$\phi-54+54\sqrt{1093}+\frac{27}{4}\left(1+\sqrt{5}-\sqrt{2\left(5+\sqrt{5}\right)}\right)$$

$$\phi-54+54\sqrt{1093}-\frac{27\left(\sqrt{10-2\sqrt{5}}-2\right)}{2\left(\sqrt{5}-1\right)}$$

Minimal polynomial

$$256\,x^8 + 95\,744\,x^7 - 3\,248\,750\,080\,x^6 - 914\,210\,725\,504\,x^5 + 15\,498\,355\,554\,921\,184\,x^4 + 2\,911\,478\,392\,539\,914\,656\,x^3 - 32\,941\,144\,911\,224\,677\,091\,680\,x^2 - 3\,092\,528\,914\,069\,760\,354\,714\,456\,x + 26\,320\,050\,609\,744\,039\,027\,169\,013\,041$$

Expanded forms

$$-\frac{187}{4} + \frac{29\sqrt{5}}{4} + 54\sqrt{1093} - \frac{27}{8}\sqrt{10 - 2\sqrt{5}} - \frac{27}{8}\sqrt{5\left(10 - 2\sqrt{5}\right)}$$

$$-\frac{107}{2} + \frac{\sqrt{5}}{2} + 54\sqrt{1093} + \frac{27}{\sqrt{5} - 1} - \frac{27\sqrt{10 - 2\sqrt{5}}}{2\left(\sqrt{5} - 1\right)}$$

Series representations

$$27\left(\sqrt{4096 + 276} - 2 - \frac{\sqrt{10 - 2\sqrt{5}} - 2}{\left(\sqrt{5} - 1\right)2}\right) + \phi = \left(162 - 108\sqrt{1093} - 2\phi - 108\sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k} + 108\sqrt{1093}\sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k} + 2\phi\sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k} - 27\sqrt{9 - 2\sqrt{5}}\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(9 - 2\sqrt{5}\right)^{-k}\right) \Big/ \left(2\left(-1 + \sqrt{4}\sum_{k=0}^{\infty} 4^{-k}\binom{\frac{1}{2}}{k}\right)\right)$$

$$27\left(\sqrt{4096+276}\ -2-\frac{\sqrt{10-2\sqrt{5}}\ -2}{\left(\sqrt{5}\ -1\right)2}\right)+\phi=$$
$$\left(162-108\sqrt{1093}\ -2\phi-108\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}+\right.$$
$$108\sqrt{1093}\ \sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}+2\phi\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}-$$
$$\left.27\sqrt{9-2\sqrt{5}}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(9-2\sqrt{5}\right)^{-k}}{k!}\right)\Big/$$
$$\left(2\left(-1+\sqrt{4}\ \sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\right)$$

$$27\left(\sqrt{4096+276}\ -2-\frac{\sqrt{10-2\sqrt{5}}\ -2}{\left(\sqrt{5}\ -1\right)2}\right)+\phi=$$
$$\left(162-108\sqrt{1093}\ -2\phi-108\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}+\right.$$
$$108\sqrt{1093}\ \sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}+$$
$$2\phi\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}-$$
$$\left.27\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(10-2\sqrt{5}\ -z_0\right)^k\,z_0^{-k}}{k!}\right)\Big/$$
$$\left(2\left(-1+\sqrt{z_0}\ \sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k\,z_0^{-k}}{k!}\right)\right)$$
for (not ($z_0 \in \mathbb{R}$ and $-\infty < z_0 \le 0$))

From which:

(27((4372)^1/2-2-1/2(((√(10-2√5) -2))/((√5-1))))+φ)^1/15

Input

$$\sqrt[15]{27\left(\sqrt{4372}-2-\frac{1}{2}\times\frac{\sqrt{10-2\sqrt{5}}-2}{\sqrt{5}-1}\right)+\phi}$$

ϕ is the golden ratio

Exact result

$$\sqrt[15]{\phi+27\left(-2+2\sqrt{1093}-\frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)}$$

Decimal approximation

1.643818568584986279990230131703681005418575687350518480483418312 4

…

$1.64381856858\ldots \approx \zeta(2) = \frac{\pi^2}{6} = 1.644934\ldots$

Alternate forms

$$\sqrt[15]{\phi-54+54\sqrt{1093}-\frac{27\left(\sqrt{10-2\sqrt{5}}-2\right)}{2\left(\sqrt{5}-1\right)}}$$

$$\frac{1}{\sqrt[15]{\frac{2\left(\sqrt{5}-1\right)}{166-108\sqrt{5}-108\sqrt{1093}+108\sqrt{5465}-27\sqrt{2\left(5-\sqrt{5}\right)}}}}$$

$$\sqrt[15]{\text{root of } 256\,x^8 + 95\,744\,x^7 - 3\,248\,750\,080\,x^6 - 914\,210\,725\,504\,x^5 + 15\,498\,355\,554\,921\,184\,x^4 + 2\,911\,478\,392\,539\,914\,656\,x^3 - 32\,941\,144\,911\,224\,677\,091\,680\,x^2 - 3\,092\,528\,914\,069\,760\,354\,714\,456\,x + 26\,320\,050\,609\,744\,039\,027\,169\,013\,041 \text{ near } x = 1729.05}$$

Minimal polynomial

$$256\,x^{120} + 95\,744\,x^{105} - 3\,248\,750\,080\,x^{90} - 914\,210\,725\,504\,x^{75} + 15\,498\,355\,554\,921\,184\,x^{60} + 2\,911\,478\,392\,539\,914\,656\,x^{45} - 32\,941\,144\,911\,224\,677\,091\,680\,x^{30} - 3\,092\,528\,914\,069\,760\,354\,714\,456\,x^{15} + 26\,320\,050\,609\,744\,039\,027\,169\,013\,041$$

Expanded forms

$$\sqrt[15]{\frac{1}{2}\left(1+\sqrt{5}\right) + 27\left(-2 + 2\sqrt{1093} - \frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)}$$

$$\sqrt[15]{-\frac{187}{4} + \frac{29\sqrt{5}}{4} + 54\sqrt{1093} - \frac{27}{8}\sqrt{10-2\sqrt{5}} - \frac{27}{8}\sqrt{5\left(10-2\sqrt{5}\right)}}$$

All 15th roots of ϕ + 27 (-2 + 2 sqrt(1093) - (sqrt(10 - 2 sqrt(5)) - 2)/(2 (sqrt(5) - 1)))

$$e^0\sqrt[15]{\phi + 27\left(-2 + 2\sqrt{1093} - \frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)} \approx 1.64382 \text{ (real, principal root)}$$

$$e^{(2i\pi)/15}\sqrt[15]{\phi + 27\left(-2 + 2\sqrt{1093} - \frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)} \approx 1.50170 + 0.6686\,i$$

$$e^{(4\,i\,\pi)/15}\sqrt[15]{\phi+27\left(-2+2\sqrt{1093}-\frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)}\approx 1.0999+1.2216\,i$$

$$e^{(2\,i\,\pi)/5}\sqrt[15]{\phi+27\left(-2+2\sqrt{1093}-\frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)}\approx 0.5080+1.5634\,i$$

$$e^{(8\,i\,\pi)/15}\sqrt[15]{\phi+27\left(-2+2\sqrt{1093}-\frac{\sqrt{10-2\sqrt{5}}-2}{2\left(\sqrt{5}-1\right)}\right)}\approx -0.17183+1.63481\,i$$

Series representations

$$\sqrt[15]{27\left(\sqrt{4372}-2-\frac{\sqrt{10-2\sqrt{5}}-2}{\left(\sqrt{5}-1\right)2}\right)+\phi}=$$
$$\frac{1}{\sqrt[15]{2}}\left(\left(\left(162-108\sqrt{1093}-2\phi-108\sqrt{4}\sum_{k=0}^{\infty}4^{-k}\binom{\frac{1}{2}}{k}+108\sqrt{1093}\sqrt{4}\right.\right.\right.$$
$$\left.\sum_{k=0}^{\infty}4^{-k}\binom{\frac{1}{2}}{k}+2\phi\sqrt{4}\sum_{k=0}^{\infty}4^{-k}\binom{\frac{1}{2}}{k}-27\sqrt{9-2\sqrt{5}}\right.$$
$$\left.\left.\left.\sum_{k=0}^{\infty}\binom{\frac{1}{2}}{k}\left(9-2\sqrt{5}\right)^{-k}\right)\Big/\left(-1+\sqrt{4}\sum_{k=0}^{\infty}4^{-k}\binom{\frac{1}{2}}{k}\right)\right)\right)\wedge(1/15)\Bigg)$$

$$\sqrt[15]{27\left(\sqrt{4372}-2-\frac{\sqrt{10-2\sqrt{5}}-2}{\left(\sqrt{5}-1\right)2}\right)+\phi}=$$
$$\frac{1}{\sqrt[15]{2}}\left(\left(\left(162-108\sqrt{1093}-2\phi-108\sqrt{4}\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}+\right.\right.\right.$$
$$108\sqrt{1093}\sqrt{4}\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}+2\phi\sqrt{4}\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}-$$
$$\left.27\sqrt{9-2\sqrt{5}}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(9-2\sqrt{5}\right)^{-k}}{k!}\right)\Big/$$
$$\left.\left.\left(-1+\sqrt{4}\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{4}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)\right)\wedge(1/15)\right)$$

$$\sqrt[15]{27\left(\sqrt{4372}-2-\frac{\sqrt{10-2\sqrt{5}}-2}{\left(\sqrt{5}-1\right)2}\right)+\phi}=$$
$$\frac{1}{\sqrt[15]{2}}\left(\left(\left(162-108\sqrt{1093}-2\phi-108\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}+\right.\right.\right.$$
$$108\sqrt{1093}\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}+$$
$$2\phi\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}-$$
$$\left.27\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k\left(10-2\sqrt{5}-z_0\right)^k z_0^{-k}}{k!}\right)\Big/$$
$$\left.\left.\left(-1+\sqrt{z_0}\sum_{k=0}^{\infty}\frac{(-1)^k\left(-\frac{1}{2}\right)_k(5-z_0)^k z_0^{-k}}{k!}\right)\right)\wedge(1/15)\right)$$

for $(\text{not } (z_0 \in \mathbb{R} \text{ and } -\infty < z_0 \le 0))$

Integral representation

$$(1+z)^a=\frac{\int_{-i\infty+\gamma}^{i\infty+\gamma}\frac{\Gamma(s)\,\Gamma(-a-s)}{z^s}\,ds}{(2\pi i)\,\Gamma(-a)} \quad \text{for } (0<\gamma<-\mathrm{Re}(a) \text{ and } |\arg(z)|<\pi)$$

From:

An Update on Brane Supersymmetry Breaking - *J. Mourad and A. Sagnotti* - arXiv:1711.11494v1 [hep-th] 30 Nov 2017

From the following vacuum equations:

$$T\, e^{\gamma_E \phi} = -\frac{\beta_E^{(p)}\, h^2}{\gamma_E}\, e^{-2(8-p)C + 2\beta_E^{(p)}\phi}$$

$$16\, k'\, e^{-2C} = \frac{h^2 \left(p + 1 - \frac{2\beta_E^{(p)}}{\gamma_E}\right) e^{-2(8-p)C + 2\beta_E^{(p)}\phi}}{(7-p)}$$

$$(A')^2 = k\, e^{-2A} + \frac{h^2}{16(p+1)} \left(7 - p + \frac{2\beta_E^{(p)}}{\gamma_E}\right) e^{-2(8-p)C + 2\beta_E^{(p)}\phi}$$

we have obtained, from the results almost equals of the equations, putting

$4096\, e^{-\pi\sqrt{18}}$ instead of

$$e^{-2(8-p)C + 2\beta_E^{(p)}\phi}$$

a new possible mathematical connection between the two exponentials. Thence, also the values concerning *p*, *C*, β_E and ϕ correspond to the exponents of *e* (i.e. of exp). Thence we obtain for p = 5 and β_E = 1/2:

$$e^{-6C+\phi} = 4096 e^{-\pi\sqrt{18}}$$

Therefore, with respect to the exponentials of the vacuum equations, the Ramanujan's exponential has a coefficient of 4096 which is equal to 64^2, while -6C+ϕ is equal to -$\pi\sqrt{18}$. From this it follows that it is possible to establish mathematically, the dilaton value.

For

exp((-Pi*sqrt(18)) we obtain:

Input:

$$\exp\left(-\pi\sqrt{18}\right)$$

Exact result:

$$e^{-3\sqrt{2}\,\pi}$$

Decimal approximation:

$$1.627201622607250929294215673911797954183858113695401 6\ldots \times 10^{-6}$$

1.6272016… * 10^{-6}

Property:

$e^{-3\sqrt{2}\,\pi}$ is a transcendental number

Series representations:

$$e^{-\pi\sqrt{18}} = e^{-\pi\sqrt{17}\,\sum_{k=0}^{\infty} 17^{-k}\binom{1/2}{k}}$$

$$e^{-\pi\sqrt{18}} = \exp\left(-\pi\sqrt{17}\,\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{17}\right)^k\left(-\frac{1}{2}\right)_k}{k!}\right)$$

$$e^{-\pi\sqrt{18}} = \exp\left(-\frac{\pi \sum_{j=0}^{\infty} \operatorname{Res}_{s=-\frac{1}{2}+j} 17^{-s}\,\Gamma\left(-\frac{1}{2}-s\right)\Gamma(s)}{2\sqrt{\pi}}\right)$$

Now, we have the following calculations:

$$e^{-6C+\phi} = 4096e^{-\pi\sqrt{18}}$$

$$e^{-\pi\sqrt{18}} = 1.6272016\ldots * 10^{-6}$$

from which:

$$\frac{1}{4096}e^{-6C+\phi} = 1.6272016\ldots * 10^{-6}$$

$$0.000244140625\ e^{-6C+\phi} = e^{-\pi\sqrt{18}} = 1.6272016\ldots * 10^{-6}$$

Now:

$$\ln\left(e^{-\pi\sqrt{18}}\right) = -13.328648814475 = -\pi\sqrt{18}$$

And:

(1.6272016* 10^-6) *1/ (0.000244140625)

Input interpretation:

$$\frac{1.6272016}{10^6} \times \frac{1}{0.000244140625}$$

Result:

0.0066650177536

0.006665017...

Thence:

$$0.000244140625\, e^{-6C+\phi} = e^{-\pi\sqrt{18}}$$

Dividing both sides by 0.000244140625, we obtain:

$$\frac{0.000244140625}{0.000244140625}\, e^{-6C+\phi} = \frac{1}{0.000244140625}\, e^{-\pi\sqrt{18}}$$

$$e^{-6C+\phi} = 0.0066650177536$$

((((exp((-Pi*sqrt(18)))))))*1/0.000244140625

Input interpretation:

$$\exp\left(-\pi\sqrt{18}\right) \times \frac{1}{0.000244140625}$$

Result:

0.00666501785...

0.00666501785…

Series representations:

$$\frac{\exp\left(-\pi\sqrt{18}\right)}{0.000244141} = 4096\exp\left(-\pi\sqrt{17}\sum_{k=0}^{\infty}17^{-k}\binom{\frac{1}{2}}{k}\right)$$

$$\frac{\exp\left(-\pi\sqrt{18}\right)}{0.000244141} = 4096\exp\left(-\pi\sqrt{17}\sum_{k=0}^{\infty}\frac{\left(-\frac{1}{17}\right)^{k}\left(-\frac{1}{2}\right)_{k}}{k!}\right)$$

$$\frac{\exp\left(-\pi\sqrt{18}\right)}{0.000244141} = 4096\exp\left(-\frac{\pi\sum_{j=0}^{\infty}\operatorname{Res}_{s=-\frac{1}{2}+j}\,17^{-s}\,\Gamma\left(-\frac{1}{2}-s\right)\Gamma(s)}{2\sqrt{\pi}}\right)$$

Now:

$$e^{-6C+\phi} = 0.0066650177536$$

$$\exp\left(-\pi\sqrt{18}\right)\times\frac{1}{0.000244140625} =$$

$$e^{-\pi\sqrt{18}}\times\frac{1}{0.000244140625}$$

$$= 0.00666501785\ldots$$

From:

ln(0.00666501784619)

Input interpretation:

$\log(0.00666501784619)$

Result:

–5.010882647757…

-5.010882647757…

Alternative representations:

$\log(0.006665017846190000) = \log_e(0.006665017846190000)$

$\log(0.006665017846190000) = \log(a)\log_a(0.006665017846190000)$

$\log(0.006665017846190000) = -\mathrm{Li}_1(0.993334982153810000)$

Series representations:

$$\log(0.006665017846190000) = -\sum_{k=1}^{\infty} \frac{(-1)^k\,(-0.993334982153810000)^k}{k}$$

$$\log(0.006665017846190000) = 2\,i\,\pi\left\lfloor \frac{\arg(0.006665017846190000 - x)}{2\,\pi} \right\rfloor + \log(x) - \sum_{k=1}^{\infty} \frac{(-1)^k\,(0.006665017846190000 - x)^k\,x^{-k}}{k} \quad \text{for } x < 0$$

$$\log(0.006665017846190000) = \left\lfloor \frac{\arg(0.006665017846190000 - z_0)}{2\pi} \right\rfloor \log\left(\frac{1}{z_0}\right) + \log(z_0) + \left\lfloor \frac{\arg(0.006665017846190000 - z_0)}{2\pi} \right\rfloor \log(z_0) - \sum_{k=1}^{\infty} \frac{(-1)^k \, (0.006665017846190000 - z_0)^k \, z_0^{-k}}{k}$$

Integral representation:

$$\log(0.006665017846190000) = \int_1^{0.006665017846190000} \frac{1}{t}\, dt$$

In conclusion:

$$-6C + \phi = -5.010882647757\ldots$$

and for C = 1, we obtain:

$$\phi = -5.010882647757 + 6 = 0.989117352243 = \boldsymbol{\phi}$$

Note that the values of n_s (spectral index) 0.965, of the average of the Omega mesons Regge slope 0.987428571 and of the dilaton 0.989117352243, are also connected to the following two Rogers-Ramanujan continued fractions:

$$\frac{e^{-\frac{\pi}{5}}}{\sqrt{(\varphi-1)\sqrt{5}} - \varphi + 1} = 1 - \cfrac{e^{-\pi}}{1 + \cfrac{e^{-2\pi}}{1 + \cfrac{e^{-3\pi}}{1 + \cfrac{e^{-4\pi}}{1 + \ldots}}}} \approx 0.9568666373$$

$$\frac{e^{-\frac{\pi}{\sqrt{5}}}}{\frac{\sqrt{5}}{1+\sqrt[5]{\sqrt{\varphi^5\sqrt[4]{5^3}}-1}}-\varphi+1}=1-\frac{e^{-\pi\sqrt{5}}}{1+\frac{e^{-2\pi\sqrt{5}}}{1+\frac{e^{-3\pi\sqrt{5}}}{1+\frac{e^{-4\pi\sqrt{5}}}{1+...}}}}\approx 0.9991104684$$

(http://www.bitman.name/math/article/102/109/)

Also performing the 512th root of the inverse value of the Pion meson rest mass 139.57, we obtain:

((1/(139.57)))^1/512

Input interpretation:

$$\sqrt[512]{\frac{1}{139.57}}$$

Result:

0.990400732708644027550973755713301415460732796178555551684...

0.99040073.... result very near to the dilaton value 0.989117352243 = $\boldsymbol{\phi}$ and to the value of the following Rogers-Ramanujan continued fraction:

$$\frac{e^{-\frac{\pi}{\sqrt{5}}}}{\frac{\sqrt{5}}{1+\sqrt[5]{\sqrt{\varphi^5\sqrt[4]{5^3}}-1}}-\varphi+1}=1-\frac{e^{-\pi\sqrt{5}}}{1+\frac{e^{-2\pi\sqrt{5}}}{1+\frac{e^{-3\pi\sqrt{5}}}{1+\frac{e^{-4\pi\sqrt{5}}}{1+...}}}}\approx 0.9991104684$$

From

Properties of Nilpotent Supergravity - *E. Dudas, S. Ferrara, A. Kehagias and A. Sagnotti* - arXiv:1507.07842v2 [hep-th] 14 Sep 2015

We have that:

Cosmological inflation with a tiny tensor–to–scalar ratio r, consistently with PLANCK data, may also be described within the present framework, for instance choosing

$$\alpha(\Phi) \;=\; i\,M\left(\Phi \;+\; b\,\Phi\,e^{ik\,\Phi}\right)\,. \tag{4.35}$$

This potential bears some similarities with the Kähler moduli inflation of [32] and with the poly–instanton inflation of [33]. One can verify that $\chi = 0$ solves the field equations, and that the potential along the $\chi = 0$ trajectory is now

$$V \;=\; \frac{M^2}{3}\left(1 \;-\; a\,\phi\,e^{-\gamma\,\phi}\right)^2\,. \tag{4.36}$$

We analyzing the following equation:

$$V \;=\; \frac{M^2}{3}\left(1 \;-\; a\,\phi\,e^{-\gamma\,\phi}\right)^2\,.$$

$$\phi \;=\; \varphi \;-\; \frac{\sqrt{6}}{k}\,,$$

$$a \;=\; \frac{b\gamma}{e} \;<\; 0\,, \qquad \gamma \;=\; \frac{k}{\sqrt{6}} \;<\; 0\,.$$

We have:

(M^2)/3*[1-(b/euler number * k/sqrt6) * (φ- sqrt6/k) * exp(-(k/sqrt6)(φ- sqrt6/k))]^2

i.e.

V = (M^2)/3*[1-(b/euler number * k/sqrt6) * (φ- sqrt6/k) * exp(-(k/sqrt6)(φ- sqrt6/k))]^2

For k = 2 and φ = 0.9991104684, that is the value of the scalar field that is equal to the value of the following Rogers-Ramanujan continued fraction:

$$\frac{e^{-\frac{\pi}{\sqrt{5}}}}{\frac{\sqrt{5}}{1+\sqrt[5]{\sqrt{\varphi^5\sqrt[4]{5^3}}-1}}-\varphi+1}=1-\frac{e^{-\pi\sqrt{5}}}{1+\frac{e^{-2\pi\sqrt{5}}}{1+\frac{e^{-3\pi\sqrt{5}}}{1+\frac{e^{-4\pi\sqrt{5}}}{1+...}}}}\approx 0.9991104684$$

we obtain:

V = (M^2)/3*[1-(b/euler number * 2/sqrt6) * (0.9991104684- sqrt6/2) * exp(-(2/sqrt6)(0.9991104684- sqrt6/2))]^2

Input interpretation:

$$V=\frac{M^2}{3}\left(1-\left(\frac{b}{e}\times\frac{2}{\sqrt{6}}\right)\left(0.9991104684-\frac{\sqrt{6}}{2}\right)\exp\left(-\frac{2}{\sqrt{6}}\left(0.9991104684-\frac{\sqrt{6}}{2}\right)\right)\right)^2$$

Result:

$$V=\frac{1}{3}(0.0814845\,b+1)^2\,M^2$$

Solutions:

$$b = \frac{225.913\left(-0.054323\,M^2 \pm 6.58545\times 10^{-10}\sqrt{M^4}\right)}{M^2} \quad (M \neq 0)$$

Alternate forms:

$$V = 0.00221324\,(b + 12.2723)^2\,M^2$$

$$V = 0.00221324\left(b^2\,M^2 + 24.5445\,b\,M^2 + 150.609\,M^2\right)$$

$$-0.00221324\,b^2\,M^2 - 0.054323\,b\,M^2 - \frac{M^2}{3} + V = 0$$

Expanded form:

$$V = 0.00221324\,b^2\,M^2 + 0.054323\,b\,M^2 + \frac{M^2}{3}$$

Alternate form assuming b, M, and V are positive:

$$V = 0.00221324\,(b + 12.2723)^2\,M^2$$

Alternate form assuming b, M, and V are real:

$$V = 0.00221324\,b^2\,M^2 + 0.054323\,b\,M^2 + 0.333333\,M^2 + 0$$

Derivative:

$$\frac{\partial}{\partial b}\left(\frac{1}{3}\,(0.0814845\,b + 1)^2\,M^2\right) = 0.054323\,(0.0814845\,b + 1)\,M^2$$

Implicit derivatives

$$\frac{\partial b(M,V)}{\partial V} = \frac{154\,317\,775\,011\,120\,075}{36\,961\,748\,(226\,802\,245 + 18\,480\,874\,b)\,M^2}$$

$$\frac{\partial b(M,V)}{\partial M} = -\frac{\frac{226\,802\,245}{18\,480\,874} + b}{M}$$

$$\frac{\partial M(b,V)}{\partial V} = \frac{154\,317\,775\,011\,120\,075}{2\,(226\,802\,245 + 18\,480\,874\,b)^2\,M}$$

$$\frac{\partial M(b,V)}{\partial b} = -\frac{18\,480\,874\,M}{226\,802\,245 + 18\,480\,874\,b}$$

$$\frac{\partial V(b,M)}{\partial M} = \frac{2\,(226\,802\,245 + 18\,480\,874\,b)^2\,M}{154\,317\,775\,011\,120\,075}$$

$$\frac{\partial V(b,M)}{\partial b} = \frac{36\,961\,748\,(226\,802\,245 + 18\,480\,874\,b)\,M^2}{154\,317\,775\,011\,120\,075}$$

Global minimum:

$$\min\left\{\frac{1}{3}\,(0.0814845\,b + 1)^2\,M^2\right\} = 0 \text{ at } (b, M) = (-16, 0)$$

Global minima:

$$\min\left\{\frac{1}{3}\,M^2\left(1 - \frac{(b\,2)\left(0.9991104684 - \frac{\sqrt{6}}{2}\right)\exp\left(-\frac{2\left(0.9991104684 - \frac{\sqrt{6}}{2}\right)}{\sqrt{6}}\right)}{e\,\sqrt{6}}\right)^2\right\} = 0$$

$$\text{for } b = -\frac{226\,802\,245}{18\,480\,874}$$

$$\min\left\{\frac{1}{3}M^2\left(1-\frac{(b\ 2)\left(0.9991104684-\frac{\sqrt{6}}{2}\right)\exp\left(-\frac{2\left(0.9991104684-\frac{\sqrt{6}}{2}\right)}{\sqrt{6}}\right)}{e\sqrt{6}}\right)^2\right\}=0$$

for $M=0$

From:

$$b=\frac{225.913\left(-0.054323\,M^2\pm 6.58545\times 10^{-10}\sqrt{M^4}\right)}{M^2}\quad (M\neq 0)$$

we obtain:

(225.913 (-0.054323 M^2 + 6.58545×10^-10 sqrt(M^4)))/M^2

Input interpretation:

$$\frac{225.913\left(-0.054323\,M^2+6.58545\times 10^{-10}\sqrt{M^4}\right)}{M^2}$$

Result:

$$\frac{225.913\left(6.58545\times 10^{-10}\sqrt{M^4}-0.054323\,M^2\right)}{M^2}$$

Plots:



(*M* from −1 to 0.2)



(*M* from −4.6 to 3.9)

Alternate form assuming M is real:

-12.2723

-12.2723 result very near to the black hole entropy value 12.1904 = ln(196884)

Alternate forms:

$$-\frac{12.2723\left(M^2-1.21228\times10^{-8}\sqrt{M^4}\right)}{M^2}$$

$$\frac{1.48774\times10^{-7}\sqrt{M^4}-12.2723\,M^2}{M^2}$$

Expanded form:

$$\frac{1.48774\times10^{-7}\sqrt{M^4}}{M^2}-12.2723$$

Property as a function:

Parity

even

Series expansion at M = 0:

$$\left(\frac{1.48774\times10^{-7}\sqrt{M^4}}{M^2}-12.2723\right)+O\left(M^6\right)$$

(generalized Puiseux series)

Series expansion at M = ∞:

$$-12.2723$$

Derivative:

$$\frac{d}{dM}\left(\frac{225.913\left(6.58545\times10^{-10}\sqrt{M^4}-0.054323\,M^2\right)}{M^2}\right)=\frac{3.55271\times10^{-15}}{M}$$

Indefinite integral:

$$\int\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}\,dM=$$
$$\frac{1.48774\times10^{-7}\sqrt{M^4}}{M}-12.2723\,M+\text{constant}$$

Global maximum:

$$\max\left\{\frac{225.913\left(6.58545\times10^{-10}\sqrt{M^4}-0.054323\,M^2\right)}{M^2}\right\}=$$
$$-\frac{140\,119\,826\,723\,990\,341\,497\,649}{11\,417\,594\,849\,251\,000\,000\,000}\ \text{at}\ M=-1$$

Global minimum:

$$\min\left\{\frac{225.913\left(6.58545\times10^{-10}\sqrt{M^4}-0.054323\,M^2\right)}{M^2}\right\}=$$
$$-\frac{140\,119\,826\,723\,990\,341\,497\,649}{11\,417\,594\,849\,251\,000\,000\,000}\ \text{at}\ M=-1$$

Limit:

$$\lim_{M\to\pm\infty}\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}=-12.2723$$

Definite integral after subtraction of diverging parts:

$$\int_0^{\infty}\left(\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}--12.2723\right)dM=0$$

From b that is equal to

$$\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}$$

From:

$$V = \frac{1}{3}(0.0814845\,b + 1)^2\,M^2$$

we obtain:

1/3 (0.0814845 ((225.913 (-0.054323 M^2 + 6.58545×10^-10 sqrt(M^4)))/M^2) + 1)^2 M^2

Input interpretation:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}+1\right)^2 M^2$$

Result:

0

Plots: **(possible mathematical connection with an open string)**



(M from −1 to 0.2)

M = -0.5; M = 0.2

(possible mathematical connection with an open string)



(M from −4.6 to 3.9)

M = 2 ; M = 3

Root:

$M = 0$

Property as a function:

Parity

even

Series expansion at M = 0:

$O\left(M^{62\,194}\right)$

(Taylor series)

Series expansion at M = ∞:

$1.75541 \times 10^{-15}\, M^2 + O\left(\left(\frac{1}{M}\right)^{62\,194}\right)$

(Taylor series)

Definite integral after subtraction of diverging parts:

$$\int_0^{\infty} \left(\frac{1}{3} M^2 \left(1 + \frac{18.4084 \left(-0.054323\, M^2 + 6.58545 \times 10^{-10} \sqrt{M^4} \right)}{M^2} \right)^2 - 1.75541 \times 10^{-15}\, M^2 \right) dM = 0$$

For M = - 0.5 , we obtain:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}+1\right)^2 M^2$$

1/3 (0.0814845 ((225.913 (-0.054323 (-0.5)^2 + 6.58545×10^-10 sqrt((-0.5)^4)))/(-0.5)^2) + 1)^2 * (-0.5^2)

Input interpretation:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\,(-0.5)^2+6.58545\times10^{-10}\sqrt{(-0.5)^4}\right)}{(-0.5)^2}+1\right)^2 (-0.5^2)$$

Result:

$$-4.388513449474645453489707833780880208333333333333333333\ldots\times10^{-16}$$

-4.38851344947*10^-16

For M = 0.2:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}+1\right)^2 M^2$$

1/3 (0.0814845 ((225.913 (-0.054323 0.2^2 + 6.58545×10^-10 sqrt(0.2^4)))/0.2^2) + 1)^2 0.2^2

Input interpretation:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\times 0.2^2+6.58545\times 10^{-10}\sqrt{0.2^4}\right)}{0.2^2}+1\right)^2\times 0.2^2$$

Result:

$7.021621519159432725583532534049408333333333333333333333333\ldots\times 10^{-17}$

7.021621519159*10^{-17}

For M = 3:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\,M^2+6.58545\times 10^{-10}\sqrt{M^4}\right)}{M^2}+1\right)^2 M^2$$

1/3 (0.0814845 ((225.913 (-0.054323 3^2 + 6.58545×10^-10 sqrt(3^4)))/3^2) + 1)^2 3^2

Input interpretation:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\times 3^2+6.58545\times 10^{-10}\sqrt{3^4}\right)}{3^2}+1\right)^2\times 3^2$$

Result:

$1.579864841810872363256294820161116875\times 10^{-14}$

1.57986484181*10^{-14}

For M = 2:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\,M^2+6.58545\times10^{-10}\sqrt{M^4}\right)}{M^2}+1\right)^2 M^2$$

1/3 (0.0814845 ((225.913 (-0.054323 2^2 + 6.58545×10^-10 sqrt(2^4)))/2^2) + 1)^2 2^2

Input interpretation:

$$\frac{1}{3}\left(0.0814845\times\frac{225.913\left(-0.054323\times2^2+6.58545\times10^{-10}\sqrt{2^4}\right)}{2^2}+1\right)^2\times2^2$$

Result:

$7.021621519159432725583532534049408333333333333333333333333\ldots\times10^{-15}$

$7.021621519*10^{-15}$

From the four results

7.021621519*10^-15 ; 1.57986484181*10^-14 ; 7.021621519159*10^-17 ;

-4.38851344947*10^-16

we obtain, after some calculations:

sqrt[1/(2Pi)(7.021621519*10^-15 + 1.57986484181*10^-14 +7.021621519*10^-17 - 4.38851344947*10^-16)]

Input interpretation:

$$\sqrt{\left(\frac{1}{2\pi}\left(7.021621519\times10^{-15}+1.57986484181\times10^{-14}+7.021621519\times10^{-17}-4.38851344947\times10^{-16}\right)\right)}$$

Result:

$5.9776991059\ldots\times10^{-8}$

$5.9776991059*10^{-8}$ result very near to the Planck's electric flow 5.975498×10^{-8} that is equal to the following formula:

$$\phi_P^E=\mathbf{E}_P l_P^2=\phi_P l_P=\sqrt{\frac{\hbar c}{\varepsilon_0}}$$

We note that:

1/55*(([(((1/[(7.021621519*10^-15 + 1.57986484181*10^-14 +7.021621519*10^-17 -4.38851344947*10^-16)]))^1/7]-((log^(5/8)(2))/(2 2^(1/8) 3^(1/4) e log^(3/2)(3)))))

Input interpretation:

$$\frac{1}{55}\left(\left(1/\left(7.021621519\times10^{-15}+1.57986484181\times10^{-14}+7.021621519\times10^{-17}-4.38851344947\times10^{-16}\right)\right)\text{ ^ }(1/7)-\frac{\log^{5/8}(2)}{2\sqrt[8]{2}\,\sqrt[4]{3}\,e\log^{3/2}(3)}\right)$$

$\log(x)$ is the natural logarithm

Result:

1.6181818182…

1.6181818182… result that is a very good approximation to the value of the golden ratio 1.618033988749...

From the Planck units:

Planck Length

$$l_{\mathrm{P}} = \sqrt{\frac{4\pi\hbar G}{c^3}}$$

5.729475 * 10^{-35} Lorentz-Heaviside value

Planck's Electric field strength

$$\mathbf{E}_{\mathrm{P}} = \frac{F_{\mathrm{P}}}{q_{\mathrm{P}}} = \sqrt{\frac{c^7}{16\pi^2\varepsilon_0\hbar G^2}}$$

1.820306 * 10^{61} V*m Lorentz-Heaviside value

Planck's Electric flux

$$\phi_{\mathrm{P}}^{E} = \mathbf{E}_{\mathrm{P}} l_{\mathrm{P}}^2 = \phi_{\mathrm{P}} l_{\mathrm{P}} = \sqrt{\frac{\hbar c}{\varepsilon_0}}$$

5.975498*10^{-8} V*m Lorentz-Heaviside value

Planck's Electric potential

$$\phi_P = V_P = \frac{E_P}{q_P} = \sqrt{\frac{c^4}{4\pi\varepsilon_0 G}}$$

1.042940*10^{27} V Lorentz-Heaviside value

Relationship between Planck's Electric Flux and Planck's Electric Potential

$E_P * l_P = (1.820306 * 10^{61}) * 5.729475 * 10^{-35}$

Input interpretation:

$$\frac{(1.820306 \times 10^{61}) \times 5.729475}{10^{35}}$$

Result:

1 042 939 771 935 000 000 000 000 000

Scientific notation:

$1.042939771935 \times 10^{27}$

$1.042939771935*10^{27} \approx 1.042940*10^{27}$

Or:

$E_P * l_P^2 / l_P = (5.975498*10^{-8})*1/(5.729475 * 10^{-35})$

Input interpretation:

$$5.975498 \times 10^{-8} \times \frac{1}{\frac{5.729475}{10^{35}}}$$

Result:

$1.042939885417075735560413475929295441554418162222542205001 33\ldots \times 10^{27}$

$1.042939885417*10^{27} \approx 1.042940*10^{27}$

Fig. 1



It is therefore possible to consider the vortices of the "quantum vacuum" schematized as cubes or octahedrons (the + sign inside a given vortex indicates its counterclockwise rotation, while the - sign indicates its clockwise rotation). Between vortex and vortex there is a layer of "bubbles"-universes (or universes-spheres), which flows, as in the simplified two-dimensional drawing, from A to B

Fig. 2

Immagine dello Spazio-Tempo a scala quantistica: le cironferenze in rosso rappresentano i punti corrispondenti alle dimensioni compattificate e gli esagoni in blu, rappresentano le “fluttuazioni” (universi potenziali – cerchi verdi) del vuoto quantistico (2D)



Proposal

Image of space-time at quantum scale: the circles in red represent the points corresponding to the compactified dimensions and the hexagons in blue, represent the "fluctuations" (potential universes - green circles) of the quantum vacuum (2D). In reality, we will have n-dimensional hyperspheres in which the compactified dimensions "roll up" and octahedrons representing the "fluctuations", containing spheres (bubbles of potential universes), of the quantum vacuum

Acknowledgments

We would like to thank Professor **Augusto Sagnotti** theoretical physicist at Scuola Normale Superiore (Pisa – Italy) for his very useful explanations and his availability.

References

A Number Theoretic Analysis of the Enthalpy, Enthalpy Energy Density, Thermodynamic Volume, and the Equation of State of a Modified White Hole, and the Implications to the Quantum Vacuum Spacetime, Matter Creation and the Planck Frequency. - *Nardelli, M., Kubeka, A.S. and Amani, A.* (2024) - Journal of Modern Physics , 15, 1-50. - https://doi.org/10.4236/jmp.2024.151001

Modular equations and approximations to π - *Srinivasa Ramanujan* - Quarterly Journal of Mathematics, XLV, 1914, 350 – 372

An Update on Brane Supersymmetry Breaking - *Jihad Mourad and Augusto Sagnotti* - arXiv:1711.11494v1 [hep-th] 30 Nov 2017

Properties of Nilpotent Supergravity - *E. Dudas, S. Ferrara, A. Kehagias and A. Sagnotti* - arXiv:1507.07842v2 [hep-th] 14 Sep 2015

See also:

The Geometry of the MRB constant *by Marvin Ray Burns*

https://www.academia.edu/22271085/The_Geometry_of_the_MRB_constant

(See also Page 29 the applications of the CMRB in various sectors of Theoretical Physics (String Theory) and Cosmology)

http://xoom.virgilio.it/source_filemanager/na/ar/nardelli/michele%20and%20antonio%20papers/Try%20to%20beat%20these%20MRB%20constant%20records!%20-%20Online%20Technical%20Discussion%20Groups%E2%80%94Wolfram%20Community%20b.pdf