





Nucleic Acids Research

Article Navigation

Termination of DNA synthesis by novel 3'modifieddeoxyribonucleoside 5'-triphosphates

Michael L. Metzker ➡, Ramesh Raghavachari, Stephen Richards, Swanee E. Jacutin, Andrew Civitello, Kevin Burgess, Richard A. Gibbs

Nucleic Acids Research, Volume 22, Issue 20, 11 October 1994, Pages 4259–4267, https://doi.org/10.1093/nar/22.20.4259 Published: 11 October 1994 Article history ▼

🎸 Cite 🛛 🔑 Permissions 🛛 < Share 🔻

Abstract

Eight 3'-modified-dNTPs were synthesized and tested in two different DNA template assays for incorporation activity. From this enzymatic screen, two 3'-O-methyldNTPs were shown to terminate DNA syntheses mediated by a number of polymerases and may be used as alternative terminators in Sanger sequencing. 3-O-(2-Nitrobenzyl) -dATP is a UV sensitive nucleotide and was shown to be incorporated by several thermostable DNA polymerases. Base specific termination and efficient photolytic removal of the 3'-protecting group was demonstrated. Following deprotection, DNA synthesis was reinitiated by the incorporation of natural nucleotides into DNA. The identification of this labile terminator and the demonstration of a one cycle stopstart DNA synthesis are initial steps in the development of a novel sequencing strategy.

Issue Section: CHEMISTRY

Illumina Ex. 1097 Illumina v. Columbia IPR2018-00291, -00318, -00322, -00385, and -00797