McGraw-Hill Dictionary of CHEMISTRY

Sybil P. Parker

Editor in Chief

McGraw-Hill Book Company

New York St. Louis San Francisco

Auckland Bogotá Guatemala Hamburg Johannesburg Lisbon London Madrid New Delhi Montreal Panama Paris São Paulo Singapore Sydney Tokyo

Mexico San Juan Toronto

> Illumina Ex. 1103 Illumina v. Columbia IPR2018-00291, -00318, -00322, -00385, and -00797

McGRAW-HILL DICTIONARY OF CHEMISTRY The material in this Dictionary has been published previously in the McGRAW-HILL DICTIONARY OF SCIENTIFIC AND TECHNICAL TERMS, Third Edition, copyright © 1984 by McGraw-Hill, Inc. All rights reserved. Philippines copyright 1984 by McGraw-Hill, Inc. Printed in the United States of America. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a data base or retrieval system, without the prior written permission of the publisher.

1234567890 DODO 8910987654

ISBN 0-07-045420-5

Library of Congress Cataloging in Publication Data

McGraw-Hill dictionary of chemistry.

1. Chemistry—Dictionaries. I. Parker, Sybil P. II. McGraw-Hill Book Company. III. Title: Dictionary of chemistry. QD5.M357 1984 540'.3'21 84-12205 ISBN 0-07-045420-5

24 allycarbamide

allycarbamide See allylurea.

- allyl- A prefix used in names of compounds whose structure contains an allyl cation.
- allylacetone CH₂CHCH₂CH₂COCH₃ A colorless liquid, soluble in water and organic solvents; used in pharmaceutical synthesis, perfumes, fungicides, and insecticides.
- allyi alcohoi CH₂CHCH₂OH Colorless, pungent liquid, boiling at 96°C; soluble in water; made from allyl chloride by hydrolysis. Also known as 2-propen-1-ol.
- allylamine CH₂CHCH₂NH₂ A yellow oil that is miscible with water; boils at 58°C; prepared from mustard oil. Also known as 2-propen-1-amine.
- allyi bromide C₃H₅Br A colorless to light yellow, irritating toxic liquid with a boiling point of 71.3°C; soluble in organic solvents; used in organic synthesis and for the manufacture of synthetic perfumes. Also known as bromoally-lene; 3-bromo-1-propene.
- ally cation A carbonium cation with a structure usually represented as $CH_2 = CH CH_2^+$; attachment site is the saturated carbon atom.
- allyi chloride CH₂CHCH₂Cl A volatile, pungent, toxic, flammable, colorless liquid, boiling at 46°C; insoluble in water; made by chlorination of propylene at high temperatures. Also known as 3-chloro-1-propene.
- allyl cyanide C_4H_5N A liquid with an onionlike odor and a boiling point of 119°C; slightly soluble in water; used as a cross-linking agent in polymerization. Also known as 3-butenenitrile; vinylacetonitrile.
- allylene $CH_3C:CH$ An acetylenic, three-carbon hydrocarbon; a colorless gas boiling at $-24^{\circ}C$; soluble in ether. Also known as methyl acetylene; propine; propyne.
- allylic rearrangement In a three-carbon molecule, the shifting of a double bond from the 1,2 carbon position to the 2,3 position, with the accompanying migration of an entering substituent or substituent group from the third carbon to the first.

allyl isosulfocyanate See allyl isothiocyanate.

- allyl isothiocyanate CH₂CH:CH₂NCS A pungent, colorless to pale-yellow liquid; soluble in alcohol, slightly soluble in water; irritating odor; boiling point 152°C; used as a fumigant and as a poison gas. Also known as allyl isosulfocyanate; mustard oil.
- allyl mercaptan CH₂CHCH₂SH A colorless liquid with a boiling point of 67–68°C; soluble in ether and alcohol; used as intermediate in pharmaceutical manufacture. Also known as allyl thiol; 2-propene-1-thiol.

allyl plastic See allyl resin.

- allyl resin Any of a class of thermosetting synthetic resins derived from esters of allyl alcohol or allyl chloride; used in making cast and laminated products. Also known as allyl plastic.
- allyl sulfide $(CH_2CHCH_2)_2S$ A colorless liquid with a garliclike odor and a boiling point of 139°C; used in synthetic oil of garlic. Also known as diallyl sulfide; oil garlic; thioallyl ether.

allylsulf allyl sulf allyl thio

allyithio solu

bam allyltric

poin

allylure wati

inhi

allyxyc for

lyla

cell

by

tei

po

alpha c and

alpha

alpha

alpha

alpha

alpha

ne rac

> alpha tic alpha

> > of co

altern

alum

alum

C

m th