

A New Palladium Precatalyst Allows for the Fast Suzuki–Miyaura Coupling Reactions of Unstable Polyfluorophenyl and 2-Heteroaryl Boronic Acids

Tom Kinzel, Yong Zhang, and Stephen L. Buchwald*

Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, Massachusetts 02139, United States

Received August 16, 2010; E-mail: sbuchwal@mit.edu

Abstract: Boronic acids which quickly deboronate under basic conditions, such as polyfluorophenylboronic acid and fivemembered 2-heteroaromatic boronic acids, are especially challenging coupling partners for Suzuki–Miyaura reactions. Nevertheless, being able to use these substrates is highly desirable for a number of applications. Having found that monodentate biarylphosphine ligands can promote these coupling processes, we developed a precatalyst that forms the catalytically active species under conditions where boronic acid decomposition is slow. With this precatalyst, Suzuki–Miyaura reactions of a wide range of (hetero)aryl chlorides, bromides, and triflates with polyfluorophenyl, 2-furan, 2-thiophene, and 2-pyrroleboronic acids and their analogues proceed at room temperature or 40 °C in short reaction times to give the desired products in excellent yields.

The Suzuki–Miyaura cross-coupling (SMC) reaction is arguably the most important and widely used method for the construction of sp^2-sp^2 carbon–carbon bonds.¹ However, since SMCs are typically performed at elevated temperatures and require several hours reaction time, the scope of usable boronic acids is limited to those that do not significantly decompose under standard conditions. Consequently, the SMCs of quickly deboronating² 2,6-difluorophenylboronic acid (1) and analog 2 as well as the SMCs of fivemembered 2-heterocyclic boronic acids 3-8 (Chart 1) are problematic.^{3–6} Being able to couple these boronic acids with functionalized aryl and heteroaryl substrates would be of great interest for the synthesis of pharmaceutical and agrochemical candidates, natural products, and materials.

Chart 1. Boronic Acids That Easily Undergo Protodeboronation in Aqueous Base To Give the Parent Arene



One viable solution to this problem is the masking of the boronic acids as MIDA boronates,⁷ cyclic triolborates,⁸ or trifluoroborate salts^{9,10} which slowly hydrolyze¹¹ to the free boronic acid under the reaction conditions. The ratio of the concentration of catalyst and the boronic acid is relatively high and, as a result, transmetalation is favored over protodeboronation. Nevertheless, these surrogates are typically prepared from the free boronic acids, and often, high palladium loadings and/or long reaction times are

10.1021/ja1073799 © XXXX American Chemical Society

required for efficient coupling. Another solution, albeit limited to aryl bromides and iodides, allows the coupling of 2 and penta-fluorophenylboronic acid by using stoichiometric amounts of Ag₂O in addition to the palladium catalyst.^{12,13}

Herein, we report the fast and efficient SMC of the free boronic acids **1-8** at room temperature or 40 °C in short reaction times of 30 min to 2 h. Key to the success was the development of a new precatalyst that generates the catalytically active LPd(0) species quickly under mild conditions under which the deboronation of the boronic acid is significantly slowed down.

In the course of mechanistic investigations, we prepared complex SPhosPdPhCl (9) that results from oxidative addition of SPhosPd(0) with chlorobenzene¹⁴ and treated it with 1 in the presence of aqueous K_3PO_4 (Scheme 1). Rapid formation of product 10 demonstrated that transmetalation and reductive elimination occur with 1 at room temperature.¹⁵

Scheme 1. Initial Observation of the Coupling of 1 with a Stoichiometric Amount of an Oxidative-Addition Complex 9



Using 2 mol % of **9** as precatalyst, 4-chloroanisole was coupled with **1** to give the desired product in 93% yield in less than 30 min using a 1:2 mixture of THF and 0.5 M aqueous K_3PO_4 . Kinetics studies showed that product formation stopped abruptly, indicating that the remainder of **1** was consumed by competing base- and/or metal-catalyzed deboronation.² Increasing the temperature resulted in a lower conversion. However, full conversion of 4-chloroanisole at room temperature was achieved by using the oxidative-addition complex **11** in which the SPhos ligand is replaced by XPhos (Chart 2).

Chart 2. XPhos-Containing Precatalysts Used in This Study



The syntheses of complexes **9** and **11** are difficult, and their use as precatalysts for the SMC of **1** with any aryl halide necessarily results in the formation of biaryl **10** in addition to the desired product. The preparation and isolation of an individual oxidativeaddition complex for each substrate is clearly impractical and often impossible. Because using $Pd(OAc)_2$ or $Pd(dba)_2$ with XPhos gave unsatisfactory results, we sought another solution to provide the

> J. AM. CHEM. SOC. XXXX, XXX, 000 = A PETITIONER NPC EX. 1020 Page 1 of 3

catalytically active XPhosPd(0) species which undergoes oxidative addition *in situ*.

Previously, we developed precatalyst **12** which, upon deprotonation by a base, forms XPhosPd(0) and indoline via C–N reductive elimination.¹⁶ While it is an excellent source of the catalytically active Pd species, its activation with weak bases is slow and occurs only at elevated temperatures. Moreover, its synthesis requires several steps. In the process of addressing these issues, we found that both drawbacks could be overcome with a new class of precatalysts where the parent aliphatic amine in **12** is replaced by 2-aminobiphenyl in **13**: First, the higher acidity of the palladiumbound aromatic amine compared to the aliphatic amine allows activation of **13** to occur almost instantaneously with weak base at room temperature. Second, **13** is easily obtained in a one-pot procedure by combining Pd(OAc)₂ with 2-aminobiphenyl, followed by addition of LiCl and the ligand (XPhos) (Scheme 2).¹⁷

Scheme 2. Synthesis of Precatalyst 13



For the SMC reaction between 4-chloroanisole and 1 in the presence of aqueous 0.5 M K_3PO_4 as base, the use of precatalysts 12 and 13 was tested. As expected, only the latter promoted the reaction, leading to full conversion to product.

Initially, several aryl halides or pseudohalides were subjected to the SMC reaction with polyfluorophenylboronic acids. We observed that aryl chlorides, bromides, and triflates are good substrates for the coupling with **1**, while only low conversions were observed with aryl iodides¹⁸ (Table 1). Substrates with coordinating groups in the ortho position, such as esters or ketones, could also not be coupled efficiently. However, even large noncoordinating ortho substituents, as in **15b**, were well tolerated. Heteroaromatic compounds readily underwent the SMC to form the desired products in excellent yields, but five-membered chloro- or bromoheteroarenes with multiple heteroatoms remained problematic.

Table 1. Coupling of Polyfluoroboronic Acids to Aryl Chlorides, Bromides, and Triflates^a



^{*a*} ArX (1 mmol), ArB(OH)₂ (1.5 mmol), **13** (2 mol %), degassed THF (2 mL), degassed 0.5 M aq K_3PO_4 (4 mL), rt, 30 min; isolated yields, average of two runs. ^{*b*} **13** (3 mol %).

B J. AM. CHEM. SOC. VOL. xxx, NO. xx, XXXX

Interestingly, we found that, while trisubstituted polyfluorophenylboronic acids 2 and 14 could be employed to give products 16 and 17, very low conversion was observed with 2,3,6-trifluorophenylboronic acid. To understand this result, we studied both the deboronation rates¹⁹ and apparent transmetalation rates²⁰ of (poly)fluorophenylboronic acids (Table 2). Transmetalation efficiency increases with the number of fluorine substituents, with ortho substitution having the greatest impact. Thus, 1 reacts about 150 times faster than simple phenylboronic acid, and almost 4 times faster than 2-fluorophenylboronic acid. In the series of $1 \rightarrow 2 \rightarrow 14$, the loss in stability is counterbalanced by higher transmetalation rates, thus allowing the coupling of 2 and 14 under essentially the same reaction conditions. In contrast, the coupling of 2,3,6trifluorophenylboronic acid is more problematic because the deboronation rate is dramatically increased, whereas the transmetalation rate does not change relative to that of 1.²¹

Table 2. Deboronation Rates and Apparent Relative Transmetalation Rates of Polyfluorophenylboronic Acids

entry	boronic acid	deboronation rate, $t_{1/2}$ (min) ^a	relative transmetalation rate, $k_{\rm rel}$
1	PhB(OH) ₂		1
2	2-F		42
3	3-F		2
4	4-F		5
5	$2,6-F_2(1)$	28	155
6	$2,4,6-F_3$ (2)	10	202
7	2,6-F ₂ -3-OnBu (14)	6	536
8	2,3,6-F ₃	2	156

^{*a*} Deboronation due to the presence of base was measured via calorimetry at room temperature in a mixture of THF and 0.5 M K_3PO_4 . The values in the column are the half-lives of the observed first-order kinetic profiles.

It is important to note that the coupling of these polyfluorophenylboronic acids proceeds by using established ligands (SPhos, XPhos) that exhibited excellent results in other SMC reactions. Thus, the previously reported unsuccessful attempts³ to couple **1** and **2** in the presence of these ligands were not due to their structure but to the reaction conditions which were necessary to provide the active catalytic species. Using precatalyst **13**, the catalytically active XPhosPd(0) species is formed rapidly at room temperature, thereby allowing the successful coupling of these unstable boronic acids.

Next, we turned our attention to heteroaryl boronic acids and found that, by using 2 mol % of 13, 2-heterocyclic boronic acids 3–8 could be coupled efficiently with a wide array of aryl and benzyl halides, mostly within 30 min at room temperature or 40 °C (Table 3). The higher stability of the 2-heterocyclic boronic acids relative to that of the polyfluorophenylboronic acids allowed the transformations of challenging substrates that could not be used with the latter. Therefore, performing the reaction at 40 °C for 2 h allowed the conversion of 4- and 5-halopyrazoles, 2-chlorobenzoxindole as well as 2-chloroacetanilide to give the coupled products 19a, 20c, 22b, and 20b in excellent yields.

The fast generation of XPhosPd(0) from precatalyst 13 is a prerequesite for the successful coupling of unstable boronic acids 1-8 but also increases the rate of reaction for other SMC reactions, as illustrated by the fast coupling of 3-furan and 3-thiopheneboronic acids to give 24 and 25.

In summary, we have developed a procedure for very fast SMC reactions at room temperature or 40 °C that allows the coupling of a wide range of (hetero)aryl halides and triflates with excellent functional group tolerance. The fast catalytic process and the extremely mild reaction conditions make the coupling of unstable polyfluorophenyl and five-membered 2-heterocyclic boronic acids

PETITIONER NPC EX. 1020 Page 2 of 3

Table 3. Coupling of 2- and 3-Heterocyclic Boronic Acids to Aryl and Benzyl Chlorides and Aryl Bromides⁴



^a Reagents and conditions: ArX (1 mmol), ArB(OH)₂ (1.5 mmol), 13 (2 mol %), degassed THF (2 mL), degassed 0.5 M aq K₃PO₄ (4 mL), rt or 40 °C, 30 min or 2 h; isolated yields, average of two runs.

with a wide range of aryl (pseudo)halides possible. The rate increase in comparison to that of typical SMC reactions relies on fast generation of the catalytically active species from an easy-toprepare, air- and moisture-stable precatalyst.

Acknowledgment. We thank the NIH for financial support of this project (GM46059) and BASF for a gift of palladium compounds. T.K. thanks the Alexander von Humboldt Foundation for a Feodor Lynen postdoctoral fellowship. We also thank Dr. Jaclyn Henderson for assistance with some experiments. The Varian NMR instrument used was supported by the NSF (Grants CHE 9808061 and DBI 9729592).

Supporting Information Available: Complete refs 4a and 4b, experimental procedures, and product characterization. This material is available free of charge via the Internet at http://pubs.acs.org.

References

- (1) (a) Miyaura, N.; Yamada, K.; Suzuki, A. *Tetrahedron Lett.* 1979, *36*, 3427.
 (b) Miyaura, N. *J. Organomet. Chem.* 2002, *653*, 54. (c) Miyaura, N. In Metal-Catalyzed Cross-Coupling Reactions; Diederich, F., de Meijere, A., Eds.; Wiley-VCH: New York, 2004.
- (2) Both base- and metal-catalyzed deboronation of boronic acids may occur under typical SMC reactions. (a) Kuivila, H. G.; Reuwer, J. F.; Mangravite, J. A. Can. J. Chem. 1963, 41, 3081. (b) Kuivila, H. G.; Reuwer, J. F.; Mangravite, J. A. J. Am. Chem. Soc. 1964, 86, 2666.

- (3) While 2,3- and 2,4-difluorophenylboronic acids could be coupled using ligand SPhos at 80–90 °C, these conditions did not allow the efficient coupling of **1** or **2**: Barder, T. E.; Walker, S. D.; Martinelli, J. R.; Buchwald, S. L. J. Am. Chem. Soc. 2005, 127, 4685.
- For some examples for the coupling of 2,6-difluorophenylboronic acid with vinyl triflates and aryl bromides, see: (a) Antonow, D.; et al. J. Med. Chem. 2010, 53, 2927. (b) Schmidt, D.; et al. Bioorg. Med. Chem. Lett. 2009, 19, 4768. (c) Palmer, B. D.; Smaill, J. B.; Rewcastle, G. W.; Dobrusin, E. M.; Kraker, A.; Moore, C. W.; Steinkampf, R. W.; Denny, W. A. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 1931.
- (a) Knapp, D. M.; Gillis, E. P.; Burke, M. D. J. Am. Chem. Soc. 2009, (3) 131, 6961. (b) Billingsley, K. L.; Buchwald, S. L. J. Am. Chem. Soc. 2007, 129, 3358. (c) Billingsley, K. L.; Anderson, K. W.; Buchwald, S. L. Angew. Chem., Int. Ed. 2006, 45, 3484. (d) Tyrell, E.; Brookes, P. Synthesis 2003,
- (6) For some representative examples of the coupling of five-membered For some representative examples of the coupling of hve-membered 2-heteroaromatic boronic acids, see: (a) Kabri, Y.; Gellis, A.; Vanelle, P. *Eur. J. Org. Chem.* **2009**, *24*, 4059. (b) Gill, G. S.; Grobelny, D. W.; Chaplin, J. H.; Flynn, B. L. *J. Org. Chem.* **2008**, *73*, 1131. (c) Organ, M.; Çalimsiz, S.; Sayah, M.; Hoi, K.; Lough, A. Angew. Chem., Int. Ed. **2009**, *48*, 2383. (d) Dang, Y.; Chen, Y. *J. Org. Chem.* **2007**, *72*, 6901. (e) Maeda, H.; Haketa, Y.; Nakanishi, T. *J. Am. Chem.* **50c**, **2007**, *129*, 13661. (f) James, C. A.; Coelho, A. L.; Gevaert, M.; Forgione, P.; Snieckus, V. J. Org. Chem. 2009, 74, 4094.
- 2-heterocyclic MIDA boronates were successfully employed for SMCs (see ref 4a), but polyfluorophenyl MIDA boronates were not described. MIDA = N-methyliminodiacetic acid.
- Yamamoto, Y.; Takizawa, M.; Yu, X.; Miyaura, N. Angew. Chem., Int. (8)Ed. 2008, 47, 928.
- Polyfluorophenyl BF₃K salts were coupled to 4-bromobenzonitrile: Molander, G. A.; Biolatto, B. J. Org. Chem. **2003**, 68, 4302. (9)
- Coupling of 2-heterocyclic BF₃K salts: Molander, G. A.; Canturk, B.; (10)
- Coupling of 2-heterocyclic B-3A sans. Morander, G. A., Canturk, B., Kennedy, L. E. J. Org. Chem. 2009, 74, 973.
 The hydrolysis of BF₃K salts to the free boronic acid likely occurs prior to cross-coupling: Butters, M.; Harvey, J. N.; Jover, J.; Lennox, A. J. J.; Lloyd-Jones, G. C.; Murray, P. Angew. Chem., Int. Ed. 2010, 49, 5156.
- (12) (a) Chen, J.; Cammers-Goodwin, A. Tetrahedron Lett. 2003, 44, 1503. (b) Korenaga, T.; Kosaki, T.; Fukumura, R.; Ema, T.; Sakai, T. Org. Lett. 2005, 7, 4915. (c) Adonin, N. Y.; Babushkin, D. E.; Parmon, V. N.; Bardin, V. V.; Kostin, G. A.; Mashukov, V. I.; Frohn, H. Tetrahedron 2008, 64, 5920.
- (13) It should be mentioned that with copper or palladium catalysts, aryl bromides and chlorides can be directly coupled to polyfluorobenzenes via C-Hactivation processes, but high temperatures and long reaction times are required. (a) Lafrance, M.; Shore, D.; Fagnou, K. Org. Lett. 2006, 8, 5097. (b) Do, H.; Daugulis, O. J. Am. Chem. Soc. 2008, 130, 1128.
- (14) Barder, T. E.; Biscoe, M. R.; Buchwald, S. L. Organometallics 2007, 26, 2183
- (15) The reductive elimination of 4-methoxy-2',6'-difluorobiphenyl from an isolated Pd(II) complex has been reported: Osakada, K.; Onodera, H.; Nishihara, Y. Organometallics 2005, 24, 190.
- (16) Biscoe, M. R.; Fors, B. P.; Buchwald, S. L. J. Am. Chem. Soc. 2008, 130, 6686.
- Albert, J.; Granell, J.; Zafrilla, J.; Font-Bardia, M.; Solans, X. J. Organomet. (17)Chem. 2005, 690, 422.
- (18) The rate of the SMC reaction under the described conditions decreases in the order ArCl > ArBr > ArI. This order is reproduced in stoichiometric transmetalation studies starting from oxidative-addition complexes. Furthermore, we found an inhibiting effect on the transmetalation rate by additional halide ions, in the order $I^- > Br^- > Cl^-$
- (19) The deboronation rates of some polyfluorophenylboronic acids in a D₂O/ pyridine mixture at elevated temperatures were reported: Frohn, H.; Adonin, N. Y.; Bardin, V. V.; Starichenko, V. F. Z. Anorg. Allg. Chem. 2002, 628, 2834.
- (20) Apparent relative transmetalation rates were determined by competition experiments using an excess of a mixture of 2-methoxyphenylboronic acid and a second boronic acid with 4-chloroanisole as substrate in the presence of 20 mol % 13. The obtained product ratios were normalized to the product ratio found for the competition experiment between 2-methoxyphenylbo-ronic acid and phenylboronic acid. To ensure that the measured values reflect the transmetalation of the free boronic acids and not of boronic acid anhydrides or boroxines present in the starting material, we modified the experiment involving phenylboronic acid by premixing the boronic acid mixture with aqueous K₃PO₄ solution prior to addition of the precatalyst and aryl chloride, and we found the same ratio of products as before.
- (21) The attempted coupling of 2,3,6-trifluorophenylboronic acid with 4-chloroacetophenone under microwave irradiation led exclusively to the deboronated compound, trifluorobenzene: Clarke, M. L.; France, M. B.; Fuentes, J. A.; Milton, E. J.; Roff, J. R. Beilstein J. Chem. 2007, 3, 18.

JA1073799

J. AM. CHEM. SOC. = VOL. xxx, NO. xx, XXXX C PETITIONER NPC EX. 1020 Page 3 of 3