



## Proven symptom reduction for your HAE attacks

In clinical trials, patients treated with KALBITOR showed significant improvement in hereditary angioedema (HAE) attack symptoms at 4 hours.

[LEARN MORE ABOUT KALBITOR › \(/how-kalbitor-works\)](#)

Please see **Full Prescribing Information**  
([http://www.shirecontent.com/PI/PDFs/Kalbitor\\_USA\\_ENG.pdf](http://www.shirecontent.com/PI/PDFs/Kalbitor_USA_ENG.pdf)), including  
**Boxed Warning and Medication Guide** ([http://www.shirecontent.com/?  
product=Kalbitor&country=USA&language=ENG&contenttype=medguide](http://www.shirecontent.com/?product=Kalbitor&country=USA&language=ENG&contenttype=medguide)).



## Patient Ambassador Events

Attend a live event and hear from someone with HAE. You can learn about his or her path to diagnosis and how they manage life with HAE.

**FIND AN EVENT NEAR YOU ›**

(/events)



## OnePath® Product Support Services

OnePath is here to support you throughout your treatment journey with KALBITOR.

**LEARN MORE ›**

(/onepath)



## Sign Up & Stay in Touch

Register to receive more information from Shire about HAE.

**SIGN UP TODAY ›**

(/join)

## Important Safety Information

### What is the most important information I should know about KALBITOR?

Serious allergic reactions may happen in some people who receive KALBITOR. These allergic reactions can be life-threatening and usually happen within 1 hour after receiving KALBITOR.

- KALBITOR should be given to you by a doctor or nurse in a healthcare setting where serious allergic reactions and hereditary angioedema (HAE) can be treated.
- Symptoms of a serious allergic reaction to KALBITOR can be similar to the symptoms of HAE, the condition that you are being treated for. Your doctor or nurse should watch you for any signs of a serious allergic reaction after treatment with KALBITOR.
- **Tell your doctor or nurse right away if you have any of these symptoms of a serious allergic reaction during or after treatment with KALBITOR:**
  - wheezing, shortness of breath, cough, chest tightness, or trouble breathing
  - dizziness, fainting, fast or weak heartbeat, or feeling nervous
  - reddening of the face, itching, hives, or feeling warm
  - swelling of the throat or tongue, throat tightness, hoarse voice, or trouble swallowing
  - runny nose, nasal congestion, or sneezing

### Who should not receive KALBITOR?

Do not receive KALBITOR if you are allergic to KALBITOR. Before receiving KALBITOR, tell your doctor if you have ever had an allergic reaction to KALBITOR.

### What else should I tell my doctor before I receive KALBITOR?

Tell your doctor if you are pregnant, plan to become pregnant, breast-feeding, or plan to breast-feed. It is not known if KALBITOR will harm your unborn baby or pass into your breast milk.

### What are the possible side effects?

Common side effects of KALBITOR include headache, nausea, diarrhea, fever, injection site reactions, such as redness, rash, swelling, itching, or bruising, and stuffy nose. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) (<http://www.fda.gov/medwatch>), or call 1-800-FDA-1088.

Talk to your doctor for more information about your medical condition or treatment.

Please see the Full Prescribing Information  
([http://www.shirecontent.com/PI/PDFs/Kalbitor\\_USA\\_ENG.pdf](http://www.shirecontent.com/PI/PDFs/Kalbitor_USA_ENG.pdf)), including Boxed Warning and  
Medication Guide ([http://www.shirecontent.com/?  
product=Kalbitor&country=USA&language=ENG&contenttype=medguide](http://www.shirecontent.com/?product=Kalbitor&country=USA&language=ENG&contenttype=medguide)).



## Hereditary Angioedema

TM or ® items and the associated logos are trademarks or registered trademarks of Shire or its affiliates.

© 2017 Shire US Inc., Lexington, MA 02421 1-800-828-2088

All rights reserved. SHIRE and the Shire Logo are trademarks or registered trademarks of Shire  
Pharmaceutical Holdings Ireland Limited or its affiliates.

S28528 06/17