The Role of More than 40 Years of Improvement in Protein A Chromatography in the Growth of the Therapeutic Antibody Industry

Glen R. Bolton

Amgen, 360 Binney Street, Cambridge, MA 02142

Krunal K. Mehta

Amgen, 360 Binney Street, Cambridge, MA 02142

DOI 10.1002/btpr.2324

Published online July 23, 2016 in Wiley Online Library (wileyonlinelibrary.com)

Protein A chromatography has been used as the mAb capture step in the majority of FDA submissions. In this study, the performance of protein A chromatography, as indicated by capacity, operational flow rate, and productivity (rate of mAb production per liter of resin) was examined over its full history to gain insights into the reasons for its consistent use. Protein A productivity and capacity have increased 4.3 and 5.5% a year, respectively, since 1978. In contrast, protein A operational flow rate increased between 1978 and 2001 and then remained constant or declined as further improvements provided only marginal benefits. The productivity of protein A resin and also the mAb bioreactor titer (14% growth) rapidly improved starting in about 1990 to economically provide material for clinical trials. Technology improvement is typically driven by product sales. The sales of protein A resin, as indicated by sales of protein A ligand (21% growth), have closely paralleled the sales of mAbs (20% growth). Both increased rapidly in 2000 after the first major mAb therapeutics were approved and the markets were developed. It is likely that alternatives to protein A chromatography have not been implemented because of the order of magnitude improvement in protein A performance. Protein A membrane adsorbers and monoliths have higher productivity than packed columns due to their short bed heights and high operational flow rates. These devices are not currently practical for large-scale manufacturing but may represent a format for future improvements in protein A productivity. © 2016 American Institute of Chemical Engineers Biotechnol. Prog., 32:1193-1202, 2016

Keywords: protein A, SpA, mAb, monoclonal antibody, affinity chromatography, purification

Introduction

The standard process for the purification of a mAb from harvested cell culture fluid typically involves at least two chromatography steps: protein A affinity and ion-exchange chromatography. Protein A chromatography separates proteins from impurities based on a reversible interaction between the Fc portion of a molecule and a protein A ligand immobilized on a chromatography matrix.¹ Impurities flow through the matrix while the mAb stays bound to the column. The column is subsequently washed to achieve additional impurity removal.² Product is eluted from the matrix by a reduction in pH. Subsequently, the mAb is typically either bound to a cation-exchange chromatography resin or flowed through an anion-exchange resin to remove additional host-cell proteins, DNA, leached protein A, media derived impurities, potential viruses, and aggregated species.³ This is often followed by an additional polishing ion-exchange or hydrophobic interaction chromatography step.⁴

The binding of protein A to a wide range of human and rabbit serum antibodies was first observed by Jensen,⁵ who

attributed the binding to natural antibody specificity for a bacterial antigen (antigen A). Subsequently, the antigen was incorrectly described as a bacterial polysaccharide⁶ and then correctly shown to be a bacterial cell wall protein.⁷ This protein was first called protein A in 1964 to distinguish it from other polysaccharides and proteins.⁸

Subsequently, the theory of natural antibody specificity for a bacterial antigen was disproven. Protein A was shown to bind to antibodies lacking bacterial antigen specificity and also to bind to Fc portions of antibodies.⁹ In addition, it was shown that protein A bound more strongly to Fc fragments of antibodies than Fab regions or light chains. Subsequently, use of protein A immobilized on a solid support for the affinity purification of antibodies was developed¹⁰ and patented.¹¹ The first protein A resin was commercialized in 1978.¹²

Protein A will bind to a range of antibody isotypes derived from different species.¹³ The first therapeutic antibody, Orthoclone, was approved in 1986 and utilized protein A chromatography as a capture step in the purification process.¹⁴ Protein A was subsequently used as the mAb capture step in 86% of FDA submissions.¹⁵ Early protein A resins were expensive and had limited capacity, flow rate, and life-time. A number of studies have described replacing the

Correspondence concerning this article should be addressed to G. Bolton at gbolton@amgen.com.

protein A chromatography step with alternative purification methods. Ion-exchange chromatography is a frequently discussed alternative mAb capture strategy.^{15–18} Antibody or impurity precipitation^{4,14,19,20} or crystallization²¹ have also been suggested. Alternative protein²² or synthetic molecule²³ affinity resins have also been proposed to replace protein A.

However, since 1978, both the performance and availability of protein A resins have steadily increased. Newer beads are more rigid yet highly porous, allowing fast flow rates, low pressure drops, and high mAb capacities.²⁴ The sizes of both beads and bead pores have been optimized to improve performance. A number of base bead chemistries including cross-linked agarose,²⁵ controlled pore glass,²⁶ polystyrenedivinylbenzene,²⁴ ceramic material,²⁷ polymethacrylate,²⁸ cross-linked cellulose,²⁹ and polyvinylether³⁰ have been developed to improve protein A performance.

Protein A resins have benefited from the removal of animal derived production materials.³¹ Improvements in how protein A ligands are immobilization on beads, e.g., via a single or multipoint linkage,²⁴ has helped improve ligand accessibility and increase capacity. Changes to the number of protein A domains and the amino acids in those domains has led to improved protein A capacity, lifetime, caustic stability, elution pH, and mAb Fc specificity.^{32–34} These improvements have led to an improvement in the amount of mAb that can be produced per liter of protein A resin per hour (productivity).

In this study, the growth of protein A productivity, capacity, and operational flow rate is tracked using available literature data. This comparison provides insights into how fast the performance of protein A chromatography has improved and how this improvement compares to the rate of improvement in bioreactor titers. In order to get insights into the timing and reasons for current and potential future improvements in protein A chromatography, the growth of protein A performance is then compared to the growth of protein A resin sales and therapeutic mAb sales.

Review of Protein A Evolution

Data from citable studies that provide the protein A column bed height, flow rate, and capacity at 10% breakthrough were compiled and used to evaluate growth in protein A productivity, capacity, and operational flow rate. The resin launch date was obtained from the resin vendors or estimated from the earliest date of publication of resin information.

The flow rate used to obtain the reported capacity data (operational flow rate) was compiled. The maximum flow rate is not summarized here because in some studies only maximum pressure is reported, and in other studies maximum flow rate values are provided without also providing the corresponding capacity data.

There is limited data available for some of the key attributes of protein A media. Unfortunately, pricing information is rarely published by vendors or customers³⁵ and therefore was not evaluated in this study. The global protein A media market was estimated to generate annual revenues between 350 and 400 million USD annually in the 2015 Repligen annual report. In addition, limited data are available for resin lifetime, nonspecific product and impurity adsorption,² ligand leaching, and reproducibility

The high pH stability of protein A ligands has improved through selective mutation of the protein A ligand.³³ The **Page 2 of 10**

chemical stability data provided in the references are listed in Table 1. However, very few references provide the expected loss in performance based on exposure to a specific pH value for a fixed duration. This makes it difficult to measure the rate at which chemical stability has improved.

Productivity is defined as the grams of mAb that can be purified per hour on a liter of protein A resin. Most, but not all studies provided the antibody load concentration. Therefore, a mAb titer of 1 g/L was assumed during loading. The wash, elution, and regeneration steps were assumed to be 10, 3, and 5 column volumes (CV), respectively. The total CVs per run were determined from the reported resin capacity, the bioreactor titer, the number of wash CV, the number of elution CV, and the number regeneration CV according to Eq. 1:

$$Total \ CV = \frac{Resin \ Capacity}{Bioreactor \ Titer} + Wash \ CV + Elution \ CV + Regen \ CV$$
(1)

The productivity was then calculated from the total CV using the reported resin capacity, linear flow rate, and bed height according to Eq. 2:

$$Productivity = \frac{Resin\ Capacity * Linear\ Flow\ Rate}{Total\ CV * Bed\ Height}$$
(2)

The amount and the quality of protein A performance data varied. Data for early protein A resins typically had only one or a few low capacity and/or low flow rate data points. Therefore, it was not possible to compare all resins at a single reference flow rate, or to compare flow rates at a single reference capacity. Therefore, data from all flow rates reported in each study are used to plot and evaluate the growth in productivity, capacity, and operational flow rate. Some of the vendor literature provided limited experimental details and may have utilized conditions favorable to their products. One study utilized an IgG3, which may have impaired the capacity of some resins.²⁵ The load antibody concentration ranged from 0.33 to 10 mg/mL and was not reported in some studies. Higher concentration has been reported to improve⁴⁹ or not affect protein A capacity.^{43,54,58} A load concentration of 1 mg/mL was assumed to calculate productivity. The bed heights used in the studies ranged from 2 to 30 cm, with lower bed heights typically allowing higher flow rates and productivity. Lower bed heights however are less representative of a mAb manufacture process and can result in lower capacities.

Protein A productivity growth

The resin productivity is plotted vs. resin launch year in Figure 1. A compounded annual productivity growth rate r of 4.3 \pm 1.4% and an initial value Y_0 of 5.8 \pm 2.4 g/L/h was determined by fitting all data points except the membrane adsorber and monolith data to Eq. 3, where t is the number of years:

$$Y = Y_0 (1+r)^t$$
(3)

The growth rate was somewhat sensitive to the assumed titer but not the assumed CV for the various process steps. Assuming a load titer of 0.1 or 10 g/L, instead of 1 g/L, changed the growth rate to either 2.4 ± 1.2 or $7.7 \pm 1.8\%$. Doubling the assumed wash, elution, and regeneration CV did not alter the productivity growth rate as all data was changed to a similar degree.

Table 1. Protein A Launch	Date, Bed Height, Ope	rational Flo	w Rate, Capacity	at 10% Breakt	through, Produc	ctivity, Feed Co	omposition, and Caustic Stability	*	
Protein A Resin	Manufacturer	Launch Year	Capacity (g/L) at 10% BT	Flow Rate (cm/h)	Productivity (g/L/h)	Bed Height (cm)	Feed Information	Caustic Stability	Reference
Protein A Sepharose TM	GEHC	1978	5.0	100	0.7	30	Mouse Serum IgG		12
Protein A Sepharose TM	GEHC	1978	10.0	09	4.3	5	0.5 mg/mL Monoclonal		25
Protein A Ultrogel	Pall/IBF	1982	0.7	09	0.4	5	0.5 mg/mL Monoclonal		25
Protein A Sepharose TM	GEHC	1983	20.1	09	6.3	5	mouse IgO5 at рн 8.9 0.5 mg/mL Monoclonal		25
o FF Affi-Gel [®] Protein A	Bio-Rad	1984	12.8	09	5.0	S.	mouse 1905 at pH 5.9 0.5 mg/mL Monoclonal		25
$\operatorname{ProSep}^{\otimes}$ -A	Millipore	1988	10.0	09	4.3	5	mouse IgG3 at pH 8.9 0.5 mg/mL Monoclonal		25
Eupergit [®] C Protein A	Rohm Pharma	1989	0.5	60	0.3	S	mouse IgG3 at pH 8.9 0.5 mg/mL Monoclonal		25
Protein A Sepharose TM 4 FF	GEHC	1989	18.5	09	6.1	5	0.5 mg/mL Monoclonal mouse loc3 at nH 8.0		25
Affi-Prep [®] Protein A nProtein A Sepharose TM	Bio-Rad GEHC	1990 1991	11.0 6.0	543 300	48.8 15.0	4.2 5	1 mg/mL Humanized mAb 0.33 mg/mL Human IgG	Short-term pH 2-10	36 37
4.rr nProtein A Sepharose TM 4.rr	GEHC	1991	10.0	200	14.3	3	0.33 mg/mL Human IgG	Short-term pH 2-10	37
4гг nProtein A Sepharose TM лтг	GEHC	1991	17.0	100	9.7	5	0.33 mg/mL Human IgG	Short-term pH 2-10	37
nProtein A Sepharose TM 4FF	GEHC	1991	24.0	30	3.4	Ś	0.33 mg/mL Human IgG	Short-term t pH 2-10	37
ProSep®-A rProtein A Sepharose TM	Millipore GEHC	1993 1995	16.0 50.0	$1,000 \\ 100$	29.4 7.4	16 10	0.8 mg/mL Chimeric antibody 0.5 mg/mL Human IgG pH 7.0	Cleaning in place pH 2-11	26 38
(4) FF	GEHC	1995	42.0	200	14.0	10	0.5 mg/mL Human IgG pH 7.0	Cleaning in place pH 2-11	38
rProtein A Sepharose TM (4) FF	GEHC	1995	31.0	300	19.0	10	0.5 mg/mL Human IgG pH 7.0	Cleaning in place pH 2-11	38
StreamLine TM A Poros [®] LP	GEHC Thermo Fisher	1998 1998	11.0 16.0	500 500 500	19.0 23.5	10	0.35 mg/mL Humanized mAb 0.35 mg/mL Humanized mAb		2 2 2 4 4 2
rotos JUA rmp Protein A Senharose TM 4FF	GEHC	2000	20.0	200	25.7	4.1	5 mg/mL Human IgG pH 7.4	Cleaning in place 2-11	31
rmp Protein A Senharose TM 4FF	GEHC	2000	32.0	100	15.6	4.1	5 mg/mL Human IgG pH 7.4	Cleaning in place 2-11	31
rmp Protein A Sepharose TM 4FF	GEHC	2000	41.0	60	10.2	4.1	5 mg/mL Human IgG pH 7.4	Cleaning in place 2-11	31
Protein A Ceramic HvnerD [®] F	Pall	2000	30.0	100	5.3	12	2.55 mg/mL Human IgG	Cleaning pH 2–13	27
MabSelect TM MabSelect TM	GEHC GEHC	2001 2001	29.0 28.0	171 300 200	5.3 9.1	20 20 20	3 mg/mL IgG 3 mg/mL IgG		39 39
MabSelect MabSelect TM MabSelect TM	GEHC GEHC	2001 2001 2001	20.0 10.0 50.0	000 1,500 167	26.8 26.8 12.3	20 10	o mg/mL igo 3 mg/mL igo 1.8 mg/mL mAb		90 39 40
Prosep [®] -VA HiCap	Millipore	2003	30.0	500	16.4	19	1 mg/mL Polyclonal human IgG		41
Prosep [®] -VA Ultra	Millipore	2004	18.0	1,200	30.0	20	2 mg/mL mAb		42

	Protein A Resin	Manufacturer	Launch Year	Capacity (g/L) at 10% BT	Flow Rate (cm/h)	Productivity (g/L/h)	Bed Height (cm)	Feed Information	Caustic Stability	Reference
	Prosep [®] -VA Ultra	Millipore	2004	26.0	600	17.7	20	2 mg/mL mAb		42
	Prosep®-VA Ultra	Millipore	2004	34.0	300	9.8	20	2 mg/mL mAb		42
	Prosep [®] -VA Ultra	Millipore	2004	37.0	200	6.7	20	2 mg/mL mAb		42
With the constant of th	MabSelect SuRe TM	GEHC	2005	20.0	600	31.6	10	Human IgG		43
With Solution (Market Solution) GETC (Market Solution) Open (Market Solution) Monter (Market Solution) Monter (Market Solution) Monter (Market Solution)	MabSelect Suke	GEHC	5002	39.0 40.0	001	1./1	10	Human IgG		4. 2. c.
	MabSelect SuRe TM	GEHC	2002	49.0 50.0	75	י. א איי	10	Human IoG		0 4 4 0 6 6
Mache Control	Mebeloce June		2002	0.00	51	0.0	00			0.0
Massensity, international and the state of the	MabSelect Atra MabSelect Xtra TM	CEHC	2002	40.0	300	9.0 10.3	07	2 mg/mL IgG 3 mg/mL IgG		90 05
	Mabselect Alla Mabselect Vina TM		5002	40.0 34.0	000	2.01 70 f	07	2 mg/mL IgO		90 02
Matrix Matrix	MabSelect Alfa MabSelect Vira TM	CEHC	2005	0.4.0 0.00	1 500	30.5	07	2 mg/mL IgG		90 05
	Sartobind [®] Protein A	Sartorius	2005	7.5	120	88.2	0.4 0.4	1 mg/mL Polyclonal human	Short term pH 2-10	44 44
Porce: MatCipute ¹¹ Themo Fisher 201 430 700 241 20 196 in Fisher 201 450 Porce' MatCipute ¹¹ Themo Fisher 201 450 700 241 20 166 in Fisher 201 45 Porce' MatCipute ¹¹ Themo Fisher 201 450 400 144 20 5 gen Flystenal human 45 Porce' MatCipute ¹¹ Themo Fisher 201 400 144 20 5 gen Flystenal human 45 Porce' MatCipute ¹¹ Themo Fisher 201 400 144 20 5 gen Flystenal human 45 Porce' MatCipute ¹¹ MatCipute ¹¹ 201 400 144 20 5 gen Flystenal human 45 Porce' MatCipute ¹¹ MatCipute ¹¹ 201 401 41 41 41 Porce' MatCipute ¹¹ MatCipute ¹¹ 201 20 301 41 42 Porce' MatCipute ¹¹ MatCipute ¹¹ 201 20 20 5 mg/m. Migc in Pis 41	(Membrane Adsorber)							IgG	- - 	: !
	Poros [®] MabCapture ^{1,14}	Thermo Fisher	2007	43.0	700	24.7	20	5 mg/mL Polyclonal human Inc. in pRS		45
	Poros [®] MabCapture TM A	Thermo Fisher	2007	45.0	500	17.9	20	5 mg/mL Polyclonal human IoG in DRS		45
	Poros [®] MabCapture TM	Thermo Fisher	2007	46.0	400	14.4	20	5 mg/mL Polyclonal human Infi in DRS		45
	Poros [®] MabCapture TM	Thermo Fisher	2007	49.0	200	7.3	20	5 mg/mL Polyclonal human		45
ProSep [®] Unia Plus Millipre 208 3.0 1.200 4.07 20 2 mg/nL mAb 4.2 ProSep [®] Unia Plus Millipre 208 5.0 0.0 1.4 20 2 mg/nL mAb 4.2 ProSep [®] Ulua Plus Millipre 208 5.0 0.00 1.4 20 2 mg/nL mAb ProSep [®] Ulua Plus Millipre 208 5.0 100 1.7 20 2 mg/nL mAb ProSep [®] Ulua Plus Bio-Rad 208 2.0 100 17 5 1 mg/nL Polyclonal huma 47 ProSep [®] Ulua Plus Bio-Rad 208 2.00 190 10 1 mg/nL Polyclonal huma 47 UNOphere SUPA TM Bio-Rad 208 2.00 100 10 1 mg/nL Polyclonal huma 47 UNOphere SUPA TM Bio-Rad 208 2.00 190 10 1 mg/nL Polyclonal huma 47 UNOphere SUPA TM Bio-Rad 208 2.00 180 10 1 mg/nL Polyclonal huma 47 <	A Bio-Monolith Protein A (Monolith)	Agilent Technologies	2008	8.0	297	184.6	0.5	0.5 mg/mL hIgG in PBS	Cleaning in place nH 2 to 13	46
	ProSep [®] Ultra Plus	Millipore	2008	38.0	1,200	40.7	20	2 mg/mL mAb		42
Rosep Ullipore 2008 500 714 20 2 mg/mL mAb 42 ProSep ⁶ Uhra Plus Millipore 2008 500 717 5 1 mg/mL Polyclonal human 42 ProSep ⁶ Uhra Plus Bio-Rad 2008 500 150 177 5 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 230 300 11.0 20 2 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 150 30 18.0 400 400 5 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 208 270 300 18.0 10 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 208 270 300 18.0 10 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 208 270 300 18.0 10 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 20	ProSep [®] Ultra Plus	Millipore	2008	45.0	600	21.4	20	2 mg/mL mAb		42
	ProSep [®] Ultra Plus	Millipore	2008	50.0	300	11.0	20	2 mg/mL mAb		42
Oxyophere SUPA TM Bio-Rad 200 100 110 100 110 100 111 110 110 110 110 110 110 110 110 110 110 110 110 110 110 111	ProSep [©] Ultra Plus	Millipore Bio Dod	2008	52.0 26.0	200 150	7.4	20 *	2 mg/mL mAb 1 mg/mL Dolvelonal human	Working nH ronge 2 11	42
$ \begin{array}{ c c c c c c c c c c c c c c c c c c $	VI IOG SISIIdeONIO	D10-1/40	2000	0.02	001	1.1.1	Û	I IIIS/IIIL I UIYCIVIIAI IIMIIAII IgG	11-C Shall Lid gillan w	Ť
UNOsphere SUPA TM Bio-Rad 2008 18.0 400 5 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 29.0 150 9.3 10 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 27.0 300 18.0 10 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 24.0 400 22.9 10 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 32.0 150 4.8 20 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 31.0 300 9.5 20 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 31.0 300 9.5 20 1 mg/mL Polyclonal human 47 UNOsphere SUPrA TM Bio-Rad 2008 21.0 1 mg/mL Polyclonal human 167	UNOsphere SUPrA TM	Bio-Rad	2008	23.0	300	33.7	S	1 mg/mL Polyclonal human IoG		47
UNOsphere SUPA TM Bio-Rad 2008 29.0 150 9.3 10 $1 \frac{100}{100}$ I Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008 27.0 300 18.0 10 1 mg/mL Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008 27.0 300 18.0 10 1 mg/mL Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008 32.0 150 4.8 20 1 mg/mL Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008 31.0 300 9.5 20 1 mg/mL Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008 31.0 300 9.5 20 1 mg/mL Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008 31.0 300 9.5 20 1 mg/mL Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008 21.0 1 mg/mL Polyclonal human 47 UNOsphere SUPA TM Bio-Rad 2008	UNOsphere SUPrA TM	Bio-Rad	2008	18.0	400	40.0	5	1 mg/mL Polyclonal human		47
UNOsphere SUPrATMBio-Rad200827.030018.010111 <td>UNOsphere SUPrATM</td> <td>Bio-Rad</td> <td>2008</td> <td>29.0</td> <td>150</td> <td>9.3</td> <td>10</td> <td>1 mg/mL Polyclonal human</td> <td></td> <td>47</td>	UNOsphere SUPrA TM	Bio-Rad	2008	29.0	150	9.3	10	1 mg/mL Polyclonal human		47
UNOSphere SUPrATMBio-Rad200824.040022.910Ing/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200832.01504.8201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200831.03009.5201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200831.03009.5201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200827.040012.0201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad201010.060042.952 mg/mL Polyclonal human47CaptivATM PriMABRepligen201017.030029.152 mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030029.152 mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030029.152 mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.038.010013.652 mg/mL Polyclonal human48	UNOsphere SUPrA TM	Bio-Rad	2008	27.0	300	18.0	10	1 mg/mL Polyclonal human		47
UNOsphere SUPrATMBio-Rad200832.01504.8201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200831.03009.5201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200827.040012.0201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200827.040012.0201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad201010.060042.952mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030029.152mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030029.152mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030029.152mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030013.652mg/mL Polyclonal human48	UNOsphere SUPrA TM	Bio-Rad	2008	24.0	400	22.9	10	یں 1 mg/mL Polyclonal human درج		47
UNOsphere SUPrATMBio-Rad200831.03009.5201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200827.040012.0201 mg/mL Polyclonal human47UNOsphere SUPrATMBio-Rad200827.040012.0201 mg/mL Polyclonal human47CaptivATM PriMABRepligen201010.060042.952 mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030029.152 mg/mL Polyclonal human48CaptivATM PriMABRepligen201017.030029.152 mg/mL Polyclonal human48CaptivATM PriMABRepligen201010.010.013.652 mg/mL Polyclonal human48	UNOsphere SUPrA TM	Bio-Rad	2008	32.0	150	4.8	20	یں 1 mg/mL Polyclonal human ترقیر		47
UNOsphere SUPrA TM Bio-Rad200827.040012.0201 m/mL Polyclonal human47CaptivA TM PriMABRepligen201010.060042.952 m/mL Polyclonal human48CaptivA TM PriMABRepligen201017.030029.152 m/mL Polyclonal human48CaptivA TM PriMABRepligen201017.030029.152 m/mL Polyclonal human48CaptivA TM PriMABRepligen201038.010013.652 m/mL Polyclonal human48	UNOsphere SUPrA TM	Bio-Rad	2008	31.0	300	9.5	20	یں 1 mg/mL Polyclonal human تحر		47
CaptivA TM PriMAB Repligen 2010 10.0 600 42.9 5 2 mg/mL Polyclonal human Cleaning in place pH 2-11 48 CaptivA TM PriMAB Repligen 2010 17.0 300 29.1 5 2 mg/mL Polyclonal human 48 CaptivA TM PriMAB Repligen 2010 17.0 300 29.1 5 2 mg/mL Polyclonal human 48 CaptivA TM PriMAB Repligen 2010 17.0 38.0 100 13.6 5 2 mg/mL Polyclonal human	UNOsphere SUPrA TM	Bio-Rad	2008	27.0	400	12.0	20	ا mg/mL Polyclonal human درج		47
CaptivA TM PriMAB Repligen 2010 17.0 300 29.1 5 2 mg/mL Polyclonal human 48 CaptivA TM PriMAB Repligen 2010 38.0 100 13.6 5 2 mg/mL Polyclonal human 48 CaptivA TM PriMAB Repligen 2010 38.0 100 13.6 5 2 mg/mL Polyclonal human 48	CaptivA TM PriMAB	Repligen	2010	10.0	600	42.9	5	یں 2 mg/mL Polyclonal human 127 نے میں د	Cleaning in place pH 2-11	48
CaptivA TM PriMAB Repligen 2010 38.0 100 13.6 5 2 mg/mL Polyclonal human 48	CaptivA TM PriMAB	Repligen	2010	17.0	300	29.1	S	2 mg/mL Polyclonal human		48
	CaptivA TM PriMAB	Repligen	2010	38.0	100	13.6	5	العن المراجعة المراجعة المراجعة المراجعة 1922 arg/mL 12 mbcs المراجعة المراجعة المراجعة المراجعة 1922 arg/mL		48

TABLE 1. Continued									
Protein A Resin	Manufacturer	Launch Year	Capacity (g/L) at 10% BT	Flow Rate (cm/h)	Productivity (g/L/h)	Bed Height (cm)	Feed Information	Caustic Stability	Reference
CaptivA TM PriMAB	Repligen	2010	45.0	50	7.1	5	2 mg/mL Polyclonal human IoG in PRS		48
Toyopearl [®] AF-rProtein A-650F	Tosoh	2010	30.5	100	12.6	5	10 mg/mL Human polyclonal IgG in PBS		28
Toyopearl [®] AF-rProtein A-650F	Tosoh	2010	24.0	200	22.9	Ś	10 mg/mL Human polyclonal IgG in PBS;	80% of initial binding capacity after 50 cycles with 17 min of 0.5 M NaOH	28
Monofinity A Resin TM	Genscript	2010	52.0	231	11.1	15.4	5 mg/mL hIgG	80% of initial binding capacity after 200 cycles with 0.5 M NaOH	49
MabSelect SuRe TM LX	GEHC	2011	18.3	009	30.2	10	Human polyclonal IgG		43
MabSelect SuRe TM LX	GEHC	2011	60.5	100	10.8 7.7	10	Human polyclonal IgG Human polyclonal IgG		64 64 64
MabSelect SuRe ^{1M} LX	GEHC	2011	67.7	75	5.9	10	Human polyclonal IgG	-	43
AbSolute [®] High Cap	Novasep	2011	44.0	200	28.4	S	1.5 mg/mL IgG	150 cycles with 20 min of 0.05 M NaOH	50
AbSolute [®] High Cap	Novasep	2011	42.0	300	42.0	5	1.5 mg/mL IgG	Working pH 1.5–9.0	50
AbSolute [®] High Cap	Novasep	2011	32.0	009	76.8 20.5	S S	1.5 mg/mL IgG	Working pH 1.5–9.0	50
Amsphere Protein A	NCL	7107	20.02	001	C. <i>KC</i>	10	at pH 7.3 at pH 7.3		10
Amsphere TM Protein A	JSR	2012	30.0	600	37.5	10	3.8 mg/mL IgG1 from CHO at nH 7.3		51
Amsphere TM Protein A	JSR	2012	40.0	188	12.9	10	3.8 mg/mL IgG1 from CHO at pH 7.3		51
KANEKA KanCapA TM	Kaneka	2012	24.0	667	19.0	20	Not provided	100 cycles with 15 min of 0.5 M NaOH	29
KANEKA KanCapA TM	Kaneka	2012	42.0	400	14.0	20	Not provided	100 cycles with 15 min of 0.5 M NaOH	29
KANEKA KanCapA TM	Kaneka	2012	51.0	240	8.9	20	Not provided	100 cycles with 15 min of 0.5 M NaOH	29
Eshmuno [®] A	Millipore	2013	45.0	400	14.3	20	mAb	Cleaning stability pH 1.5-13.5	30
Toyopreal [®] AF-rProtein A HC-650F	Tosoh	2013	102.3	120	51.0	5	10 mg/mL mAb		52
Toyopreal [®] AF-rProtein A HC-650F	Tosoh	2013	124.2	60	26.2	5	10 mg/mL mAb		52
Toyopreal [®] AF-rProtein A HC-650F	Tosoh	2013	143.3	24	10.7	2	10 mg/mL mAb		52
Toyopreal [®] AF-rProtein A HC-650F	Tosoh	2013	58.7	400	15.7	19.5	3 mg/mL mAb	15 min with 0.1 M NaOH	53
Toyopreal [®] AF-rProtein A HC-650F	Tosoh	2013	52.2	400	35.4	8.4	3 mg/mL mAb	15 min with 0.1 M NaOH	53
Toyopreal [®] AF-rProtein A HC-650F	Tosoh	2013	36.2	400	76.3	3.5	3 mg/mL mAb	15 min with 0.1 M NaOH	53
Provance TM Protein A	Grace	2014	28.0	600	36.5	10	Not provided		54
CIM [®] n or r-Protein	Grace BIA Separations	2014 2014	10.0 10.0	357 357	20.4 107.1	1.2	Not provided 1.0 mg/mL human IgG	Cleaning in place pH 1-13	55 55
A-1 TUBE MONOLITHIC COLUMN (Monolith)							20 mM Tris-HCl buffer, pH 7.4		

1197

Page 5 of 10

IADLE I. COMMAN									
		Launch	Capacity (g/L)	Flow	Productivity	Bed			
Protein A Resin	Manufacturer	Year	at 10% BT	Rate (cm/h)	(g/L/h)	Height (cm)	Feed Information	Caustic Stability	Reference
CIM [®] r-Protein A-1	BIA Separations	2014	4.2	8	5.0	0.3	Human polyclonal IgG		56
TUBE MONOI ITUIC									
COLUMN									
(Monolith)									
CIM [®] r-Protein A-1	BIA Separations	2014	3.7	30	17.2	0.3	Human polyclonal IgG		56
TUBE									
MONOLITHIC									
COLUMN (Monolith)									
CIM [®] r-Protein A-1	BIA Separations	2014	3.2	80	39.9	0.3	Human polyclonal IgG		56
TUBE									
MONOLITHIC									
COLUMN (Monolith)									
CIM [®] r-Protein A-1	BIA Separations	2014	2.7	160	70.4	0.3	Human polyclonal IgG		56
TUBE									
MONOLITHIC									
Mab Salaat SuDaTM and	Unation	2015	60.0	750	01	10.0	Unmon notice Long	Chort form all 2 12 7	57
Managered aure pec	OEAC	C107	00.0	007	19.2	10.0	пишап рогустопат таст	NUOLI-LEITII PIT 2-13./	10
*Productivity is defined a	s the grams of mAb the	ut can be pur	ified per hour per	liter of protein	A resin include	l loading, wash	, elution, and regeneration. Pro	oductivity was calculated with Eq.	2 using the resin
capacity and flow rate data as	ssuming a mAb titer of	1 g/L and 10	. 3. and 5 column	volumes of was	sh. elution. and r	egeneration. res	pectively.		



Figure 1. Protein A resin productivity as a function of product launch date.

Productivity was determined by the amount of grams of antibody that could be purified on a liter of protein A resin within the time required for column loading (assuming a 1 g/L titer), washing (10 CV), elution (3 CV), and regeneration (5 CV). Data are connected by solid (solid and dotted when there are two studies in the same year) lines for studies providing capacities at multiple flow rates. A compounded annual productivity growth rate of $4.3 \pm 1.4\%$ and an initial value Y_0 of 5.8 ± 2.4 g/L/h was determined by fitting all data except the membrane adsorber and monolith data to Eq. 3. Resins are depicted with squares and membrane adsorbers and monoliths are depicted by crosses.

Protein A capacity growth

The improvement in protein A capacity vs. year is plotted in Figure 2. A compounded annual capacity growth rate of $5.5 \pm 1.2\%$ and an initial value Y_0 of 7.2 ± 2.6 g/L was determined by fitting all data points except the membrane adsorber and monolith data to Eq. 3.

Operational flow rate growth

The improvement in operational flow rate vs. year is plotted in Figure 3. In contrast to protein A capacity and productivity, operational flow rate increased between 1978 and 2001, as new rigid and highly porous beads were introduced,²⁴ and then remained constant or declined.

Protein A sales growth

There is limited data describing the growth in protein A production and sales in the literature or in the annual reports of the major protein A resin producers: GE Healthcare, Millipore, and Applied Biosystems. However, Repligen Incorporated, the largest supplier of protein A ligand to the resin manufacturers, provided the percent of revenue from sales to GE healthcare, the largest protein A producer, and Applied Biosystems in annual reports. In addition, Repligen provided data for total protein A ligand sales from 2003 to 2011. These sales included protein A ligands used for production resins and small-scale research and diagnostic purposes. Sales to GE Healthcare grew from \$310,800 in 1999 to \$24,148,373 in 2014. A compounded annual sales growth rate of $21 \pm 2\%$ and an initial value Y_0 of





Data are connected by solid (solid and dotted when there are two studies in the same year) lines for studies providing capacities at multiple flow rates. A compounded annual capacity growth rate of $5.5 \pm 1.2\%$ and an initial value Y_0 of 7.2 ± 2.6 g/L was determined by fitting all data except the membrane adsorber and monolith data to Eq. 3. Resins are depicted with squares and membrane adsorbers and monoliths are depicted by crosses.



Figure 3. Protein A operational flow rate (the flow rate used for the reported capacity data) as a function of protein A launch date.

Data are connected by solid (solid and dotted when there are two studies in the same year) lines for studies providing capacities at multiple flow rates. Resins are depicted with squares and membrane adsorbers and monoliths are depicted by crosses.

 $$1,300,000 \pm 390,000$ was determined by fitting the GE Healthcare sales data to Eq. 3 and is plotted in Figure 4. This was similar to the over 20% ligand sales growth rate reported by Repligen in their 2015 annual report.

Repligen's annual reports indicated that GE Healthcare, their biggest protein A ligand customer by a large margin, primarily purchased protein A ligand for resin production.



Figure 4. The growth of Repligen protein A ligand sales.





Figure 5. The growth of protein A productivity, protein A capacity, mAb titer, protein A sales, and mAb sales vs. year.

mAb sales were determined from the sum of the sales of the top five mAbs (Avastin[®], Humira[®], Rituxan[®], Remicade[®], Herceptin[®]) and Enbrel[®] in each year. The growth curves that best fit the protein A productivity and capacity data in Figures 1 and 2, respectively, are plotted here. The sales of protein A ligand from Repligen to GE Healthcare in Figure 4 are plotted here.

Based on this, it is estimated here that total sales of protein A resin grew at the same rate as sales of Repligen protein A ligand to GE Healthcare: 21% between 1999 and 2014.

The growth of protein A productivity, protein A capacity, bioreactor mAb titer,^{59,60} protein A sales and mAb sales were compared vs. year in Figure 5. The growth curves that best fit the protein A productivity and capacity data in Figures 1 and 2, respectively, are plotted in Figure 5. The sales of the top five mAbs (Avastin[®], Humira[®], Rituxan[®], Remicade[®], Herceptin[®]) and Enbrel^{®61} in each year were obtained from annual reports and then added and plotted. The sales of protein A ligand from Repligen to GE Healthcare in Figure 4 are plotted in Figure 5.

Discussion

Growth in productivity, capacity, and operational flow rate

Protein A productivity (Figure 1) and capacity (Figure 2) have increased steadily since 1978 and have improved by an order of magnitude. It is believed that protein A productivity and capacity are still improving. One indicator of this is the fact that two of the highest performing resins, MabSelect SuReTM LX and Toyopreal[®] AF-rProtein A HC-650F, have been launched recently (2011 and 2013, respectively).

Surprisingly, the operational flow rate increased from 1978 to 2001 and then remained constant or declined (Figure 3). Operation at or above 1000 cm/h was first reported in 1993 and last reported in the Table 1 data in 2008. It is possible that this is because operating above 1000 cm/h provides a marginal improvement in the total time required for a protein A step. For example, at high flow rate it takes 70 min to do one protein A cycle (including the load, wash, elution, and regeneration steps), assuming a 20 cm bed height, 1000 cm/h flow rate, 40 g/L capacity, and 1 g/L titer. Given that the harvest and low pH viral inactivation steps that occur before and after the protein A step typically take more than an hour, and the antibody titers are now typically above 1 g/L, there is limited incentive to reduce the protein A cycle time below 70 min. The protein A flow rate can also be limited by pumping capacity.⁴

The highest productivity for a protein A resin always occurred at the highest flow rate in cases where data for multiple flow rates are reported. For example, the capacity of one resin was reported to be 10 g/L at 1500 cm/h and 20 g/L at 600 cm/h.³⁹ The higher flow rate condition is more productive on a g/L/h basis, but it is unlikely a capacity of 10 g/L would be acceptable in a manufacturing process.

It is interesting in Figure 2 to speculate on the theoretical limits of protein A capacity. Cation-exchange chromatography capacities approaching 200 g/L have been reported.^{18,62} It is far easier to produce a cation-exchange resin with a high density of properly orientated ligands than a protein A resin. However, it is likely that the capacity of protein A resins will continue to increase. It is likely that the size of the protein A ligand will be further reduced by eliminating all amino acids not required for structure, expression, or binding. This would allow for an increase in ligand density on the beads and an increase in space available for mAb binding. More porous but structurally rigid beads will also increase the space available for ligand immobilization and mAb binding. The number of protein A domains per ligand can be altered to optimize resin capacity. Optimal ligand immobilization, using more selective reactions and spacer arms, can improve capacity.

Protein A monoliths and membrane adsorbers can be operated at far higher flow rates and shorter bed heights than Page 8 of 10 packed columns and can therefore provide very high productivity.^{63,64} They have a number of disadvantages however that have prevented their use in manufacturing. They are currently only available up to the 8 mL scale and have low capacities (2.7-10 g/L in this study) and therefore dilute elution peaks.⁶⁵ For example, if three column volumes are used for elution, the elution pool concentration would range between 1 and 3 g/L. Breakthrough of mAb has been reported during loading of protein A membrane adsorbers.⁶⁶ Despite these disadvantages, they are included in this evaluation because of their potential. They may represent a format for future improvements in protein A productivity. The residence times, and therefore also the cycle times of protein A membrane adsorbers and monoliths are dramatically lower than those of columns, which would be beneficial for multicolumn chromatography operations.⁶⁷

Growth of protein A performance in the context of the mAb industry

It is interesting to consider the reasons for protein A productivity growth and to compare it to other industries. The technology improvement in 62 different technology industries has been described to be associated mainly with cumulative production growth in that industry.⁶⁸ Since protein A resin is mainly used for the production of mAbs, the growth of protein A production will be associated with the growth of the mAb industry. The average compound annual revenue growth rate for the top five mAb products and Enbrel over the 2004–2013 period was 20%.⁶¹ The growth rate of protein A sales (estimated here to be 21%) is higher than that reported for other advanced chromatography resins (5.8%⁶⁹), but similar to the sales growth rate of the mAb industry (20%), which is not surprising since protein A resin is used for mAb production.

Complicating the comparison of antibody and protein A resin sales is the fact that mAb revenues have increased in part due to higher drug prices, which would not be associated with increased protein A use. This is possibly compensated for by the growing use of protein A resin to make clinical phase antibodies which do not generate mAb revenue.

The mAb titer in bioreactors has increased from about 500 mg/L in 1987 to 5 g/L in 2006 in one study,⁶⁰ an annual growth rate of 14%. In another study,⁵⁹ the titer increased from 0.2 g/L in 1985 to 2.64 g/L in 2015, also an annual growth rate of 14%, which is higher than the rate of growth of protein A productivity (4.3%). This difference is not surprising because limitations in titer and bioreactor capacity were a significant threat to the early growth of the therapeutic mAb industry.^{70,71} Limitations in protein A productivity never threatened mAb industry growth and required large development investments. In addition, since roughly 2000, revenue from protein A sales have been far lower than revenue from mAb sales (Figure 5). Because of this, the financial resources available to protein A vendors for product improvements are far below the resources available to mAb producers to improve bioreactor titers.

It is notable in Figure 5 that protein A productivity and mAb titer both increased rapidly starting in about 1990, with the mAb titer increasing faster than protein A productivity. This coincides with the beginning of early clinical development of the highest selling mAbs. The sales of protein A and therapeutic mAbs started to increase rapidly about ten

years later in 2000. This gap roughly coincides with the time required to perform all clinical testing, obtain regulatory approval, and develop the market for a mAb. The fact that growth in bioreactor titer and growth in protein A performance happened before growth in sales indicates that investments in development were occurring at risk assuming that future sales would be strong.

Conclusions

While the sequence of steps used for mAb purification has not changed since the first therapeutic antibody was licensed, the performance of protein A chromatography, as indicated by capacity and productivity has steadily improved. Protein A productivity and capacity have increased 4.3 and 5.5% a year, respectively, since 1978 and are likely to continue to increase. The improvements have been caused in part by improvements in bead rigidity, size, polymers, and pore structure. In addition, changes in ligand immobilization, the number of protein A domains, and the protein A amino acid sequence have led to improvements.

In contrast, protein A operational flow rate increased between 1978 and 2001 and then remained constant or declined as further improvements provided only marginal benefits. The productivity of protein A resin rapidly improved starting in about 1990, in parallel with improvements in bioreactor titer, to economically provide material for clinical trials. The sales of protein A resin, as indicated by sales of the protein A ligand (21% growth), have closely paralleled the sales of mAbs (20% growth). Both increased rapidly in 2000 after the first major mAbs were approved and the markets were developed. It is likely that the many alternatives to protein A chromatography that have been explored have not been implemented because of the steady improvement in protein A performance.35 Protein A membrane adsorbers and monoliths have higher productivity than packed columns due to their short bed heights and high operational flow rates. These devices are not currently practical for large-scale manufacturing, but may represent a format for future improvements in protein A productivity.

Acknowledgments

The authors acknowledge the help and assistance of Ganesh Vedantham, Paul O'Neill, John Royce, and Alejandro Becerra-Arteaga for their contributions to this work.

Literature Cited

- 1. Shukla AA, Hubbard B, Tressel T, Guhan S, Low D. Downstream processing of monoclonal antibodies-application of platform approaches. J Chromatogr B. 2007;848:28-39.
- 2. Zhang Q, Goetze AM, Cui H, Wylie J, Trimble S, Hewig A, Flynn GC. Characterization of the co-elution of host cell proteins with monoclonal antibodies during protein A purification. Biotechnol Prog. 2016; 32:708-717.
- 3. Purdie JL, Kowle RL, Langland AL, Patel CN, Ouyang A, Olson DJ. Cell culture media impact on drug product solution stability. Biotechnol Prog. 2016; Apr 25. doi: 10.1002/ btpr.2289. [Epub ahead of print]
- 4. Zheng J, Wang L, Twarowska B, Laino S, Sparks C, Smith T, Russell R, Wang M. Caprylic acid-induced impurity precipitation from protein A capture column elution pool to enable a two-chromatography-step process for monoclonal antibody purification. Biotechnol Prog. 2015;31:1515-1525.

- 5. Jensen K. A normally occurring Staphylococcus antibody in human serum. Acta Pathol Microbiol Scand. 1958;44:421-428.
- 6. Jensen K, Neter E, Gorzynski EA, Anzai H. Studies on toxic products of Staphylococcus. Acta Pathol Microbiol Scand. 1961;53:191-200.
- 7. Löfkvist T, Sjöquist J. Chemical and serological analysis of antigen preparations from Staphylococcus aureus. Acta Pathol Microbiol Scand. 1962;56:295-304.
- 8. Grov A, Myklestad B, Oeding P. Immunochemical studies on antigen preparations from Staphylococcus aureus. Acta Pathol Microbiol Scand. 1964;61:588-596.
- 9. Forsgren A, Sjöquist J. "Protein A" from S. aureus: I. Pseudoimmune reaction with human y-globulin. J Immunol. 1966;97: 822-827
- 10. Hjelm H, Hjelm K, Sjöquist J. Protein a from Staphylococcus aureus. Its isolation by affinity chromatography and its use as an immunosorbent for isolation of immunoglobulins. FEBS Lett. 1972:28:73-76.
- 11. Sjoquist JA. Method of Binding Immunoglobulin Employing a Polypeptide from Microorganisms. United States Patent 3,995,018; 1976.
- 12. Ey PL, Prowse SJ, Jenkin CR. Isolation of pure IgG1, IgG2a and IgG2b immunoglobulins from mouse serum using protein A-sepharose. Immunochemistry. 1978;15:429-436. Jul
- 13. Roque ACA, Silva CSO, Taipa MA. Affinity-based methodologies and ligands for antibody purification: advances and perspectives. J Chromatogr A. 2007;1160:44-55.
- 14. Lain B. Protein A: the life of a disruptive technology. BioProcess Int. 2013;11:29-38.
- 15. Miesegaes GR, Lute S, Strauss DM, Read EK, Venkiteshwaran A, Kreuzman A, Shah R, Shamlou P, Chen D, Brorson K. Monoclonal antibody capture and viral clearance by cation exchange chromatography. Biotechnol Bioeng. 2012;109:2048-2058.
- 16. Lain B, Cacciuttolo MA, Zarbis-Papastoitsis G. Development of a high capacity MAb capture step based on cation-exchange chromatography. BioProcess Int. 2009;7:26-34.
- 17. Follman DK, Fahrner RL. Factorial screening of antibody purification processes using three chromatography steps without protein A. J Chromatogr A. 2004;1024:79-85.
- 18. Tao Y, Ibraheem A, Conley L, Cecchini D, Ghose S. Evaluation of high capacity cation exchange chromatography for direct capture of monoclonal antibodies from high-titer cell culture processes. Biotechnol Bioeng. 2014;111:1354-1364.
- 19. Kuczewski M, Schirmer E, Lain B, Zarbis-Papastoitsis G. A singleuse purification process for the production of a monoclonal antibody produced in a PER.C6 human cell line. *Biotechnol J.* 2011;6:56–65.
- 20. Eggersgluess JK, Richter M, Dieterle M, Strube J. Multi-stage aqueous two-phase extraction for the purification of monoclonal antibodies. Chem Eng Technol. 2014;37:675-682.
- 21. Trilisky E, Gillespie R, Osslund TD, Vunnum S. Crystallization and liquid-liquid phase separation of monoclonal antibodies and fc-fusion proteins: screening results. Biotechnol Prog. 2011;27:1054-1067.
- 22. Detmers F, Hermans P, Jiao J-A, McCue JT. Novel affinity ligands provide for highly selective primary capture. BioProcess Int. 2010; February: 50-54
- 23. Newcombe AR, Cresswell C, Davies S, Watson K, Harris G, O'Donovan K, Francis R. Optimised affinity purification of polyclonal antibodies from hyper immunised ovine serum using a synthetic Protein A adsorbent, MAbsorbent® A2P. J Chromatogr B. 2005;814:209-215. 1/25/
- 24. Fahrner RL, Whitney DH, Vanderlaan M, Blank GS. Performance comparison of protein A affinity-chromatography sorbents for purifying recombinant monoclonal antibodies. Biotechnol Appl Biochem. 1999;30:121-128.
- 25. Fuglistaller P. Comparison of immunoglobulin binding capacities and ligand leakage using eight different protein A affinity chromatography matrices. J Immunol Methods. 1989;124:171-177.
- 26. Shandle P, Mills G, Erickson J, Scott R, Smith T. Antibody purification. United States Patent 5,429,746; 1995.
- 27. Pall Life Sciences. BIOSEPRA® Protein A Ceramic HYPERD® F. Product note-LPN PN702-004-12/2004.
 28. Tosoh Bioscience. TOYOPEARL[®] AF-rProtein A-650F. *F10P11B*.
- 29. Iritani K, Kitahashi H, Sato T, Bando K. KANEKA KanCapA(TM). A New mAb Purification Platform. BioProcess. International Industry Yearbook 2013-2014, 64-66.

- Merck Millipore. Eshmuno[®] A Chromatography Media. Lit. No. DS5553EN00 PS-13-08674 Rev.B. 2013.
- Amerhsam Biosciences. Affinity chromatography—rmp Protein A Sepharose Fast Flow. Data file 18-1141-34 AB, 2000-04.
- 32. GE Healthcare. MabSelect SuRe. Data file 11-0011-65 AC.
- Ghose S, Allen M, Hubbard B, Brooks C, Cramer SM. Antibody variable region interactions with Protein A: implications for the development of generic purification processes. *Biotechnol Bio*eng. 2005;92:665–673.
- Müller E, Vajda J. Routes to improve binding capacities of affinity resins demonstrated for Protein A chromatography. *J Chromatogr B*. 2016;1021:159–168.
- 35. Kelley B. Very large scale monoclonal antibody purification: the case for conventional unit operations. *Biotechnol Prog.* 2007;23:995–1008.
- Hale G, Drumm A, Harrison P, Phillips J. Repeated cleaning of protein A affinity column with sodium hydroxide. *J Immunol Methods*. 1994;171:15–21. May 2
- GE Healthcare. nProtein A Sepharose 4 Fast Flow. Data File 18-1125-19 AD.
- GE Healthcare. Antibody affinity chromatography—rProtein A SepharoseTM Fast Flow. Instructions 71-5000-09 AD.
- 39. Keener RN, Fernandez EJ, Maneval JE, Hart RA. Advancement in the modeling of pressure-flow for the guidance of development and scale-up of commercial-scale biopharmaceutical chromatography. J Chromatogr A. 2008;1190:127–140.
- 40. Yang L, Harding JD, Ivanov AV, Ramasubramanyan N, Dong DD. Effect of cleaning agents and additives on Protein A ligand degradation and chromatography performance. *J Chromatogr A*. 2015;1385:63–68.
- EMD Millipore. ProSep[®]—vA High Capacity Chromatography Media—Designed for large-scale purification of therapeutic antibodies. Data Sheet DS1013EN00RevB.
- 42. Wang C, Mann F. Increasing MAb capture productivity: use of transport and process models in development and operation of improved chromatography media. *BioProcess Int.* 2009; 56–61.
- GE Healthcare. Dynamic binding capacity study on MabSelect SuRe LX for capturing high-titer monoclonal antibodies. Application Note 28-9875.
- 44. Sartorius. Sartobind[®] Protein A 2 ml—Laboratory Scale Affinity Membrane Adsorbers Publication No.: SL-2050-e08031.
- 45. Gebski C, Surka S, Leete T, Xu M, McCoy M, Lynch P, Hunt T, Purtell C. Characterization of Poros Mabcapture A chromatography media. *Bioprocess Online* 2011; October:1.
- Agilent Technologies. Bio-Monolith Protein A Column. September 16, 2008;5973–1722.
- Bio-Rad Laboratories. UNOsphere SUPrATM Affinity Chromatography Media. 5729 RevB.
- Repligen. Captiva Primab—Protein A affinity resin—Regulatory Support File RSF-1003-01. March 2010.
- Genscript. Monofinity A Resin for Antibody Purification PBM1407212010.
- Novasep. AbSoluteTM by Novasep: High Performance Protein A Media For the capture of monoclonal and polyclonal antibodies 10-2009.
- McCaw TR, Koepf EK, Conley L. Evaluation of a novel methacrylate-based Protein A resin for the purification of immunoglobulins and Fc-fusion proteins. *Biotechnol Prog.* 2014;30: 1125–1136.
- Regina R. Protein A chromatography with High-Titer Feedstocks. *BioProcess International Industry Yearbook* 2014–2015, 60.

- 53. Tosoh Bioscience. Comparison of TOYOPEARL[®] AF-rProtein A HC-650F Binding Capacity at Various Bed Heights and Constant Linear Velocity. Application note AN78 0115.
- 54. Preisig C. Innovations in purification: lowering cost and accelerating drug development with pre-packed columns for single use. *Bioprocess Online*. 2015; January:1–5
 55. BIA Separations GmbH. CIM[®] Tube Monolithic Columns for
- BIA Separations GmbH. CIM[®] Tube Monolithic Columns for Antibody Affinity Chromatography. 2015;Publication #: PSISnArArGrL-1-1503.
- Herigstad MO, Dimartino S, Boi C, Sarti GC. Experimental characterization of the transport phenomena, adsorption, and elution in a protein A affinity monolithic medium. *J Chromatogr A*. 2015; 1407:130–138.
- 57. GE Healthcare. Affinity chromatography GE Healthcare MabSelect SuRe[™] pcc Data file, 29177558 AA2015.
- 58. Chakrabarti A, Muller E, Vajda J, Wacker A. Characterization of a newly developed high capacity, alkaline resistant, recombinant protein A resin. *Tosoh Biosciences TP198*. American Chemical Society BIOT Conference, Dallas, Texas 2014.
- 59. Rader RA, Lander ES. Biopharmaceutical manufacturing: historical and future trends in titers, yields, and efficiency in commercial-scale bioprocessing. *BioProcessing J.* 2014/2015;13:1–9.
- Seth G, Hossler P, Yee J, Hu W-S. Engineering cells for cell culture bioprocessing—physiological fundamentals. In: Hu W-S, editor. Cell Culture Engineering. Vol 101, Berlin: Springer; 2006:119–164.
- 61. Ecker DM, Jones SD, Levine HL. The therapeutic monoclonal antibody market. *mAbs.* 2015;7:9.
- 62. Gagnon P. How to choose an industrial cation exchanger for IgG purification. *BioProcess Int.* 2010:22–34.
- Mayolo-Deloisa K, González-Valdez J, Rito-Palomares M. PEGylated protein separation using different hydrophobic interaction supports: conventional and monolithic supports. *Biotechnol Prog.* 2016;32:702–707.
- 64. Hou Y, Brower M, Pollard D, Kanani D, Jacquemart R, Kachuik B, Stout J. Advective hydrogel membrane chromatography for monoclonal antibody purification in bioprocessing. *Biotechnol Prog.* 2015;4:974–982.
- 65. Brorson K, Brown J, Hamilton E, Stein KE. Identification of protein A media performance attributes that can be monitored as surrogates for retrovirus clearance during extended re-use. *J Chromatogr A*. 2003;989:155–163.
- Morrow JK. Improving protein production processes upstream manufacture increases, while downstream operations tend to falter. *Genet Eng Biotechnol News* 2007;27:1–5.
- 67. Kaltenbrunner O, Diaz L, Hu X, Shearer M. Continuous bindand-elute protein A capture chromatography: optimization under process scale column constraints and comparison to batch operation. *Biotechnol. Prog.* 2016 Apr 25. doi: 10.1002/btpr.2291. [Epub ahead of print]
- Nagy B, Farmer JD, Bui QM, Trancik JE. Statistical basis for predicting technological progress. *PloS One* 2013;8:e52669.
- 69. www.beroe-inc.com. Implementation of advanced chromatography techniques to mitigate purification concerns in bispecific monoclonal antibody manufacturing. 2012:1–16.
- Mallik A, Pinkus GS, Sheffer S. Biopharma's capacity crunch. The McKinsey Quarterly 2002; 2.
- Butler M. Animal cell cultures: recent achievements and perspectives in the production of biopharmaceuticals. *Appl Microbiol Biotechnol.* 2005;68:283–291.

Manuscript received May 24, 2016, and revision received July 3, 2016.

1202