




Fig. 1.—The log of α as a function of $10^3/T$.

Results and Discussion

The results of these experiments are summarized in Table I. The uncertainty in the isotopic equilibrium constants is estimated to be less than one digit in the last decimal place shown. As expected, the data show an inverse dependence upon temperature.

By plotting $\log \alpha$ vs. $1/T$, a straight line was obtained, as shown in Fig. 1. The heat of exchange calculated from the slope of this line was 2.26 cal./mole and $-\Delta S$ was 1.81×10^{-3} cal./mole degree at 25° . These results are slightly higher than those given by Glueckauf³ for the $\text{Na}^{22}\text{--Na}^{24}$ exchange, and doubtless reflect the greater mass difference ratio ($\Delta M/M$) which exists between the isotopes of lithium as compared to Na^{22} and Na^{24} .

USE OF GLASS ELECTRODES TO MEASURE ACIDITIES IN DEUTERIUM OXIDE^{1,2}

BY PAUL K. GLASOE³ AND F. A. LONG

Department of Chemistry, Cornell University, Ithaca, New York

Received August 26, 1959

Recent studies have shown that a conventional glass electrode apparatus can be used to give fairly precise values of acidity in aqueous solutions, particularly when only relative values of the acidity are needed.^{4,5} One situation where it would be helpful to have additional data on relative acidities is for solutions of weak acids and bases in the solvents D_2O and H_2O . We have therefore investigated the applicability of the glass electrode to measurement of acidity in D_2O solutions. A general study is already available on some of the properties of glass electrodes in D_2O ,⁶ but it does not give much con-

sideration to the measurement of acidity in D_2O with ordinary electrodes. We have recently learned that this latter problem was considered some time ago by Hart, and by R. B. Fischer and R. A. Potter but the results are available only in special publications.^{7,8}

Experimental

All measurements were made with a Beckman Model G pH meter at 25° . The Beckman No. 39166 (old No. 19166) glass electrode and saturated calomel electrode combination was used for all measurements, unless otherwise noted. The pH meter was standardized with conventional buffer mixtures chosen to be close to the range of the pH measurements. Reagent grade chemicals were used without further purification. Formic acid was redistilled and maleic acid, aniline hydrochloride and pyridinium perchlorate were recrystallized. The D_2O was 99.5 atom % deuterium. Solutions of NaOD were made by passing water vapor over sodium. Other solutions in D_2O were made by dissolving the anhydrous hydrogen compound (except for hydrochloric acid where a concentrated aqueous solution was diluted). For the low concentrations studied, this did not change the deuterium content by a significant amount. All concentrations are in moles (or equivalents) per liter.

Results and Discussion

Glass Electrode Measurements in Solutions of Strong Acids.—Solutions of roughly 0.01 M hydrochloric acid in H_2O and in D_2O were prepared by dilution and the exact concentration of acid was determined by titration with standard base. The averages of several pH meter readings with the standard electrode combination are given under (a) of Table I. For solutions of comparable acidity

TABLE I

GLASS ELECTRODES IN SOLUTIONS OF HYDROCHLORIC ACID IN H_2O AND IN D_2O

Solutions are 0.00976 M HCl in H_2O and 0.00983 M DCl in D_2O ; pH meter standardized with pH 4.00 phthalate buffer

	—Av. rdgs.—		Dif-
	H_2O soln.	D_2O soln.	ference
(a) Standard glass electrode— Beckmann No. 39166; 39168 calomel	2.08	1.69	0.40
(b) Beckmann 1190-80 glass electrode; 1170 calomel	2.08	1.69	.39
(c) Beckmann type E glass electrode; 1170 calomel	2.02	1.62	.40
(d) "Synthetic" 1190-80 glass electrode; 1170 calomel	2.13	1.73	.40

the pH meter reading in D_2O solution is 0.40 pH unit lower than in H_2O solution. To determine whether this difference was characteristic only of this particular electrode system, these same solutions were measured using various glass electrode-calomel electrode combinations. The results are given in (b), (c) and (d) of Table I. It is apparent that the difference is constant for the different electrodes, within the error of reading the meter. The "synthetic 1190-80" electrode in (d), Table I, is an ordinary Beckman 1190-80 electrode from which the internal solution was poured out and replaced with a solution of 0.100 M HCl in H_2O . The

(1) Supported in part by a grant from the Atomic Energy Commission.

(2) Presented at the 136th meeting of the American Chemical Society, Atlantic City, September, 1959.

(3) National Science Foundation Science Faculty Fellow at Cornell University, 1958-1959.

(4) E. Grunwald, *J. Am. Chem. Soc.*, **73**, 4934 (1951).

(5) A. L. Bacarella, E. Grunwald, H. F. Marshall and E. L. Purlee, *THIS JOURNAL*, **62**, 856 (1958).

(6) D. Hubbard and G. W. Cleek, *J. Research Natl. Bur. Standards*, **49**, 267 (1952).

(7) R. G. Hart, *Nat. Res. Council, Canada*, CRE 423, June 1949.

(8) R. B. Fischer and R. A. Potter, A.E.C. document NODC-715, Sept. 12, 1945.

TABLE II
MEASUREMENTS WITH GLASS ELECTRODE FOR STRONG BASES IN H₂O AND IN D₂O

	Base	Acidity		Δ calcd.	pH Reading		Δ exp.	ΔpD (Δ calcd. - Δ exp.)
		$-\log C_{H^+}$	$-\log C_{D^+}$		In H ₂ O	In D ₂ O		
(a)	NaOH NaOD	12.00	12.86	0.86	11.83	12.21	0.38	0.48
(b)	NaOH NaOD	12.00	12.86	.86	12.00 ^a	12.47 ^a	.47	.39
(c)	Ba(OH) ₂ Ba(OD) ₂	12.46	13.29	.83	12.38	12.82	.44	.39
(d)	Ca(OH) ₂ Ca(OD) ₂	12.28	13.14	.86	12.22	12.67	.45	.41

^a Measurements made with type E (low sodium ion error) electrode.

actual potential of this electrode was markedly different from a normal 1190-80 electrode but even so the difference for the two solvents remains at 0.40 pH unit.⁹

These measurements indicate that the difference of 0.40 pH meter unit is not dependent on the particular type of glass electrode and that the pD of a solution in D₂O can be determined by use of the equation

$$pD = pH \text{ meter reading} + 0.40 \quad (1)$$

where the "pH meter reading" is obtained with an apparatus standardized to read pH in H₂O solutions. For convenience, we shall refer to the difference, pD minus pH meter reading, in this case 0.40, as ΔpD .¹⁰

Fisher and Potter⁷ made studies of a glass electrode system using buffer solutions in D₂O and concluded that an equation like 1 could be used to give pD . However, they proposed a numerical constant of 0.25 rather than 0.40. Hart⁸ did a similar study with solutions of hydrochloric acid in the two solvents and also with phosphate buffers (pH of from 5 to 7). He found values for the constant of equation 1 which range from 0.40 to 0.47. Hart also gave further consideration to the Fisher and Potter results and noted difficulties in the latter's work, both in the extrapolations and in the calculation of ionic strength. We conclude that the value of 0.40 for the constant of equation 1 agrees within experimental error with all of the earlier results.

Glass Electrode Measurements in Solutions of Strong Bases.—pH meter readings using the standard electrodes were made on solutions of 0.001009 *M* NaOH in H₂O and 0.001104 *M* NaOD in D₂O. The resulting average values are given under (a) in Table II. Calculation of the value of $-\log C_{D^+}$ or $-\log C_{H^+}$ for solutions of known concentration of hydroxide ion involves the ion product of water. This was taken to be 0.15×10^{-14} and 1.01×10^{-14} for D₂O and H₂O, respectively.¹¹

The value of ΔpD of 0.48 for the sodium hydroxide solutions is significantly higher than that obtained for acid solutions. However, it is well known that

(9) In view of the frequent recommendation that a glass electrode should be equilibrated in water for several hours before use, it is worth noting that when a glass electrode was transferred from a water solution to a D₂O solution (or vice versa) the "pH" difference of 0.40 appeared to be established as rapidly as measurements could be made.

(10) As is well known, the definition of pH is somewhat fuzzy. One can assume that the definition of pD is equally so. Our implicit definition of pD is that it "means the same thing for solutions in pure D₂O that pH does for H₂O."

(11) R. W. Kingerley and V. K. LaMer, *J. Am. Chem. Soc.*, **63**, 3256 (1941).

the sodium ion error of the standard glass electrode depends on the pH at a fixed concentration of sodium ion. From the Beckman chart of sodium ion error it is found that for 0.01 *M* sodium ion at 25° the correction is 0.13 pH unit at pH 12.80 and 0.05 unit at pH 12.00. The value of ΔpD taking this into account is $0.48 - 0.08 = 0.40$ pH unit, the same as the value for acid solutions. In agreement with this, a ΔpD value of 0.39 (Table IIb) was found for these same solutions with the Beckman type E electrode which has a negligible sodium ion error at these concentrations. Further confirmation of the similarity of ΔpD in acid and in base solutions is given by the results obtained with solutions of approximately 0.029 *N* barium hydroxide and 0.02 *N* calcium oxide in H₂O and D₂O (Table II, (c) and (d)). These solutions would be expected to show an insignificant cation effect at these low concentrations. In both cases the ΔpD is 0.40 pH unit, within experimental error.

K_H/K_D Ratios for Some Weak Acids and Bases.—To determine whether the correction term 0.40 is applicable in the pH range between the extremes of strong acids and strong bases, the pK_a values for some weak acids were determined in H₂O and in D₂O using the glass electrode to measure pH and pD . For every acid, values of pK_H' or pK_D' (where K' is the concentration equilibrium constant) were calculated from the pH or pD value and the molar ratio of acid and conjugate base for each of several buffer ratios. Values of $\Delta pK'$, defined as $pK_D' - pK_H'$ were always calculated from pK_D' and pK_H' values for solutions of equal ionic strength. In most measurements the ionic strength was less than 0.10. Values of pK_H or pK_D were calculated from pK' using the activity coefficient expression

$$-\log f_{\pm} = \frac{0.509\sqrt{I}}{1 + \sqrt{I}}$$

where I is ionic strength for molar concentrations. It made no difference, within experimental error, whether the K_H/K_D ratio was calculated from the average of the $\Delta pK'$ or the ΔpK values.

Table III summarizes the values of K_H/K_D obtained for a variety of weak acids. In general the agreement between the K_H/K_D ratios from glass electrode studies and from other methods of measurement is quite satisfactory, and it seems safe to conclude that the factor of 0.40 for ΔpD is applicable to measurements at intermediate acidities.




Fig. 1.—Dependence of pH meter reading for 0.01 M hydrochloric acid on atom per cent. deuterium of the aqueous solvent. Meter standardized to read pH for H₂O solutions.

TABLE III

K_D/K_H FROM MEASUREMENTS OF pH AND pD WITH GLASS ELECTRODE

Acid	pK^H	pK^D	ΔpK^a	K^H/K^D	K^H/K^D other
Acetic	4.73	5.25	0.52	3.3	3.33 ¹²
Formic	3.75	4.20	.45	2.8	2.5 ¹³ ; 2.7 ¹⁴
Aniline-HCl	4.55	5.13	.58	3.8	3.1 ¹³
Maleic, pK_1	1.98	2.53	.55	3.6	4.2 ¹⁵
Maleic, pK_2	6.28	6.61	.33	2.1	2.4 ¹⁵
Phosphoric, pK_1	2.11	2.31	.20	1.6	1.6 ¹³
Carbonic, pK_2	10.33	10.96	.63	4.3	4.4 ¹⁶
<i>m</i> -Nitroaniline-HCl	2.48	2.96	.48	3.0	3.8 ¹⁷

^a Values of $\Delta pK'$, calculated from measurements of similar ionic strengths are the same, within experimental error.

Dependence of ΔpD on Deuterium Content of Solution.—Varying quantities of 0.010 M HCl in H₂O and 0.010 M DCl in D₂O were mixed to form a series of solutions of the same acid concentration but with different deuterium content. The “pH’s” of these solutions were measured with the standard glass electrode. As Fig. 1 shows the reading of the pH meter is very nearly a linear function of the atom per cent. deuterium.

Source of Potential Difference Giving Rise to ΔpD .—The correction term ΔpD could arise from a change of potential in D₂O compared to H₂O for either or both the glass electrode and the calomel electrode. In an exploratory experiment on this point, after the pH meter was standardized with the standard electrodes, the calomel electrode was replaced by one containing a saturated solution of potassium chloride in D₂O. There was no significant change in the meter reading. It thus appears that the potential change is due solely to the glass electrode.

(12) S. Korman and V. K. LaMer, *J. Am. Chem. Soc.*, **58**, 1396 (1936).

(13) G. Schwarzenbach, *Z. Elektrochem.*, **44**, 46 (1938).

(14) J. C. Hornel and J. A. V. Butler, *J. Chem. Soc.*, 1361 (1936).

(15) G. Dahlgren and F. A. Long, *J. Am. Chem. Soc.*, **82**, in press (1960).

(16) J. Curry and Z. F. Hugus, *ibid.*, **66**, 653 (1944). This reference lists 3.95 but recalculation for molar concentrations leads to the value 4.4 (J. Curry, private communication).

(17) E. Hogfeldt and J. Bigeleisen, *J. Am. Chem. Soc.*, **82**, 15 (1960).

THE NATURE OF THE S-O BOND IN DISULFUR MONOXIDE


By P. A. GIGUÈRE

Department of Chemistry, Laval University, Québec, Canada

Received October 14, 1959

Recently Meschi and Myers¹ have reported the detailed microwave spectrum and molecular structure of disulfur monoxide, S₂O. Mass spectrometric and other measurements² had previously led them to positive identification of that unstable compound, which at one time³ was mistaken for dimeric sulfur monoxide, (SO)₂. From analysis of the spectra they found for the molecule a bent S-S-O structure with the following parameters: S-S distance, 1.884 Å.; S-O, 1.465 Å.; S-S-O angle, 118 ± 0.5°. The authors also concluded: “the S-O bond is probably fairly close to a single bond.” However, this view is not supported by the existing data. A single covalent S-O bond should be appreciably longer than the above, namely, 1.70 Å., from the sum of the covalent radii of the S and O atoms.⁴ (The revised values corrected for partial ionic character of the bond⁵ lead to nearly the same bond length, 1.69 Å.) The distance 1.51 Å. quoted by Meschi and Myers for the single S-O bond is in fact the interatomic distance in the tetrahedral SO₄²⁻ ion where considerable double bond character is expected.

As for the double-bond radii of the atoms, their “normal” values are much more uncertain than those of the single bonds. At any rate the S-O bonds in SO₂, 1.432 Å., are definitely shorter than the sum of the Pauling’s double-bond radii, 1.49 Å. The explanation for this has been given by Moffitt⁶ who analyzed the electronic structure of various sulfoxides and sulfones by means of his self-consistent LCAO method and showed that in these molecules the S-O bonds are largely double, with the 3d orbitals of sulfur contributing appreciably to the π -bonding. The resulting hybrid bond is stronger than one involving a pure p- or d-bonding orbital of sulfur. Obviously, such is the case for disulfur monoxide where the central S atom can use its 3d orbital in hybridization (form I) rather than forming so-called coordinate bonds with large formal charge separation as in the canonical forms II and III



From the known vibrational frequencies of the S₂O molecule⁷ the force constant for the S-O bond may be estimated roughly at 9.3×10^5 dynes cm.⁻¹

(1) D. J. Meschi and R. J. Myers, *J. Mol. Spectroscopy*, **3**, 405 (1959).

(2) D. J. Meschi and R. J. Myers, *J. Am. Chem. Soc.*, **78**, 6220 (1956).

(3) For a review see P. W. Schenk, *Chem. Z.*, **67**, 273 (1943).

(4) L. Pauling, “The Nature of the Chemical Bond,” Cornell University Press, Ithaca, N. Y., 1945.

(5) V. Schomaker and D. P. Stevenson, *J. Am. Chem. Soc.*, **63**, 37 (1941).

(6) W. Moffitt, *Proc. Roy. Soc. (London)*, **A200**, 409 (1950).

(7) A. V. Jones, *J. Chem. Phys.*, **18**, 1263 (1950).