

History of Changes for Study: NCT00729664

Phase 1 Open Label Study of MDX-1105 in Subjects With Solid Tumors

[Latest version \(submitted September 3, 2015\) on ClinicalTrials.gov](#)

- A study version is represented by a row in the table.
- Select two study versions to compare. One each from columns A and B.
- Choose either the "Merged" or "Side-by-Side" comparison format to specify how the two study versions are to be displayed. The Side-by-Side format only applies to the Protocol section of the study.
- Click "Compare" to do the comparison and show the differences.
- Select a version's Submitted Date link to see a rendering of the study for that version.
- The yellow A/B choices in the table indicate the study versions currently compared below. A yellow table row indicates the study version currently being viewed.
- Hover over the "Recruitment Status" to see how the study's recruitment status changed.
- Study edits or deletions are displayed in **red**.
- Study additions are displayed in **green**.

Study Record Versions

Version	A	B	Submitted Date	Changes
1	<input checked="" type="radio"/>	<input type="radio"/>	August 4, 2008	None (earliest Version on record)
2	<input type="radio"/>	<input checked="" type="radio"/>	April 17, 2009	Recruitment Status, Contacts/Locations, Study Status, Arms and Interventions, Outcome Measures, Study Description, Eligibility and Oversight
3	<input type="radio"/>	<input type="radio"/>	June 18, 2009	Study Status and Sponsor/Collaborators
4	<input type="radio"/>	<input type="radio"/>	June 24, 2009	Contacts/Locations, Arms and Interventions, Study Design, Study Identification, Study Description, Study Status, Eligibility, Outcome Measures and Conditions
5	<input type="radio"/>	<input type="radio"/>	July 22, 2009	Contacts/Locations, Study Status, Conditions and Study Identification
6	<input type="radio"/>	<input type="radio"/>	July 31, 2009	Contacts/Locations and Study Status
7	<input type="radio"/>	<input type="radio"/>	October 27, 2009	Contacts/Locations, Arms and Interventions, Study Status, Eligibility, Study Design, Conditions and Study Description
8	<input type="radio"/>	<input type="radio"/>	January 22, 2010	Study Status and Contacts/Locations
9	<input type="radio"/>	<input type="radio"/>	March 12, 2010	Sponsor/Collaborators, Contacts/Locations, Study Status and Study Identification
10	<input type="radio"/>	<input type="radio"/>	March 29, 2010	Study Status and Study Identification
11	<input type="radio"/>	<input type="radio"/>	April 22, 2010	Study Status
12	<input type="radio"/>	<input type="radio"/>	June 24, 2010	Study Status, Eligibility and Study Design
13	<input type="radio"/>	<input type="radio"/>	December 2, 2010	Study Status
14	<input type="radio"/>	<input type="radio"/>	January 19, 2011	Arms and Interventions, Study Status, Study Identification, Conditions, Eligibility and Study Design
15	<input type="radio"/>	<input type="radio"/>	May 24, 2011	Contacts/Locations, Study Status and Study Design
16	<input type="radio"/>	<input type="radio"/>	July 25, 2011	Study Status and Eligibility
17	<input type="radio"/>	<input type="radio"/>	January 11, 2012	Sponsor/Collaborators, Study Status and Contacts/Locations
18	<input type="radio"/>	<input type="radio"/>	January 16, 2012	Study Identification and Study Status
19	<input type="radio"/>	<input type="radio"/>	February 2, 2012	Contacts/Locations and Study Status
20	<input type="radio"/>	<input type="radio"/>	February 15, 2012	Study Status
21	<input type="radio"/>	<input type="radio"/>	March 1, 2012	Study Status
22	<input type="radio"/>	<input type="radio"/>	March 14, 2012	Contacts/Locations and Study Status
23	<input type="radio"/>	<input type="radio"/>	March 30, 2012	Study Status and Contacts/Locations
24	<input type="radio"/>	<input type="radio"/>	April 16, 2012	Study Status
25	<input type="radio"/>	<input type="radio"/>	April 30, 2012	Contacts/Locations and Study Status
26	<input type="radio"/>	<input type="radio"/>	May 10, 2012	Outcome Measures, Study Status and Study Description
27	<input type="radio"/>	<input type="radio"/>	June 18, 2012	Study Status
28	<input type="radio"/>	<input type="radio"/>	February 1, 2013	Study Status, Contacts/Locations and References
29	<input type="radio"/>	<input type="radio"/>	February 7, 2013	Study Status
30	<input type="radio"/>	<input type="radio"/>	February 22, 2013	Study Status

Version	A	B	Submitted Date	Changes
31	<input type="radio"/>	<input type="radio"/>	March 14, 2013	Recruitment Status, Contacts/Locations and Study Status
32	<input type="radio"/>	<input type="radio"/>	March 28, 2013	Study Status
33	<input type="radio"/>	<input type="radio"/>	May 7, 2014	Study Status
34	<input type="radio"/>	<input type="radio"/>	June 25, 2014	Study Status
35	<input type="radio"/>	<input type="radio"/>	November 13, 2014	Study Status
36	<input type="radio"/>	<input type="radio"/>	November 20, 2014	Study Status
37	<input type="radio"/>	<input type="radio"/>	December 5, 2014	Study Status
38	<input type="radio"/>	<input type="radio"/>	April 27, 2015	Study Status
39	<input type="radio"/>	<input type="radio"/>	May 5, 2015	Study Status
40	<input type="radio"/>	<input type="radio"/>	May 19, 2015	Study Status
41	<input type="radio"/>	<input type="radio"/>	June 1, 2015	Study Status
42	<input type="radio"/>	<input type="radio"/>	September 3, 2015	Recruitment Status, Study Status and Study Design

Compare

Comparison Format:

☒ Merged☐ Side-by-Side

[Scroll up to access the controls](#)

Changes (Merged) for Study: NCT00729664

August 4, 2008 (v1) -- April 17, 2009 (v2)

No earlier versions to compare

Compare v2 to v3

Changes in: [Study Status](#), [Oversight](#), [Study Description](#), [Arms and Interventions](#), [Outcome Measures](#), [Eligibility](#) and [Contacts/Locations](#)

☐ Show only changed modules

▼ Study Identification

Unique Protocol ID: MDX 1105-01
Brief Title: Phase 1 Open Label Study of MDX-1105 in Subjects With Solid Tumors
Official Title: A Phase 1, Open-Label, Dose-Escalation, Multidose Study of MDX-1105 Administered Every 14 Days in Subjects With Selected Advanced or Recurrent Solid Tumors
Secondary IDs:

▼ Study Status

Record Verification: ~~August 2008~~ April 2009
Overall Status: ~~Not yet recruiting~~ Recruiting
Study Start: ~~November 2008~~ April 2009
Primary Completion: November 2011 [Anticipated]
Study Completion:
First Submitted: August 4, 2008
First Submitted that Met QC Criteria: August 4, 2008
First Posted: August 7, 2008 [Estimate]
Last Update Submitted that Met QC Criteria: ~~August 4, 2008~~ April 17, 2009
Last Update Posted: ~~August 7, 2008~~ April 21, 2009 [Estimate]

▼ Sponsor/Collaborators

Sponsor: Medarex
Responsible Party:
Collaborators:

▼ Oversight

U.S. FDA-regulated Drug:
U.S. FDA-regulated Device:
Data Monitoring: No

▼ Study Description

Brief Summary: The primary purpose of this study is to determine the safety of MDX- 1105, including the maximum dose that can be tolerated when administered for 14 days in subjects with solid tumors- 1105. The study will also evaluate the preliminary effectiveness of the study drug in various types of solid tumors. How MDX-1105 is absorbed, distributed, metabolized and eliminated from the body will also be studied.

Detailed Description: This is a Phase 1 study of a fully human monoclonal IgG4 antibody targeting the Programmed Death-Ligand 1 (PD-L1). Up to 40 subjects with measurable advanced or recurrent solid tumors (renal cell carcinoma, non-small cell lung cancer, malignant melanoma, or epithelial ovarian cancer) will be enrolled in the study.

An accelerated titration design (ATD) will be utilized for this Phase 1 study. Five- Four dose levels are planned and include: 0.3, 1, 3, 40- and 46- 10 mg/kg/dose. Subjects will be assigned to a dose level in the order of study entry.

The maximum total study participation duration for a subject is 2.5 years.

The first cohort of 1 subject will receive MDX-1105 0.3 mg/kg/dose administered as an intravenous infusion every 14 days for a total of 3 infusions. Enrollment into the next cohort cannot begin until 2 weeks have elapsed since the last subject's dosing in the previous cohort. One subject will be enrolled in each cohort until 1 subject experiences a greater than or equal Grade 2 drug-related adverse event, whereupon an additional 2 subjects will be enrolled at that dose level. Thereafter, the study will become the traditional 3+3 design with 3 or 6 subjects treated at this dose level and at all subsequent dose levels depending on the incidence of DLT's. If no DLT's occur in a cohort of 3 subjects, a new cohort of 3 subjects will be treated at the next higher dose level. If 1 of 3 subjects in a cohort experiences a DLT, that cohort will be expanded to 6 subjects. If only 1 of the 6 subjects has a DLT, then the next cohort of 3 subjects will be treated at the next higher dose level. If 2 or more DLT's occur within a cohort, then that dose level will be above the MTD (the highest dose where no more than 1 of 6 subjects has experienced a DLT), and new subjects will be enrolled at the previous lower (tolerated) dose level until that cohort has 3 or 6 subjects. This lower dose level will be considered the MTD if less than or equal to 1 in 6 subjects has a DLT. Depending on toxicity, an intermediate dose level may be defined by a protocol amendment. A subject who withdraws from the study during cycle 1 for reasons other than a DLT will be replaced.

A total of 16 subjects (6 in the dose escalation cohort + 10 in the expansion cohort) will be enrolled at the MTD or the highest dose studied if the MTD is not identified. For the expansion cohort, if 3 or more subjects experience a DLT, enrollment will be stopped and the data will be reviewed by the Medarax Medical Monitor before any subject continues in the study.

A DLT is defined as a greater than or equal to Grade 3 drug-related adverse event occurring within the first cycle (42 days) of dosing (excluding tumor flare defined as local pain, irritation, or rash localized at site of known or suspected tumor or a transient Grade 3 infusion adverse event) using National Cancer Institute Common Terminology Criteria for Adverse Events version 3.0. A DLT will be considered related to study drug unless there is a clear, well-documented, alternative explanation for the adverse event.

Delayed DLT's are drug-related adverse events that occur after Cycle 1; delayed DLT's will not be used to determine the MTD for dose escalation. Delayed DLT's will be collected and evaluated by the Investigators and Medarax on an ongoing basis. If 4 or more such delayed DLT's are noted, accrual will be held pending safety analysis and will be restarted only with Investigator and Medarax approval.

Tumor response will be evaluated using Response Evaluation Criteria in Solid Tumors (RECIST) with modifications. End of cycle tumor response assessments for all subjects will occur within Days 38 to 42. Subjects with initial progressive disease (PD) based on sum of the longest diameter (SLD) or development of new lesions at the end of a cycle, but with otherwise stable or improved performance and clinical status, will continue to be treated with study drug for 1 additional cycle. At the next scheduled imaging evaluation, if there is further increase in the SLD, or development of additional new lesions, then PD has been confirmed and the subject will stop treatment and complete the off-treatment procedures. If the SLD remains the same or decreased and no additional new lesions develop and the subject is otherwise stable or improved in performance and clinical status, then PD has not been confirmed and the subject will continue to be treated with study drug for 1 additional cycle until the next scheduled imaging evaluation.

In the absence of confirmed disease progression, subjects who discontinue MDX-1105 treatment should continue in the study until at least 70 days following the last dose of MDX-1105 or the next scheduled visit has been performed which is even longer. Subjects who discontinue MDX-1105 because of a treatment-related adverse event will remain in the study and continue to be followed until 70 days following the last dose of MDX-1105 or until stabilization or resolution of the event, whichever is longer. Subjects who discontinue MDX-1105 administration due to an adverse event and who have a demonstrated complete response (CR) partial response (PR) or stable disease (SD) will remain in the study and be followed every 3 months for up to 6 months until relapse, confirmed disease progression, or initiation of new therapy, whichever occurs first.

Subjects who achieve a confirmed CR will not receive further cycles of MDX-1105 and will be followed every 3 months for 6 months until relapse, initiation of a new therapy, or a total of 6 months follow up, whichever occurs first. Subjects with a response of PR/SD will continue to receive MDX-1105 treatment until achievement of a confirmed CR, confirmed disease progression, development of a greater than or equal to Grade 3 CTCAE, or intolerability to therapy. If a subject is eligible to receive additional cycles, the first dose of the next cycle should be given 14 days after the last dose of the prior treatment cycle but should be given no later than 21 days. No subject will be permitted dose escalations or de-escalations. The maximum duration of study drug treatment for a subject is 2 years. Those subjects who continue to receive clinical benefit after 2 years of therapy (subjects with CR, PR, or SD) will be followed every 3 months from the last day they received study drug for 6 months until relapse, confirmed disease progression, initiation of a new therapy, or a total of 6 months follow up, whichever occurs first. The maximum total study participation duration for a subject is 2.5 years.

▼ Conditions

Conditions: Renal Cell Carcinoma
Non-Small Cell Lung Cancer
Malignant Melanoma
Epithelial Ovarian Cancer

Keywords:

▼ Study Design

Study Type: Interventional
Primary Purpose: Other
Study Phase: Phase 1
Interventional Study Model: Parallel Assignment
Number of Arms: 1
Masking: None (Open Label)
Enrollment: 40 [Anticipated]

▼ Arms and Interventions

Arms	Assigned Interventions
Experimental: 4 Study Drug 5 cohorts (0.3, 1, 3, 10, and 15 mg/kg/dose) Dosing with study drug	Biological: MDX-1105 MDX-1105 i.v.infusion, 5-4 cohorts, 0.3, 1, 3, 10, 15 mg/kg/dose every 14 days for a total of 3 infusions infusions per cycle

▼ Outcome Measures

- Primary Outcome Measures:
- To determine the safety, maximum tolerated dose(MTD and dose-limiting toxicity(DLT)of MDX-1105
[Time Frame: ~~MDX-1105 administered as an infusion every 14 days for a total of 3 infusions~~ Weekly and bi-weekly]
- Secondary Outcome Measures:
- Evaluate preliminary efficacy in solid tumors on the basis of objective responses
[Time Frame: ~~Days 1, 15 and 29~~ Day 42]

▼ Eligibility

Minimum Age: 18 Years
Maximum Age:
Sex: All
Gender Based:
Accepts Healthy Volunteers: No

Criteria: Inclusion Criteria:

1. Adults at least 18 years of age
2. Life expectancy greater than or equal to 12 weeks
3. ECOG performance status of 0 or 1
4. Criteria specific to each tumor type:

RCC: Subjects must have a histologically confirmed diagnosis of renal cell carcinoma (clear cell component) with advanced or recurrent disease that is not amenable to cure by surgery or other means, and must have failed at least 1 previous systemic therapy ~~including, but not limited to, treatment with sunitinib, temsirolimus, sorafenib, interleukin-2 and/or chemotherapy~~.

NSCLC: Subjects must have refractory or recurrent histologically or cytologically confirmed advanced NSCLC. The malignancy must be deemed unresectable, and subjects must have failed at least 1 prior systemic taxane-based therapy.

MEL: Subjects must have a histologically confirmed diagnosis of melanoma with advanced disease (previously treated, therapy-refractory or recurrent Stage III (unresectable) or Stage IV); disease no longer controlled by surgery, chemotherapy, or radiotherapy; and disease refractory to or relapsed after standard therapy (may include high-dose interleukin-2) ~~All melanomas regardless of primary site of disease will be allowed~~.

epOC: Subjects must have a histologically confirmed diagnosis of epithelial ovarian cancer with advanced or recurrent disease that is not amenable to cure by surgery or other means and must have failed at least 1 prior systemic platinum-based therapy.

5. Must have measurable disease with at least 1 unidimensional measurable lesion per RECIST with modifications
6. ~~May have been treated with up to 3 previous systemic therapies for advanced/recurrent disease or have become intolerant to a systemic therapy~~
7. ~~Prior treated brain metastases must be without MRI evidence of progression for at least 8 weeks and off steroids for at least 4 weeks before study drug administration to be eligible~~
8. ~~May have been treated with radiation therapy (TR) provided there are measurable lesions outside the field of RT~~
9. ~~Prior systemic RT completed at least 4 weeks before study administration. Prior focal radiotherapy completed at least 2 weeks before study drug administration. No radiopharmaceuticals (strontium, samarium) within 8 weeks before study drug administration~~
10. ~~At least 28 days since the last chemotherapy, immunotherapy, biological or investigational therapy, radiation, or at least 2 weeks from approved tyrosine kinase inhibitor therapy before the first dosing of MDX-1105 and have recovered from any toxicity associated with such treatment~~
11. ~~Systemic or topical corticosteroids must be discontinued at least 4 weeks before study drug administration~~
12. ~~Prior major treatment-related surgery completed at least 4 weeks before study drug administration.~~
13. ~~Screening laboratory values must meet the following criteria:~~
~~WBC greater than or equal to 3000/uL Neutrophils greater than or equal to 1500/uL Platelets greater than or equal to 100,000/uL Hemoglobin greater than or equal to 9.0g/dL Creatinine less than or equal to 2mg/dL AST less than or equal to 2.5 X ULN ALT less than or equal to 2.5 X ULN Bilirubin less than 1.5 mg/dL~~
14. Women must meet 1 of the following criteria: post-menopausal for at least 24 consecutive months; surgically incapable of bearing children or utilizing a reliable form of contraception. Women of child bearing potential must agree to use a reliable form of contraceptive during the study Treatment Period and for at least 70 days following the last dose of study drug; and
15. Men must agree to the use of male contraception during the study treatment and for at least 180 days after the last dose of study drug. -

Exclusion Criteria:

1. History of severe hypersensitivity reactions to other monoclonal antibodies
2. Prior therapy with an anti-PD-1, anti-PD-L1, or anti-CTLA-4 antibody
3. Any other malignancy, excluding basal or squamous cell carcinoma of the skin, cervical carcinoma in situ, or localized prostate cancer, from which the subject has not been disease-free for at least 5 years
4. Active infection requiring i.v. systemic therapy within 28 days prior to the first dose of MDX-1105
5. Evidence of bleeding diathesis or coagulopathy
6. Any active autoimmune disease or a documented history of autoimmune disease, or history of syndrome that required systemic steroids or immunosuppressive medications, except for subjects with vitiligo
7. Recent history of attenuated viral vaccination within 30 days prior to the first dose of study drug
8. Known current drug or alcohol abuse
9. Apparent active or latent tuberculosis infection as indicated by any of the following: PPD recently converted to positive; chest x-ray with evidence of infections infiltrate; recent unexplained changes in fever/chill patterns
10. Underlying medical conditions that, in the Principal Investigator's opinion, will make the administration of study drug hazardous or obscure the interpretation of toxicity determination or adverse events
11. Concurrent medical condition requiring the use of immunosuppressive medications, or systemic or topical corticosteroids. Inhaled or intranasal corticosteroids (with minimal systemic absorption may be continued if the subject is on a stable dose). Non-absorbed intra-articular steroid injections will be permitted
12. Use of other investigational drugs within 30 days before study administration
13. Psychiatric illness or social situation that would preclude study compliance; or
14. Pregnant or nursing. -

▼ **Contacts/Locations**

Central Contact Person: Israel Lowy, MD PhD
Telephone: 908-479-2447
Email: ilowy@mederax.com

Central Contact Backup: David Bailis
Telephone: 908-479-2602
Email: dbailis@mederax.com

Study Officials: **Henry C Pitot, MD**
Principal Investigator
Mayo Clinic

Locations: **United States, California**
Study Center
[Recruiting]
Los Angeles, California, United States, 90404

United States, Georgia
Study Center
[Recruiting]
Atlanta, Georgia, United States, 30322

United States, Maryland
Study Center
[Not yet recruiting]
Baltimore, Maryland, United States, 21231

United States, Minnesota
Mayo Clinic
Rochester, Minnesota, United States, 55905
~~Contact: Contact: Henry C Pitot, MD~~
~~Contact: Principal Investigator: Henry C Pitot, MD~~ Study Center
[Not yet recruiting]
Rochester, Minnesota, United States, 55905

United States, New York
Study Center
[Not yet recruiting]
Buffalo, New York, United States, 14623

United States, Ohio
Study Center
[Not yet recruiting]
Columbus, Ohio, United States, 43210

United States, Pennsylvania
Study Center
[Not yet recruiting]
Philadelphia, Pennsylvania, United States, 19107

United States, Texas
Study Center
[Recruiting]
Houston, Texas, United States, 77053

Study Center
[Not yet recruiting]
Houston, Texas, United States, 77203

United States, Washington
Study Center
[Not yet recruiting]
Seattle, Washington, United States, 98109

▼ **IPDSharing**

Plan to Share IPD:

▼ **References**

Links:

Available IPD/Information:

[Scroll up to access the controls](#)

[Scroll to the Study top](#)

[U.S. National Library of Medicine](#) | [U.S. National Institutes of Health](#) | [U.S. Department of Health & Human Services](#)