



YIELD ENHANCER

ACTIVE INGREDIENTS:

Cytokinin (as kinetin)	0.009%
Gibberellic acid	0.005%
Indole-3-butyric acid	0.005%

OTHER INGREDIENTS:

.....	99.981%
Total	100.000%

This product contains: 0.002698 grams cytokinin, 0.001499 grams gibberellic acid, 0.001499 grams indole-3-butyric acid per fluid ounce.

CONTAINS NON-PLANT FOOD INGREDIENTS:

- 0.009% Cytokinin
- 0.005% Gibberellic Acid
- 0.005% Indole-3-butyric Acid

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional Precautionary Statements inside booklet.

Read attached label before using.

EPA Reg. No. 57538-13-89391



Distributed By:
INNICTIS® CROP CARE, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

092316RD051917A

PLANT GROWTH REGULATOR

FIRST AID

IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC web site: www.npic.orst.edu).

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

PERSONAL PROTECTIVE EQUIPMENT

Waterproof gloves are sufficiently chemical-resistant for this product. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride,
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4,6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of all excess treated seed and seed packaging by burial away from streams and bodies of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours unless wearing appropriate PPE.

Exception: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical-resistant gloves made of any waterproof material, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter treated areas until sprays have dried.

CHEMIGATION: APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION

Apply this product only through the following types of irrigation systems: center pivot, traveler, big gun, motorized lateral move, end tow, side (wheel) roll, solid set, and hand move irrigation equipment. Do not apply through any other types of irrigation systems. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Experiment Station specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

A. Center Pivot, Traveler, Big Gun, Motorized Lateral Move, End Tow, and Side (Wheel) Roll Irrigation Equipment: Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank of injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and acreage contained in circle or run. Mix recommended amount of *ARRIVE PLUS* Yield Enhancer for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run, but continue to operate irrigation system until *ARRIVE PLUS* Yield Enhancer has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

B. Solid Set and Hand Move Irrigation Equipment: Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of *ARRIVE PLUS* Yield Enhancer for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to insure that *ARRIVE PLUS* Yield Enhancer will remain in suspension during the injection cycle. *ARRIVE PLUS* Yield Enhancer can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until *ARRIVE PLUS* Yield Enhancer is cleared from last sprinkler head.

SAFETY DEVICES

(1) The systems designated above must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. (2) All pesticide injection pipelines must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. (3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. (5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

SYSTEMS CONNECTED TO PUBLIC WATER SOURCES

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank at at least twice the inside diameter of the fill pipe. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

For additional instructions on safety precautions, refer to statements (2), (3), (4) (6) and (7) in the section SAFETY DEVICES.

GENERAL INFORMATION

ARRIVE PLUS Yield Enhancer is a biostimulant containing plant growth regulators. *ARRIVE PLUS* enhances plant growth and development by stimulating cell division, cell differentiation and enlargement; nutrient uptake and nutrient utilization. It is especially effective when applied with foliar fertilizer, but it is also compatible with pesticides.

MIXING INSTRUCTIONS: *ARRIVE PLUS* Yield Enhancer is water soluble and suitable for use in conventional liquid application systems including sprinkler irrigation systems. Shake *ARRIVE PLUS* thoroughly and dilute in sufficient water to assure adequate, even coverage without producing excessive runoff. Agitate the spray mixture during application and apply within 12 hours of dilution. *ARRIVE PLUS* can be applied tank mixed with most insecticides, fungicides, herbicides and foliar fertilizers but must be the last addition to the spray mixture.

APPLICATION INSTRUCTIONS

Apply *ARRIVE PLUS* Yield Enhancer (by ground or air) to foliage diluted in 2 to 10 gallons of water per acre. Larger volumes of water may be used if not associated with excessive runoff. For best results, apply in the early morning or late evening. When applying *ARRIVE PLUS* Yield Enhancer in a band or as a foliar-directed spray, reduce the application rate from the labeled broadcast rate in proportion to the percent of the field surface area covered by the foliar spray, but not below the minimum rate listed in the table.

VEGETABLES

Crop	No. Of Applications	Broadcast Spray		Band Spray		In Seed Furrow		Transplant Water		Spray Timing
		Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	
Beans	1	8	560	4	280					Between the 3rd trifoliate leaf stage and flower bud formation.
	2-3	4	280	3	210	2-8	140-560			Begin at the 3rd trifoliate leaf stage and then at 7-10 day intervals.
	4-6	3	210	2	140					Begin at the 2nd trifoliate leaf stage and then at 7-14 day intervals.
Broccoli	3	6	425	4	280	2-8	140-560	2-8	140-560	Begin at the 4-5 leaf stage and then at 10-14 intervals.
Brussels Sprouts	3	6	425	4	280	2-8	140-560	2-8	140-560	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Cabbage	3	6	425	4	280	2-8	140-560	2-8	140-560	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Cauliflower	3	6	425	4	280	2-8	140-560	2-8	140-560	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Corn, Sweet	1	8	560	5	350	2-8	140-560			2-6 leaf stage.
	2 or more	4	280	3	210					Begin at 2-6 leaf stage and then at 7-21 day intervals through the end of tasseling.
Cucumbers	1	8	560	4	280					Between flower bud initiation and first bloom.
	2-3	4	280	3	210	2-8	140-560	2-8	140-560	Begin at flower bud initiation and then at 7-10 day intervals.
	3-4	4	280	3	210					Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then at 7-10 day intervals.
Lettuce	3	6	425	4	280	2-8	140-560	2-8	140-560	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Melons	1	8	560	4	280					Between flower bud initiation and first bloom.
	2-3	4	280	3	210	2-8	140-560	2-8	140-560	Begin at flower bud initiation and then at 7-10 day intervals.
	4-6	3	210	2	140					Begin 2 weeks after emergence and then at 10-14 day intervals.
Onions	3	6	425	4	280	2-8	140-560			Begin 2 weeks after emergence and then at 10-14 day intervals.
Peppers	4-6	4	280	3	210	2-8	140-560	2-8	140-560	Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then at 7-14 day intervals.
Potatoes	1	16	1,135	8	560	2-8	140-560			Tuber initiation.
	3	8	560	4	280					Begin at stolon formation (8-10 leaf stage) and then at 10-14 day intervals.
Squash	1	8	560	4	280	2-8	140-560			Between flower bud initiation and then at 7-10 day intervals
	2-3	4	280	3	210					Begin 2 week after emergence and then at 7-1 day intervals.
	4-6	3	210	2	140					Begin 2 weeks after emergence and then at 7-14 day intervals.
Tomatoes	1	8	560	4	280					Between flower bud initiation and first bloom.
	2-3	4	280	3	210	2-8	140-560	2-8	140-560	Begin at flower bud initiation and then at 7-10 day intervals.
	4-6	3	210	2	140					Begin 2 weeks after emergence, or at transplant, and then at 7-14 day intervals.

FIELD CROPS

Crop	No. of Applications	Broadcast Spray		Band Spray		In Seed Furrow		Transplant Water		Spray Timing
		Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	
Beets, sugar	1	16	1,135	8	560	2-8	140-560			6-8 leaf stage.
	2-3	8	560	4	280					Begin at the 2 leaf stage and then at 7-14 day intervals.
Canola	3	6	425	4	280	2-8	140-560			Begin at the 3-5 leaf stage and then at 10-14 day intervals.
Corn	1	8	560	5	350	2-8	140-560			2-6 leaf stage.
	1	8	560	4	280	2-8	140-560			Between flower initiation and final bloom.
	2-3	4	280	3	210					Early bloom and 7-14 days later.
Cotton	3-4	4	280	3	210					Begin at pinhead square and then at 7-10 day intervals.
	4-6	4	280	3	210	2-8	140-560			Begin 30 days after planting and then at 7-14 day intervals.
Peanuts	1	8	560							2-5 leaf stage or panicle initiation.
	2	4	280							2-5 leaf stage or panicle initiation.
Rice	1	8	560	5	350	2-8	140-560			2-6 leaf stage.
Sorghum	1	8	560	5	350	2-8	140-560			2-6 leaf stage.
Soybeans	1	8	560	5	350	2-8	140-560			3-7 trifoliate leaf stage (V4-V8)
	2	4	280	3	210					3-7 trifoliate leaf stage (V4-V8) and 10-17 days later.
Tobacco	2	8	560	5	350			2-8	140-560	1 st 40 days after planting 2 nd after topping.
	1-2	8	560	6	180	2-8	140-560			Start at tillering in the fall and/or spring and when 2 to 3 leaves form on main stem.

SMALL FRUITS, VINES AND TREE FRUITS

Bananas: Apply 4 to 8 fl. oz./A (300 to 600 ml/hectare) in a band around the root mat and repeat every 10 to 14 days for a total of 10 applications.

Pineapple: Spray after transplanting with 4 to 8 fl. oz./100 gallons (120 to 240 ml/300 liters) and repeat every 10 to 14 days or apply in an irrigation system to provide 4 to 8 fl. oz./A (324 to 648 ml/hectare).

Strawberries: Begin sprays at first bloom at 4 fl oz/A (286 ml/hectare) broadcast or 2 fl oz/A (143 ml/hectare) as band sprays directed at the row. Repeat sprays at 2 to 4 week intervals for a total of 3 to 6 sprays.

Oranges: Apply at 1 to 2 pints/100 gallons (126 ml to 252 ml/100 liters) at first bloom and repeat at each flush of new growth.

Grapes: Apply 2 to 8 fl. oz./A every 7 to 21 days from bud break through veraison.

Olive: Apply 2 to 8 fl. oz./A every 7 to 21 days from bud break through harvest.

Pomegranate: Apply 2 to 8 fl. oz./A every 7 to 21 days from bud break through harvest.

POSTHARVEST

Apply *ARRIVE PLUS* Yield Enhancer as a post-harvest application at 2 to 8 ounces per acre to promote uniformity and vigor in the following cropping season.

YOUNG TREES AND ORNAMENTALS

Shrubs, Established: For increased vigor, rapid growth and healthy plant appearance, spray foliage with transplant solution (1 fl. oz. *ARRIVE PLUS* Yield Enhancer/2 gallons of water) to point of run-off two to three times per year.

Flowering Plants (Roses, Azaleas, etc.): 3-4 year old—using stock solution (1 fl. oz. *ARRIVE PLUS* Yield Enhancer/2 gallons of water) —Take 4 fl. oz. stock solution in 1 gallon of water and water in around root zone one time per year, preferably early spring.

Sod:* To improve growth, heavy rooting, broadcast 8 fl. oz./acre.

1. To speed up regrowth after harvest, broadcast 4 fl. oz./acre.

2. To boost with another 4 fl. oz./acre, broadcast six weeks later.

Turf:* For quick "lie down" after laying and to get turf off to a quick start, use *ARRIVE PLUS* Yield Enhancer as follows:

1. Broadcast 2 fl. oz./5,000 sq. ft. and water in.

2. Second application—30 days later—2 fl. oz./5,000 sq. ft. and water in.

*Do not apply this product through any type of irrigation system.

NOTICE: *ARRIVE PLUS* YIELD ENHANCER IS NOT A FERTILIZER. USE IN COMBINATION WITH A GOOD FERTILIZER PROGRAM WHERE INDICATED.

GOLF COURSES

Greens: Make an initial treatment with *ARRIVE PLUS* Yield Enhancer to promote root development and protect against "winter kill" using 2 fl. oz./5,000-7,000 sq. ft. Thereafter, use 1 fl. oz./green every 30 days.

Tees: Use *ARRIVE PLUS* Yield Enhancer at ½ fl. oz./1,200-1,500 sq. ft. of tee area every 30 days to maintain a healthy mass root growth.

Fairways: To establish the necessary root growth to fully utilize applied fertilizer, use *ARRIVE PLUS* Yield Enhancer at 8 fl. oz./acre two times the first year. Thereafter, use 8 fl. oz./acre one to two times a year.

TRANSPLANTS

ARRIVE PLUS YIELD ENHANCER MAY BE USED WHEN TRANSPLANTING ORNAMENTALS AND YOUNG TREES FOR A FAST AND HEALTHY START. Use 1 fl. oz./two gallons water stock solution as follows: At time of transplanting—

1. Bare (naked) roots - dip or spray with stock solution (1 fl. oz. **ARRIVE PLUS** Yield Enhancer/2 gallons of water).
2. Balled plants - spray ball at time of transplant.
3. Mist (not run-off) foliage lightly at time of transplant.
4. Using stock solution, apply 1 gallon **ARRIVE PLUS** Yield Enhancer in furrow/acre.

SEED TREATMENT

Use only on seeds for crops listed elsewhere on the label. Do not use treated seed for food, feed or oil purposes. If this product is intended for commercial seed treatment, treated seed must be labeled in accordance with the requirements of the Federal Seed Act and applicable State seed laws. An approved dye must be added to distinguish treated seed and prevent inadvertent use for food, feed, or oil purposes. If this product is intended for "at planting" use, treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of excess treated seed by burial away from streams and bodies of water. A dye is not required.

For crops listed: Apply 2 to 4 fluid ounces per 100 lbs. of seed (.75 to 1.5 ml per kg). Dilute with water and mist spray on seed while mixing. Do not store seed wet as germination can be reduced if not planted soon after treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Do not store in direct sunlight. Avoid freezing temperatures.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Use label language appropriate for container size and type.

Non refillable containers: Do not reuse or refill this container. Clean container promptly after emptying.

Non refillable container equal to or less than 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 120 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Non refillable container greater than 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of INNVICTIS CROP CARE LLC or Seller, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold INNVICTIS CROP CARE LLC and Seller harmless for any claims relating to such factors.

INNVICTIS CROP CARE LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or INNVICTIS CROP CARE LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. INNVICTIS CROP CARE LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither INNVICTIS CROP CARE LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF INNVICTIS CROP CARE LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF INNVICTIS CROP CARE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

INNVICTIS CROP CARE LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of INNVICTIS CROP CARE LLC.

SPECIMEN

INTENTIONALLY LEFT BLANK

SPECIMEN

INTENTIONALLY LEFT BLANK