



Arysta LifeScience

FIFRA 24C REGISTRATION  
EPA SLN No. FL-120001

REGISTRATION FOR SPECIAL LOCAL NEED FOR DISTRIBUTION  
AND USE ONLY WITHIN THE STATE OF FLORIDA

**CAPTEC 4L<sup>®</sup>**  
EPA Reg. No. 66330-239

**ACTIVE INGREDIENT:**

\*Captan ..... 38.75%  
Related Derivatives ..... 0.85%

**OTHER INGREDIENTS:** ..... 60.40%

**TOTAL:** ..... 100.0%

\*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide  
Contains 4 lbs of Captan per gallon.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
All applicable directions, restrictions and precautions on the EPA registered label are to be followed.  
This labeling must be in the possession of the user at the time of pesticide application.

CROP	DISEASE	RATE PER APPLICATION	
		qt/100 Gallons/ Application	MAXIMUM qt/A/ Application
<b>STRAWBERRIES:</b>	Botrytis, Leaf Spot	1.5 - 3.0	3.0
<b>Precautions and Restrictions</b>	Apply by broadcast in sufficient water for thorough coverage by ground or in 10 to 20 gallons of water by air. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7 to 14 day intervals. Under conditions favorable to fruit rot, continue applications through harvest period, treating immediately after each picking. Do not apply more than 72 quarts of Captec 4L per acre per year. Do not apply more than 24 quarts of Captec 4L per acre per berry crop. May be applied up to day of harvest. Restricted Entry Interval is 24 hours.		

If applying as a directed/banded spray; use band rate of Captec 4L according to the following formula:

$$\frac{\text{Plant Bed Width (inches)}}{\text{Row Spacing (inches)}} \times \text{Broadcast rate per acre} = \text{Banded rate of Captec 4L per acre.}$$

24 (c) Registrant:

Arysta LifeScience North America, LLC  
15401 Weston Parkway, Suite 150  
Cary, NC 27513

032212