

## RESTRICTED USE PESTICIDE

Due to acute toxicity. For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

The Drexel logo consists of the word "Drexel" in a white, sans-serif font, centered within a black triangle that points downwards.

# QUIK-QUAT™

(EPA Reg. No. 19713-617)

### FIFRA Section 2(ee) RECOMMENDATION FOR SOYBEANS HARVEST AID

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF LOUISIANA

#### DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

#### SOYBEANS

Use Pattern	THIS PRODUCT Rate per Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval (Days)
Harvest aid	5.4 to 10.7 fl. oz.	Ground: 10 gals.	-

#### Additional Precautions, Restrictions and Directions:

- Do not make more than 3 applications per year.
- Indeterminant varieties: Applications should be made when at least 65% of the seed pods have reached a mature brown color or when seed moisture is 30% or less.  
Determinant varieties: Apply when plants are mature, i.e., beans are fully developed, 1/2 of leaves have dropped, and remaining leaves are yellowing.
- Injury will occur on immature soybeans.
- Mature cocklebur, especially drought-stressed plants, are tolerant to this product and desiccation will not be complete. Always use the higher rate when treating cocklebur.
- Do not apply within 15 days of harvest.
- Do not graze or harvest for forage or hay.

Manufactured By:



QUIK-QUAT and the Drexel logo are trademarks or registered trademarks of Drexel Chemical Company.