FIFRA Section 2(ee) Recommendation

EVITO™ 480 SC Fungicide
EPA Reg. No. 66330-64
This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U. S. Environmental Protection Agency

FOR DISTRIBUTION ONLY IN GEORGIA, NORTH CAROLINA, SOUTH CAROLINA AND VIRGINIA

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for EVITO 480 SC Fungicide. Also observe all directions, precautions and limitations on the labeling of all other products mentioned below. This recommendation must be in the possession of the user at the time of application.

For Control or Suppression of Certain Diseases on Peanuts

Crop	Disease Control	Rates to Use	Application Timing
Peanuts	**Early leaf spot** (*Cercospora arachidicola*) **Late leaf spot** (*Cercosporidium personatum*) **Leaf rust** (*Puccinia arachidis*) **Disease Suppression** **Stem rot** **White mold** **Southern blight** (*Sclerotium rolfsii*) **Rhizoctonia limb rot** (*Rhizoctonia solani*)	3.8 to 5.7 fl oz/A (0.12 to 0.18 lbs ai/A)	For optimum results, begin applications preventively. Apply as needed on a 14-day interval. To limit the potential for development of disease resistance: • In areas with typically 1-4 sprays per year, alternate every application of a solo QoI fungicide with at least one application of another effective mode of action fungicide. • In areas with typically 5 or more fungicide sprays per year, a maximum of 2 sequential applications of a QoI fungicide followed by at least an equal number of another effective mode of action fungicide. Improved white mold control can be obtained with higher rates of Evito (5.7 fl oz/A, per federal label) or with combinations (tank mixed or alternating applications) with other products labeled for white mold.

Notes:
- Do not apply more than 22.8 fl oz (0.72 lbs ai) of EVITO 480 SC Fungicide per crop cycle.
- There is a maximum number of 4 applications per season, and a minimum interval of 14 days between applications.
- Use of a surfactant may increase coverage and efficacy of this product.
- Do not apply within 14 days of harvest.

Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Revised date: May 21, 2014
Expiration Date: May 21, 2015