



This seed has been treated with azoxystrobin, mefenoxam, fludioxonil, and sedaxane fungicides at the manufacturer's recommended rate.

**WARNING. TREATED SEED. DO NOT USE FOR FEED, FOOD OR OIL PURPOSES.**

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

**Storage:** Store treated seed away from feeds and foodstuffs. **Do not allow children, pets or livestock to have access to treated seeds.**

**Personal Protective Equipment:** Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.

**General Information:** Avoid breathing dust from treated seeds. Work upwind while loading planters. Workers may choose to wear a dust mask. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**Environmental:** Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting and in areas such as row ends. Treated seed must be incorporated into the soil at the recommended depth.

**Disposal:** Dispose of all excess treated seed by burying seed away from bodies of water. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwater. Dispose of seed packaging or containers in accordance with local requirements. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed, and (2) no measureable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

**Restrictions:** In the event of a crop failure or harvest of a crop grown from this treated seed, the field may be replanted immediately to canola, cereal grains crop group 15, corn (field, pop, sweet), cotton, dried shelled pea and bean crop subgroup 6C, peanut, potato, rapeseed crop subgroup 20A, sorghum, soybean, and sugarbeet. For all other crops, the minimum plant back interval is 30 days from the date this treated seed was planted.

The maximum number of applications per year is 2.

**Groundwater Advisory:** Azoxystrobin, a degradate of azoxystrobin, and mefenoxam are known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in ground water. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**For Emergencies** or questions related to azoxystrobin, fludioxonil, mefenoxam, and sedaxane applied to this seed, please call 1-800-424-9300.

©2018 Syngenta.

First Defense SBR Custom Blend – Helena – 10 17 18