SUPPLEMENTAL LABELING

KASUGAMYCIN	GROUP	24	FUNGICIDE

KASUMIN® 2L

Supplemental Label to Add DIRECTIONS FOR USE on CHERRY (Crop Subgroup 12-12A); Capulin; Black Cherry; Nanking Cherry; Sweet Cherry; Tart Cherry; and Cultivars, varieties and/or hybrids of these.

This supplemental label expires March 1, 2021 and must not be used or distributed after this date.

For Agricultural Use Only

ACTIVE INGREDIENT:
*Kasugamycin Hydrochloride Hydrate 2.3%
OTHER INGREDIENTS: 97.7%
TOTAL: 100.0%
*(Equivalent to 2.0% kasugamycin)
Contains 0.168 pounds kasugamycin per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **KASUMIN® 2L** BEFORE APPLYING. USE OF **KASUMIN 2L** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **KASUMIN 2L (EPA Reg. No. 66330-404).**

030118V002Su

CHERRY (Crop Subgroup 12-12A); Capulin; Black Cherry; Nanking Cherry; Sweet Cherry; Tart Cherry; and Cultivars, varieties and/or hybrids of these

Disease Control	Application Rate fl oz product (lbs ai)	Specific Use Instructions
Bacterial Blast *(Pseudomonas syringae pv syringae)*	**64 fl oz/A (0.084 lb ai/A)**	▪ Apply at a rate of **64 fl oz/A in at least 100 gal of water by ground application**. Reduced spray volumes may be utilized for small trees where complete coverage can be obtained with less than 100 gal of water/A. Spray volume must be sufficient to provide good coverage of treated foliage. ▪ Begin applications when conditions favor disease development, from white tip to petal fall. ▪ For airblast application: To help reduce off-target drift, direct spray into the canopy, and turn off outward pointing nozzles at row ends and when spraying outer rows.
Disease Suppression	**Application Rate fl oz product (lbs ai)**	**Specific Use Instructions**
Bacterial Canker (*Pseudomonas syringae pv syringae*)	**64 fl oz/A (0.084 lb ai/A)**	▪ Apply at a rate of **64 fl oz/A in at least 100 gal of water by ground application**. Reduced spray volumes may be utilized for small trees where complete coverage can be obtained with less than 100 gal of water/A. Spray volume must be sufficient to provide good coverage of treated area. ▪ Begin applications when conditions favor disease development; prior to cold; and when wet conditions occur in the fall and winter (November through February). Check with your local university cooperative extension service or other local experts for the best preventive timing. ▪ For airblast application: To help reduce off-target drift, direct spray into the canopy, and turn off outward pointing nozzles at row ends and when spraying outer rows.

RESTRICTIONS:

- Do not apply after petal fall.
- Do not apply more than 64 fl oz/A (0.084 lb ai/A) of **KASUMIN 2L** in a single application.
- Do not apply more than 256 fl oz (0.336 lb ai/A) of **KASUMIN 2L** per acre per year.
  In California: Do not apply more than 128 fl oz (0.168 lb ai/A) of **KASUMIN 2L** per acre per year.
- Do not make more than 4 applications of **KASUMIN 2L** per year with a minimum interval of 7 days between applications.
- **In California:** Do not make more than 2 applications of **KASUMIN 2L** per year.
- Do not apply more than 2 consecutive applications of **KASUMIN 2L** before alternating with another fungicide/bactericide of a different mode of action. If additional bactericide applications are needed, utilize an alternate product registered for this use. Two more consecutive **KASUMIN 2L** applications may be made subsequent to use of the alternate registered product.
- Do not use an alternate tree-row application method.
- Do not apply kasugamycin in orchards in which the soil has been fertilized with animal waste/manure or human biosolids.
- Animal grazing in treated areas is prohibited. The public must be notified by posting restriction signs along the perimeter of the treatment area.
- Do not apply **KASUMIN 2L** by aerial application.
- **KASUMIN 2L** must be applied at the equivalent of 64 fluid ounces per 100 gallons of water per acre when using less than 100 gallons per acre spray volume.
- Do not apply **KASUMIN 2L** through any type of irrigation system, including chemigation.
- Preharvest Interval (PHI) = 30 days.
- Use only the specified and full-strength application rates.

Produced For:
ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.
The **KASUMIN** logo is a registered trademark of an Arysta LifeScience Group Company.
KASUMIN is a registered trademark of Hokko Chemical Industry Co., Ltd.
©2018 Arysta LifeScience Group Company.