

**ADDITIONAL APPLICATIONS**

DO NOT apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).

DO NOT apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.

For continued weed control, additional application of Quali-Pro Oxadiazon 2G may be made 60 to 120 days after the previous application in the following situations:

2 REPEAT TREATMENTS					
Container-Grown Established		Container-Grown Newly Transplanted		Field-Grown Established and Newly Transplanted	
Species	Varieties	Species	Varieties	Species	Varieties
Abelia	Grandiflora	Common Boxwood		Holly	Nellie R. Stevens, Dwarf Burford, Burford
Common boxwood		English Ivy			
English Ivy		Juniper	Tam		
Holly	Helleri	Pine	Slash		
Juniper	Tam	Rhododendron	Obtusa Kurume Formosa		
Pine	Slash				
Rhododendron	Obtusa Kurume Formosa Kurume Snow				

3 REPEAT TREATMENTS	
Container-Grown Established	
Species	Varieties
Abelia	Grandiflora
Holly	Helleri, Rotundifolia
Rhododendron	Kurume Formosa, Kurume Snow
Greenhouse-Grown Established	
Species	Varieties
Rose	Sonia, Forever Yours, Coed, Golden Wave
Forest Nurseries Seedlings- At least 4 weeks old	
Species	Varieties
Pine	Loblolly, Slash, Eastern White

**STORAGE AND DISPOSAL**

DO NOT contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in dry area. DO NOT store near other pesticides or seeds.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:**

**Nonrefillable Container (flexible-bag-all weights):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (rigid-fifty lbs. or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Nonrefillable Container (rigid-greater than fifty lbs.):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**LIMITATION OF WARRANTY AND LIABILITY**

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Quali-Pro® is a registered trademark of Makhteshim Agan of North America, Inc.



**Oxadiazon 2G**  
*Herbicide*

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers. Not for use on home lawns.

Quali-Pro Oxadiazon 2G is a selective, preemergence herbicide for control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines, and trees.

**ACTIVE INGREDIENT:**

Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-isopropoxyphenyl)-2-1,3,4-oxadiazoline-5-one]: ..... % BY WT.

**OTHER INGREDIENTS:** ..... 98.0%

**TOTAL:** ..... 100.0%

EPA Reg. No. 66222-211

EPA Est. No. 48234-GA-001<sup>®</sup>; 37429-GA-001<sup>®T</sup>; 37429-GA-002<sup>®D</sup>  
Letter(s) in lot number correspond(s) to superscript in EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING – AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call Prosar at 1-877-250-9291.	

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for: **Makhteshim Agan of North America, Inc.**  
3120 Highwoods Blvd, Suite 100  
Raleigh, NC 27604

**Net Contents: 50 Pounds**

EPA 012607/Notif  
090408/030811/040412/Rev F

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**WARNING**  
 Causes substantial but temporary eye injury. DO NOT get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**  
**Loaders, applicators, and other handlers must wear:**  
 • Long-sleeved shirt and long pants  
 • Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**  
**Users Should:**  
 • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.  
 • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.  
 • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**  
 This pesticide is toxic to fish. DO NOT apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwater or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**  
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the worker protection standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:  
 • coveralls  
 • shoes plus socks, and  
 • chemical-resistant gloves made of any waterproof material

**GENERAL INFORMATION**  
 Quali-Pro Oxadiazon 2G is a selective, pre-emergent herbicide for control of annual grasses and many broadleaf weeds in commercial turf and on a wide variety of woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

For best results, water-in the product as soon as practical after application.

**State Specific Restrictions:**  
 • The state of Arizona has not approved this product for use on agricultural commodities. If grown for commercial production, use on the following sites/crops is prohibited: container-grown, newly transplanted, and field-grown ornamental shrubs, vines, and trees.

**DO NOT apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).**

**DO NOT apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.**

**COMMERCIAL TURF**  
 For use in established perennial Bluegrass, Bermudagrass, perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysia turf; such as fairways, parks, golf courses, and lawns. Quali-Pro Oxadiazon 2G controls Crabgrass (*digitaria* spp.), Annual Bluegrass (*Poa annua*), Goosegrass (*Eleusine indica*); and it also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis* spp.), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus officinis*), and Pigweed (*Amaranthus* spp.).

Apply to dry foliage only with a properly calibrated granular applicator at recommended rate in early spring before weed seed germination. If rain is not expected shortly after application, irrigation to remove herbicide granules from foliage will increase effectiveness.

For Crabgrass, Goosegrass, and Crowfoot control, apply 100-200 pounds Quali-Pro Oxadiazon 2G per acre (2 to 4 pounds active ingredient) or 2 1/4 to 4 1/2 pounds per 1000 sq. ft. Late applications are not recommended and may not be fully effective.

For Annual Bluegrass control, apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 1/2 pounds/1000 sq. ft. in early spring or late summer prior to seed germination.

For Pigweed, Florida Pusley, Oxalis, Stinging Nettle, and Carpetgrass control, apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 1/2 pounds per 1000 sq. ft. in early spring prior to weed seed germination.

For St. Augustinegrass turf, DO NOT exceed the rate of 150 pounds/acre (3 pounds active ingredient) of Quali-Pro Oxadiazon 2G.

Discoloration of Bermudagrass and St. Augustinegrass may result from use of this herbicide, but this discoloration is temporary and normally outgrown within 2 to 3 weeks.

Red Fescue and Bentgrass are susceptible to Quali-Pro Oxadiazon 2G and may be severely damaged. Avoid use of this product on turf containing Red Fescue or Bentgrass except where they are undesirable grasses. DO NOT apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after

treatment. DO NOT apply to Dichondra or Centipedegrass. Quali-Pro Oxadiazon 2G should not be used on putting greens or tees because of varying cultural practices used.

**WOODY ORNAMENTAL SHRUBS, VINES, AND TREES**  
**APPLICATION:**

Apply Quali-Pro Oxadiazon 2G anytime during the year prior to weed seed germination for weed control. Quali-Pro Oxadiazon 2G can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of Quali-Pro Oxadiazon 2G will improve weed control activity. Quali-Pro Oxadiazon 2G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Quali-Pro Oxadiazon 2G. DO NOT apply to wet foliage or under conditions in which granules will collect on leaves.

**AMOUNT TO USE:**  
**Ground Application-** Apply Quali-Pro Oxadiazon 2G broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 1/4 to 4 1/2 pounds per 1000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

Quali-Pro Oxadiazon 2G at 100 to 200 pounds (2.0 to 4.0 pounds of active ingredient) per acre is effective in controlling annual grasses and broadleaf weeds such as:

**Annual Broadleaf Weeds**

Bittercress  
 Carpetweed  
 Common Groundsel  
 Common Purslane  
 \*Eveningprimrose  
 \*Galinsoga  
 Golden Ragwort  
 Lambsquarters  
 Pennsylvania Smartweed  
 Petty Spurge  
 Redroot Pigweed  
 Speedwell  
 Spotted Catsear  
 \*Swinecress  
 Yellow Woodsorrel

*Cardamine* spp.  
*Mollugo verticillata*  
*Senecio vulgaris*  
*Portulaca oleracea*  
*Oenothera* spp.  
*Galinsoga* spp.  
*Senecio aureus*  
*Chenopodium album*  
*Polygonum pensylvanicum*  
*Euphorbia peplus*  
*Amaranthus Retroflexus*  
*Veronica* spp.  
*Hypochoeris radicata*  
*Coronopus didymus*  
*Oxalis stricta*

**Annual Grasses**

*Poa annua*  
*Echinochloa crus-galli*  
*Digitaria* spp.  
*Panicum dichotomiflorum*  
*Setaria viridis*

\*Use 150 to 200 pounds Quali-Pro Oxadiazon 2G (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

Note: For the control of liverwort (*Marchantia* spp.) in California, Goosegrass (*Eleusine Indica*), and Prostrate Spurge (*Euphorbia supina*), use 200 pounds Quali-Pro Oxadiazon 2G (4.0 pounds of active ingredient) per acre.

**CROP TOLERANCE**  
 Quali-Pro Oxadiazon 2G is recommended for use on a wide variety of container and field-grown woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G may be used at the recommended rates on the following ornamental species:

**ESTABLISHED Species**  
 Agapanthus  
 Andromeda  
 Arborvitae  
 Arbovitae  
 Arctostaphylos  
 Azalea

Barberry

Boxwood  
 Callistemon  
 Camellia  
 Ceanothus Cedar  
 Cotoneaster

Cypress  
 Dogwood  
 Douglas Fir  
 Eucalyptus  
 Euonymus

Forsythia  
 Gardenia  
 Ginko  
 Heath Heather  
 Hederia  
 Honeysuckle  
 Holly

Jasmine  
 Juniper

Justica  
 Leucothoe  
 Myrtle  
 Magnolia  
 Mahonia  
 Osmanthus  
 Periwinkle  
 Photinia

**CONTAINER GROWN**

Pine

Pittosporum  
 Podocarpus  
 Privet

Pyracantha  
 Quercus Raphiolepis  
 Rhododendron

Rose

Star Jasmine  
 Syzygium  
 Viburnum  
 Xylosma

**Species**  
 Andromeda  
 Arborvitae  
 Azalea

Monterey  
 Red Osier  
 Pseudotsugamenziesii  
 Sideroxylon, Nicholla  
 Greenlane, Microphylla Variegata,  
 Emeraldgaety, Dwarf, Gold spot,  
 Golden Green, Silver King,  
 Wintercreeper Vegetus, Emerald  
 Cushion, Spreading

Lynwood gold  
 Amy Yashioka, August Beauty, Mystery  
 Biloba  
 Mrs. Maxwell  
 English Ivy  
 Purpleleaf, Purpurea  
 Dwarf Okinowan, Convexa, Rotundifolia,  
 Hetzi, Helleri, Burford, English, Hellus,  
 Dwarf Yaupon, Nellie R. Stevens,  
 Boxleafed, Green Island, Japanese

Yellow Star, Star  
 Barharbor, Wiltoni, Torulosa, Pfizeriana  
 Compacti, Pfizeriana Aurea, Torulosa  
 Variegata, Pfizeriana Glauca, Gulf  
 Stream, Tam, Armstrongii, Shore  
 Andora, San Jose, Blue Haven, Stricta,  
 Jade River, Broadmoor, Meyer,  
 Rocket, Plumosa, Royal Beauty,  
 Praecox, Greek, Apiculata Blackburn,  
 Plumosa Compacta, Procumbens,  
 Blue Pfizer  
 Brandegeana  
 Catesbaei, Axillaris  
 Compacta Noellii  
 Dr. Merrilli, Samuel Somers, Grandiflora  
 Oregon Grape-Holly  
 Fortunes  
 Vinca Minor  
 Frazerii

Leucothoe  
 Magnolia  
 Mahonia  
 Pine  
 Privet  
 Pyracantha  
 Rhododendron

Rose

**ESTABLISHED Species**  
 Arborvitae  
 Ash  
 Azalea  
 Barberry  
 Birch  
 Camellia

Cotoneaster  
 Crabapple  
 Dogwood

Slash, Japanese Black, Montana,  
 Scotch, Digger, Shore, Thumbergii  
 Variegated, Japanese  
 Macrophylla  
 Japanese, Waxleaf, Glossy, Vicary,  
 Golden Vicary, Curly Leaf  
 Victory, Scarlet Firethorn, Wyatt, Graberi  
 India Hawthorn  
 Cunningham, Blue Diamond, Snow,  
 Christmas, Chionoides, 'PJM', Dr.  
 Rutgers, Coral Bells, President Clay  
 Summer Snow, Bigleaf Winter Creeper,  
 Charlotte Armstrong  
 Yellow Star Jasmine  
 Paniculatum  
 Sandankwa, Laurestinus, David  
 Congestum

Euonymus

Forsythia

Holly  
 Honeysuckle  
 Juniper

Lilac  
 Maple  
 Oak  
 Periwinkle  
 Pine

Pittosporum  
 Privet  
 Rhododendron  
 Rose

Russian Olive Spruce  
 Taxus  
 Tsuga  
 Viburnum  
 Xylosma

**Ground Covers**  
 Carpobrotus  
 Osteo  
 Spermum Sedum  
 Algerian Ivy  
 Lynwood Gold  
 Rotundifolia, Helleri, English, Japanese  
 Plumosa, Plumosa Compacta, Old Gold,  
 Pfizeriana Aurea, Torulosa,  
 Armstrongii, Tam, Wiltoni, Procumbens

Axillaris  
 Dr. Merrilli  
 Oregon Grape-Holly  
 Slash  
 Japanese, Waxleaf, Glossy  
 Graberi  
 Snow, Blue Diamond, Chionoides, 'PJM',  
 Dr. Rutgers  
 Summer Snow, Bigleaf Winter Creeper,  
 Charlotte Armstrong

**FIELD GROWN**

**Varieties**  
 Pyramidal  
 Green  
 Hinodi geri  
 Red, Mentor  
 Clump, Paper  
 Tiffany, Marie Bracey, Elizabeth Dowd,  
 Tom Knudsen, Crimson Robe, R.D.  
 Wheeler  
 Lowfast, Skogholm  
 Flowing, Toringo  
 Redstemmed, Red, Flowering

Burning Bush, Dwarf Burning Bush,  
 Jewell

Showy Boarder, Lynngold, Lynwood Gold,  
 Goldenbell

Boxleafed, Green Island, Japanese  
 Zabeli

Plumosa, Mint Julip, Arcardia, Variegata;  
 Plumosa Compacta, Andora, Wiltoni,  
 Hetzi

Persian, Dwarf Korean  
 Amur  
 Red  
 Vinca Minor  
 Austrian, Scotch, Norway, Stone,  
 Japanese, Black, Lobloily, Slash,  
 Eastern White

Undulatum  
 California, Japanese  
 Rosebud  
 Texas Centennial, Sutters Gold, Show  
 Girl

Black Hills, Norway, Serbian  
 Taunton Spreading Yew, Hicks  
 Canadian Hemlock  
 European Cranberry Bush

Edule

Album

Terminalis  
 Reptans  
 Fortunei Radicans

**NEWLY TRANSPLANTED Varieties**  
 Tiffany, Marie Bracey, Elizabeth Dowd,  
 R.D. Wheeler, Tom Knudsen, Crimson  
 Robe  
 Redstemmed, Red, Flowering  
 Showy Border, Lynwood Gold,  
 Goldenbell  
 English Ivy  
 Green Island, Japanese  
 Plumosa, Andora, Wiltoni, Hetzi  
 Red  
 Vinca Minor  
 White Austrian, Eastern White  
 California, Japanese  
 Rosebud  
 Black Hills, Norway, Serbian  
 Hicks  
 Canadian Hemlock

# QUALI-PRO®

# Oxadiazon 2G

## Herbicide

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers. Not for use on home lawns.

Quali-Pro Oxadiazon 2G is a selective, preemergence herbicide for control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines, and trees.

ACTIVE INGREDIENT:	% BY WT.
Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-isopropoxyphenyl)-2-1,3,4-oxadiazoline-5-one]:	2.0%
OTHER INGREDIENTS:	98.0%
TOTAL:	100.0%

EPA Reg. No. 66222-211

EPA Est. No. 48234-GA-001<sup>F</sup>; 37429-GA-001<sup>BT</sup>; 37429-GA-002<sup>BO</sup>  
Letter(s) in lot number correspond(s) to superscript in EPA Est. No.

## KEEP OUT OF REACH OF CHILDREN

### WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call Prozar at 1-877-250-9291.	

For additional precautionary, handling, and use statements, see inside of this booklet.

**Manufactured for:** Makhteshim Agan of North America, Inc.  
3120 Highwoods Blvd, Suite 100  
Raleigh, NC 27604

## Net Contents: 50 Pounds