

**RESTRICTED USE PESTICIDE**

**Toxic to fish and aquatic organisms**

For retail sale to and use only by certified applicators or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

# REVEAL

## ENDURX

### Insecticide/Miticide

#### REVEAL ENDURX IS NOT FOR SALE OR USE IN CALIFORNIA

For use to control listed insects and mites on artichokes, beans, brassicas, caneberries, canola, cilantro, citrus, coriander, corn, cotton, crambe, cucurbits, dried beans and succulent peas and beans, fruiting vegetables, head lettuce, hops, leafy brassicas, mayhaw, okra, pears, rapeseed, spinach, tobacco, tomatoes, and tuberous and corm vegetables.

For use to control listed insect pests on Ornamentals and Trees (including Field and Container Grown Nursery Stock, Christmas Trees, Interiorscapes and Plantscapes, Lawns, Trees and Shrubs, and on Golf Courses and Sod Farms)

**ACTIVE INGREDIENT:**

|  |                |
|--|----------------|
| Bifenthrin: (2 methyl[1,1 -biphenyl]-3-yl)methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl-cyclopropanecarboxylate* | <b>%BY WT.</b> |
| .....  | 25.0%          |

|                                   |       |
|-----------------------------------|-------|
| <b>OTHER INGREDIENTS**:</b> ..... | 75.0% |
|-----------------------------------|-------|

|                    |        |
|--------------------|--------|
| <b>TOTAL</b> ..... | 100.0% |
|--------------------|--------|

\*Cis isomers 97% minimum, trans isomers 3% maximum.

\*\*Contains xylene range aromatic solvents.

This product contains 2 pounds active ingredient per gallon.

### KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

See inside label booklet for additional precautionary statements.

EPA Reg. No. 89168-19-89391



**Distributed By:**  
INNICTIS CROP CARE, LLC  
1220 West Ash, Suite B  
Windsor, CO 80550

210121121812V2

35457

## FIRST AID

|  |   |
|--|---|
| <b>IF SWALLOWED</b>  | <ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give any liquid to the person.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul> |
| <b>IF IN EYES:</b>   | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>                 |
| <b>IF ON SKIN OR CLOTHING:</b>   | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>IF INHALED:</b>   | <ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>              |
| <b>NOTE TO PHYSICIAN:</b><br>This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and should be avoided. This product contains a petroleum distillate; vomiting may cause aspiration pneumonia. |   |

## HOT LINE NUMBER

Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: [www.npic.orst.edu](http://www.npic.orst.edu)) or call your poison control center at 1-800-222-1222.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

May be fatal if swallowed. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eye-wear (goggles, face shield, or safety glasses). Harmful if inhaled or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

### Handlers who may be exposed to the dilute through application or other tasks must wear:

- Long-sleeved shirt and long pants, or coveralls worn over short-sleeved shirt and short pants.
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber, or viton
- Chemical resistant footwear plus socks

**Handlers who may be exposed to the concentrate through mixing, loading, application, or other tasks must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber, or viton
- Chemical resistant footwear plus socks
- Protective eyewear
- Chemical resistant apron when mixing and loading and cleaning equipment

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

The use of REVEAL ENDURX is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

**PHYSICAL/CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to users of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls, chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber or Viton, and shoes plus socks.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, and greenhouses.

Do not enter or allow people or pets on treated areas until the spray has dried.

### **RESISTANCE MANAGEMENT**

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

### **APPLICATIONS INSTRUCTIONS – FOOD CROPS**

Application rates will vary according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations and higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip. In New York State, this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

### **CHEMIGATION USE DIRECTIONS**

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

For LEPA irrigation, a minimum of 0.75 inch of water per acre is recommended. Where non-emulsified oils are used as the diluent, 1 to 2 pints per acre is recommended.

Results from utilizing chemigation have been variable and depend upon the set up and calibration of equipment. Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers, or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent area.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain function interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

REVEAL ENDURX should be applied continuously for the duration of the water application. REVEAL ENDURX should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

## **BUFFER ZONES**

### **Vegetative Buffer Strip**

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field

edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing bifenthrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers:

Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp. <http://www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf>

#### **Buffer Zone for Ground Application (groundboom, overhead chemigation, or airstblast)**

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

#### **Buffer Zone for ULV Aerial Application**

Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

#### **Buffer Zone for Non-ULV Aerial Application**

Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

### **SPRAY DRIFT REQUIREMENTS**

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made in cotton. Use of ultra low volume (ULV) application on corn and hops is prohibited.

#### **Wind Direction and Speed**

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph.

#### **Temperature Inversion**

Do not make aerial or ground applications into temperature inversions.

Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Do not make aerial or ground applications to corn if heavy rainfall is imminent.

#### **Droplet Size**

Use only Medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

### **Ground Applications**

For ground applications, wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

### **Aerial Applications**

The spray boom must be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

### **ROTATIONAL CROPS**

Crops with existing bifenthrin tolerances may be rotated at any time. All other crops may be rotated 30 days following the final application of REVEAL ENDURX.

### **TANK MIXTURES**

REVEAL ENDURX may be applied in tank mixtures with other products approved for use on registered crops. Observe all restrictions and precautions which appear on the labels of these products. Test for compatibility of products before mixing.

## INDEX TO CROPS LISTED ON THIS LABEL

| <u>CROP</u>                              | <u>PAGE</u> | <u>CROP</u>                          | <u>PAGE</u> |
|--|-------------|--------------------------------------|-------------|
| Artichoke                                | 8           | Cucurbits                            | 20-21       |
| Brassica Crops                           | 9           | Grapes                               | 24          |
| Caneberries                              | 10          | Hops                                 | 24-25       |
| Canola, Crambe, Rapeseed                 | 11          | Leafy Brassicas                      | 25-26       |
| Cilantro, Coriander                      | 12          | Lettuce, Head                        | 26          |
| Citrus (Not for this use in Florida)     | 12-13       | Mayhaw                               | 27          |
| Dried Beans and Peas                     | 21-22       | Okra                                 | 27          |
| Fruiting Vegetables                      | 22-24       | Peanut                               | 28          |
| (Eggplant, Pepper (Bell & Non-Bell),     |             | Pears                                | 29          |
| Groundcherry, Pepino, Tomato, Tomatillo) |             | Root Crops                           | 30          |
| Field Corn (Grain and Silage), Popcorn,  | 14          | Soybean                              | 31-32       |
| Field Corn Grown for Seed (At Plant Use) |             | Spinach                              | 32          |
| Field Corn (Grain and Silage), Popcorn,  | 15          | Succulent Peas and Beans             | 33          |
| Field Corn Grown for Seed (PRE & PPI)    |             | Tobacco                              | 34          |
| Field Corn (Grain and Silage), Popcorn,  | 15-16       | Tuberous and Corn Vegetables         | 35          |
| Field Corn Grown for Seed (Foliar Use)   |             | Ornamentals and Trees                | 35-40       |
| Sweet Corn (Grain and Silage),           | 17          | (including Field and Container Grown |             |
| Sweet Corn Grown for Seed (At Plant Use) |             | Nursery Stock, Christmas Trees,      |             |
| Sweet Corn (Grain and Silage),           | 18-19       | Interiorscapes and Plantscapes,      |             |
| Sweet Corn Grown for Seed (Foliar Use)   |             | Lawns, Trees and Shrubs,             |             |
| Cotton                                   | 19-20       | and on Golf Courses and Sod Farms)   |             |

## FOOD CROP USE INSTRUCTIONS

### ARTICHOKE

Apply as directed at a rate of 6.4 fl. oz. (0.1 lb. ai) per acre.

| PEST                 | APPLICATION INSTRUCTIONS   |
|----------------------|--|
| Artichoke Plume Moth | Apply when pest population reaches damaging threshold and repeat as necessary to maintain control, but not apply more often than 15 day intervals.   |
| Cribrate Weevil      | <p><b>Application by ground:</b> Apply a full cover spray in a minimum of 75 gallons of finished spray per acre.</p> <p><b>Application by air:</b> Apply specified rate in a minimum of 10 gallons per acre.</p> |

### RESTRICTIONS:

- Do not exceed 0.5 lb. ai per acre per season.
- A 5-day preharvest interval must be observed.

## BRASSICA CROPS

Apply as directed using the rates in the table below.

| CROP   | PEST/RATE   | APPLICATION INSTRUCTIONS   |
|--|---|--|
| Head and Stem Brassica<br>Vegetables including:<br>Broccoli<br>Chinese Broccoli (gailon, White flowering broccoli)<br>Brussels sprouts<br>Cauliflower<br>Cavalo broccoli<br>Kohlrabi<br>Cabbage<br>Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy)   | <p style="text-align: center;"><b>PESTS:</b></p> Cutworms<br>Corn Earworm<br>Tobacco Budworm<br>Saltmarsh Caterpillar<br>Leafhoppers<br>Flea Beetles<br>Imported Cabbageworm<br>Cucumber Beetles<br>Whitefly<br>Armyworms<br>Loopers<br>Stink Bugs<br>Crickets<br>Ground Beetles<br>Thrips<br>Wireworm (adults)<br>Diamondback Moth <p style="text-align: center;"><b>RATE:</b></p> 2.1 to 6.4 fl. oz./acre<br>(0.033 to 0.1 lb. ai/acre) | <p>Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.</p> <p>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.</p> <p>Thorough coverage is essential to achieve control.</p> |
| <p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 0.5 lb. active ingredient (1 quart) per acre per season.</li> <li>• Do not make more than 5 applications after bloom.</li> <li>• Do not make applications less than 7 days apart.</li> <li>• Do not apply within 7 days of harvest.</li> </ul> | Banks Grass Mite<br>Twospotted Spider Mite<br>Carmine Mite<br>Pacific Spider Mite<br>Lygus spp. <p style="text-align: center;"><b>RATE:</b></p> 5.12 to 6.4 fl. oz./acre<br>(0.08 to 0.1 lb. ai/acre)   |  |

## CANE BERRIES

Apply as directed using the rates in the table below.

| CROP  | PEST/RATE  | APPLICATION INSTRUCTIONS  |
|---|--|---|
| <p>Caneberries including:<br/>Blackberries<br/>Bingleberries<br/>Dewberries<br/>Lowberries<br/>Marionberries<br/>Olallieberries<br/>Youngberries<br/>Loganberries<br/>Raspberries</p> | <p><b>PESTS:</b><br/>Leafrollers<br/>Orange Tortrix<br/>Root Weevils</p> <p><b>RATE:</b><br/>3.2 to 6.4 fl. oz./acre<br/>(0.05 to 0.1 lb. ai/acre)</p> | <p>Apply by air or ground equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons per acre by air and 50 gallons per acre by ground).</p> <p>One application may be made pre-bloom and a second application may be made post bloom.</p>  |
|   | <p><b>PEST:</b><br/>Spider Mites<br/>Raspberry Crown Borer</p> <p><b>RATE:</b><br/>6.4 fl. oz./acre<br/>(0.1 lb. ai/acre)</p>                          | <p>For Crown Borer, apply 0.1 lb. ai per acre post-harvest (fall) or pre-bloom (spring) as a drench application directed at the crown of plants in a minimum of 200 gallons of water per acre. Greater efficacy is observed at higher water gallonages (up to 400 gallons) or in an application prior to a significant rainfall event. Do not make both pre-bloom foliar and pre-bloom drench applications.</p> |
| <p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"><li>• Do not apply within 3 days of harvest.</li><li>• Do not exceed 0.2 lb ai per acre per season.</li></ul>           |  |   |

**CANOLA, CRAMBE, RAPESEED**

Apply as directed at a rate of 2.1 to 2.6 fl. oz. (0.033 to 0.04 lb. ai) per acre.

| <b>PEST</b>   | <b>APPLICATION INSTRUCTIONS</b>  |
|---|--|
| Aphids<br>Armyworms<br>Cutworms<br>Diamondback Moth<br>Loopers<br>Other Lepidopterous Larvae<br>Lygus Bugs<br>Flea Beetle<br>Flea Hopper<br>Grasshoppers<br>Plant Bugs<br>Stink Bugs<br>Seedpod Weevil<br>Thrips<br>Whitefly  | Apply in a minimum of 2 gallons of finished spray per acre by air, or in a minimum of 10 gallons per acre with ground equipment.<br><br>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.<br><br>Thorough coverage is essential to achieve control. |
| <b>RESTRICTIONS:</b> <ul style="list-style-type: none"><li>• Do not apply more than 0.08 lb. active ingredient (5.12 ounces) per acre per season.</li><li>• Do not make applications less than 14 days apart.</li><li>• Do not apply within 35 days of harvest.</li></ul> |  |

## CILANTRO, CORIANDER

Apply as directed using the rates in the table below.

| PEST   | RATE  | APPLICATION INSTRUCTIONS   |
|--|---|--|
| Aphids<br>Beet Armyworm<br>Cabbage Looper<br>Cutworm<br>Flea Beetle<br>Grasshoppers<br>Leafminer<br>Saltmarsh caterpillar<br>Spotted Cucumber Beetle<br>Thrips<br>Whitefly   | 2.1 to 6.4 fl. oz. (0.033 to 0.1 lb. ai) per acre | Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft. |
| Two Spotted Spider Mite  | 5.12 to 6.4 fl. oz. (0.08 to 0.1 lb. ai) per acre |  |
| <b>RESTRICTIONS:</b> <ul style="list-style-type: none"><li>• Do not make applications less than 7 days apart.</li><li>• Do not apply more than 0.5 pound active ingredient per acre per season.</li><li>• Do not apply within 3 days of harvest.</li></ul> |   |  |

## CITRUS

The use of REVEAL ENDURX protects citrus tree roots from **Diaprepes** and other citrus root weevil feeding by creating a barrier. As citrus root weevil eggs hatch, the newly hatched larvae (neonates) fall to the soil surface beneath the tree and come into contact with REVEAL ENDURX as they attempt to burrow into the root zone. Disturbance of the soil beneath the tree should be minimized.

Timing of application is very important. Peak emergence of **Diaprepes** adults varies by citrus growing region, and environmental factors such as soil moisture can affect emergence. Usually, two peaks occur for **Diaprepes**, first in the spring then late summer or early fall. **Southern Blue-Green and Blue-Green Citrus Weevils** and **Fuller Rose Beetle** usually have a single emergence peak in the spring. Brown and Little Leaf Notchers usually have three emergence peaks, spring, summer and fall. Since emergence varies by region and season, the best way to time application is to observe the adults. By trapping adults when they are most active (in the morning or and late afternoon) during the spring and summer emergence periods an estimation of numbers can be obtained. Eggs are laid 8 to 10 weeks following the adult emergence from the soil; larvae invasion into the soil will begin 2 to 3 weeks following adult emergence. REVEAL ENDURX must be applied prior to the dropping of the neonates. Consult local university extension personnel for current information to protect citrus trees from **Citrus Root Weevils** and other pests.

**CITRUS cont'd**

Apply as directed using the rates in the table below.

| PEST   | RATE   | APPLICATION INSTRUCTIONS  |
|--|--|---|
| <p>Diaprepes Root Weevil<br/>(Diaprepes abbrevia tus),<br/>Southern Blue Green<br/>Citrus Root Weevil<br/>(Pachnaeus litus),<br/>Blue green Citrus Root<br/>Weevil<br/>(Pachnaeus opalus),<br/>Brown Leaf Notcher<br/>(Epicaerus mexica-<br/>nus),<br/>Little Leaf Notcher<br/>(Artipus floridanus)</p>  | <p>16 to 32 fl. oz.<br/>(0.25 to 0.5<br/>lb. ai)<br/>per acre</p>  | <p>Apply REVEAL ENDURX by ground equipment to bare soil beneath citrus trees. REVEAL ENDURX must be uniformly applied from the trunk to the drip line of the tree, apply in a minimum of 40 gallons of dilute spray per acre. Greater spray volume should insure greater uniformity of coverage. A pre- and post-application irrigation may aid in the uniformity of coverage as well.</p> <p>Apply to individual citrus resets, when not in solid planted rows, using hand-gun or shielded sprayer.</p> <p>Peak emergence of Diaprepes Root Weevil generally occurs in the spring. Depending on weather conditions, a minor emergence may also occur in the fall.</p> <p>If the citrus grove to be treated is in an area where weather conditions are conducive to primary emergence occurring in the spring, 32 fl. oz. formulated product should be used to obtain best results.</p> |
| <p>Fire Ant<br/>(Solenopsis spp.),<br/>Asian Cockroach<br/>(Blattella asahinae)</p>  | <p>6.4 to 16 fl. oz.<br/>(0.1 to 0.25<br/>lb. ai)<br/>per acre</p> | <p>If the citrus grove to be treated is in an area where weather conditions will promote more than one peak pest emergence, 16 fl. oz. formulated product can be applied early season and 16 fl. oz. formulated product can be applied later in the season.</p> <p>Follow spray drift precautions on this label.</p>  |
| <p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>• Do not apply through irrigation systems.</li> <li>• Do not allow any application of REVEAL ENDURX to contact fruit or foliage.</li> <li>• Do not apply a total of 32 fl. oz. of REVEAL ENDURX (0.5 lb. ai) per acre per year.</li> <li>• Apply the specified dosage in a minimum of 40 gallons of finished spray per acre.</li> <li>• Ground application only. Do not apply by air.</li> <li>• Do not apply within 1 day of harvest.</li> </ul> |  |   |

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED  
(AT PLANT USE)**

Apply as directed using the rates in the table below.

| PEST   | RATE  | APPLICATION INSTRUCTIONS   |           |           |  |
|--|---|--|-----------|-----------|--|
| Corn Rootworm Larvae<br>(Northern, Southern,<br>Western)<br>Army Cutworm<br>Cutworm Species<br>Grubs<br>Seed Corn Beetle<br>Seed Corn Maggot<br>True Armyworm<br>or<br>Armyworm Species<br>Wireworms   | 0.30 fl. oz. (.0046 lb. ai) per 1,000 linear feet of row                  | Apply as a 5- to 7-inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe in front of the press wheel centered over the row. Use the table below to determine the REVEAL ENDURX needs per acre. Apply in a minimum of 3 gallons of finished spray per acre. |           |           |  |
|  | 0.15 to 0.30 fl. oz. (.0023 to .0046 lb. ai) per 1,000 linear feet of row |  |           |           |  |
| <p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>• Do not apply to soil where there is greater than 30% cover of crop residue remaining.</li> <li>• Do not apply within 30 days of harvest.</li> <li>• Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment.</li> <li>• Do not apply more than 0.1 pound active per acre per season as an at-plant application.</li> </ul> |   |  |           |           |  |
| <b>Row spacings (inches)</b>   | <b>40</b>   | <b>38</b>  | <b>36</b> | <b>30</b> |  |
| REVEAL ENDURX (pounds ai per acre)   | 0.060   | 0.064  | 0.069     | 0.080     |  |
| REVEAL ENDURX (formulated product ounces per acre)   | 3.9   | 4.1  | 4.4       | 5.12      |  |

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (PRE- EMERGENT & PRE-PLANT INCORPORATED)**

Apply as directed using the rates in the table below.

| PEST  | RATE   | APPLICATION INSTRUCTIONS   |
|---|--|--|
| Black Cutworm<br>White Grub<br>Wireworm<br>Seedcorn Maggot<br>Armyworm spp.<br>Stalkborer | 3 to 4 fl. oz.<br>(0.047 to 0.062<br>lb. ai)<br>per acre<br>Pre-plant<br>Incorporated<br>(PPI) | The 3-4 oz/A rate must be applied as PPI and can be tank mixed and applied with PPI herbicides. Incorporation of REVEAL ENDURX should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth. |
| Black Cutworm<br>Armyworm spp.<br>Stalkborer  | 2.56 fl. oz. (0.040<br>lb. ai) per acre<br>Pre-emergence<br>(PRE)                              | The 2.56 oz/A rate may be applied PRE and can be tank mixed and applied with PRE herbicides.   |

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (FOLIAR USE)**

Apply as directed using the rates in the table below.

| PEST   | RATE   | APPLICATION INSTRUCTIONS  |
|--|--|---|
| Aphids<br>Army Cutworm<br>Beet Armyworm<br>Cereal Leaf Beetle<br>Chinch Bug<br>Common Stalk Borer<br>Corn Earworm<br>Corn Rootworm Adult<br>Cucumber Beetle Adults<br>Cutworm Species<br>European Corn Borer<br>Fall Armyworm<br>Flea Beetle<br>Grasshoppers<br>Greenbug<br>Japanese Beetle Adult<br>Sap Beetle<br>Southern Armyworm<br>Southern Corn Leaf Beetle<br>Southwestern Corn Borer<br>Stinkbugs<br>Tarnished Plant Bug | 2.1 to 6.4<br>fl. oz.<br>(0.033 to 0.10<br>lb. ai)<br>per acre | <p>Apply in a minimum of 2 to 5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment. To improve control by aircraft, use 5 gallons of finished spray per acre particularly when initial populations are heavier than normal. When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.</p> <p>Thorough coverage is essential to achieve control.</p> <p><b>To Control Ear-Attacking Pests:</b> Apply REVEAL ENDURX just before silking and repeat as necessary to maintain control but do not exceed maximum application rates and reapplication intervals listed elsewhere in this section.</p> <p><b>Southwestern Corn Borer, European Corn Borer:</b> Make application for corn borer control with initial application at or shortly before egg hatch.</p> <p><b>For Control Of Other Insect Pests:</b> Apply when pests first appear and repeat as necessary but do not exceed maximum application rates and reapplication intervals listed elsewhere in this section.</p> |

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (FOLIAR USE) *cont'd***

| <b>PEST</b>  | <b>RATE</b>  | <b>APPLICATION INSTRUCTIONS</b>   |
|--|--|---|
| True Armyworm<br>or<br>Armyworm Species<br>Webworms<br>Western Bean Cut-<br>worm<br>Yellowstriped Army-<br>worm  | 2.1 to 6.4<br>fl. oz.<br>(0.033 to 0.10<br>lb. ai)<br>per acre |   |
| Banks Grass Mite<br>Carmine Mite<br>Twospotted Spider Mite   | 5.12 to 6.4 fl.<br>oz. (0.08-0.10<br>lb. ai)<br>per acre       | <p>Apply for <b>Banks Grass Mite</b> control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.</p> <p>For <b>Twospotted Spider Mite</b> and <b>Carmine Mite</b> control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy. Higher rates will be necessary for heavier initial populations and corn under heat or drought stress. Field experience with dimethoate at 0.5 lb. active per acre in tank mixture has demonstrated good control under these conditions.</p> <p><b>For Mite Control In Texas, New Mexico, Oklahoma, and Arizona:</b> Apply in a minimum of 5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment.</p> |
| <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 0.3 pound active ingredient per acre per season including pre and ppi, at-plant, plus foliar applications.</li> <li>• Do not apply within 30 days of harvest.</li> <li>• Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.</li> <li>• Use of ultra low volume (ULV) application on corn is prohibited.</li> <li>• Do not make aerial or ground applications to corn if heavy rainfall is imminent.</li> <li>• Use of REVEAL ENDURX on corn is prohibited in all coastal counties.</li> </ul> |  |   |

## SWEET CORN (GRAIN AND SILAGE), SWEET CORN GROWN FOR SEED (AT PLANT USE)

Apply as directed using the rates in the table below.

| PEST   | RATE  | APPLICATION INSTRUCTIONS  |           |           |  |
|--|---|---|-----------|-----------|--|
| Corn Rootworm Larvae (Northern, Southern, Western)   | 0.30 fl. oz. (0.0046 lb. ai) per 1,000 linear feet of row                   | Apply as a 5- to 7-inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the REVEAL ENDURX needs per acre. Apply in a minimum of 3 gallons of finished spray per acre.   |           |           |  |
| Army Cutworm<br>Cutworm Species<br>Grubs<br>Seed Corn Beetle<br>Seed Corn Maggot<br>True Armyworm<br>or<br>Armyworm species<br>Wireworms   | 0.15 to 0.30 fl. oz. (0.0023 to 0.0046 lb. ai) per 1,000 linear feet of row | <p>Mix REVEAL ENDURX with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of REVEAL ENDURX, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.</p> <p>Applications of REVEAL ENDURX alone or in recommended tank mixtures, in conjunction with in-furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of REVEAL ENDURX and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.</p> |           |           |  |
| <b>RESTRICTIONS:</b>   |   |   |           |           |  |
| <ul style="list-style-type: none"> <li>• Do not apply to soil where there is greater than 30% cover of crop residue remaining.</li> <li>• Do not apply within 30 days of harvest.</li> <li>• Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment.</li> <li>• Do not apply more than 0.1 pound active per acre per season as an at-plant application.</li> </ul> |   |   |           |           |  |
| <b>Row spacings (inches)</b>   | <b>40</b>   | <b>38</b>   | <b>36</b> | <b>30</b> |  |
| REVEAL ENDURX (pounds per acre)  | 0.060   | 0.064   | 0.069     | 0.080     |  |
| REVEAL ENDURX(formulated product ounces per acre)  | 3.9   | 4.1   | 4.4       | 5.12      |  |

**SWEET CORN (GRAIN AND SILAGE), SWEET CORN GROWN FOR SEED (FOLIAR USE)**

Apply as directed using the rates in the table below.

| PEST   | RATE   | APPLICATION INSTRUCTIONS  |
|--|--|---|
| <p>Aphids<br/> Army Cutworm<br/> Beet Armyworm<br/> Cereal Leaf Beetle<br/> Chinch Bug<br/> Common Stalk Borer<br/> Corn Earworm<br/> Corn Rootworm Adults<br/> Cucumber Beetle Adult<br/> Cutworm Species<br/> European Corn Borer<br/> Fall Armyworm<br/> Flea Beetle<br/> Grasshoppers<br/> Greenbugs<br/> Japanese Beetle Adult<br/> Sap Beetle<br/> Southern Armyworm<br/> Southern Corn Leaf Beetle<br/> Southwestern Corn Borer<br/> Stinkbugs<br/> Tarnished Plant Bug<br/> True Armyworm<br/> or Armyworm Species<br/> Webworms<br/> Western Bean Cutworm<br/> Yellowstriped Armyworm</p> | <p>2.1 to 6.4 fl. oz.<br/> (0.033-0.10 lb. ai)<br/> per acre</p> | <p>Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.</p> <p>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.</p> <p>Thorough coverage is essential to achieve control.</p> <p><b>To Control Ear-Attacking Pests:</b> Apply REVEAL ENDURX when silking begins and repeat as necessary to maintain control but do not exceed maximum application rates and reapplication intervals listed elsewhere in this section.</p> <p><b>Southwestern Corn Borer, European Corn Borer:</b> Make 2 applications for corn borer with the initial application at or shortly before egg hatch.</p> <p><b>For Control Of Other Insect Pests:</b> Apply when pests first appear and repeat as necessary but do not exceed maximum application rates and reapplication intervals listed elsewhere in this section.</p> |
| <p>Banks Grass Mite<br/> Carmine Mite<br/> Twospotted Spider Mite</p>  | <p>5.12 to 6.4 fl. oz.<br/> (0.08-0.10 lb. ai)<br/> per acre</p> | <p>Apply for <b>Banks Grass Mite</b> control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.</p> <p>For <b>Twospotted Spider Mite</b> and <b>Carmine Mite</b> control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.</p> <p>Higher rates will be necessary for heavier initial populations and corn under heat or drought stress.</p>   |

**SWEET CORN (GRAIN AND SILAGE), SWEET CORN GROWN FOR SEED (FOLIAR USE) *cont'd***

**RESTRICTIONS:**

- Do not apply more than 0.2 pound active ingredient (12.8 ounces formulated product) per acre per season.
- Do not apply within one day of harvest.
- Do not graze livestock in treated areas or cut treated crops for feed within 1 day of last application.
- Use of ultra low volume (ULV) application on corn is prohibited.
- Do not make aerial or ground applications to corn if heavy rainfall is imminent.
- Use of REVEAL ENDURX on corn is prohibited in all coastal counties.

**COTTON**

Apply as directed using the rates in the table below.

| PEST  | RATE  | APPLICATION INSTRUCTIONS  |
|---|---|---|
| European Corn Borer<br>Soybean (Banded)<br>Thrips<br>Tobacco Thrips   | 1.3-6.4 fl. oz./acre<br>(0.02-0.10 lb. ai/<br>acre) | REVEAL ENDURX may be applied in water or refined vegetable oil (soybean/cottonseed).  |
| Boll Weevil<br>Bollworm<br>Cabbage Looper<br>Cotton Aphid<br>Cotton Fleahopper<br>Cotton Leafperforator<br>Cutworms<br>Fall Armyworm<br>Plant Bugs<br>Saltmarsh Caterpillar<br>Southern Garden<br>Leafhopper<br>Stink Bugs<br>Tobacco Budworm<br>Whitefly<br>Yellowstriped Armyworm | 2.6-6.4 fl. oz./acre<br>(0.04-0.10 lb. ai/<br>acre) | <p><b>Application in Water:</b> Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray.</p> <p><b>ULV Application:</b> Apply the recommended rate of REVEAL ENDURX in refined vegetable oil in a minimum of 1 quart of finished spray per acre with aircraft calibrated to give adequate coverage.</p> <p><b>To Control Boll Weevil:</b> Apply REVEAL ENDURX at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels.</p> <p><b>To Control Mites and Aphids:</b> Apply when pests first appear. Repeat as necessary to maintain control but do not exceed maximum application rates and reapplication intervals listed elsewhere in this section. Higher rates will be required once a damaging threshold is established.</p> |
| Beet Armyworm<br>Carmine Spider Mite<br><i>Lygus</i> spp.<br>Pink Bollworm<br>Twospotted Spider Mite  | 3.8-6.4 fl. oz./acre<br>(0.06-0.10 lb. ai/<br>acre) |   |

**COTTON cont'd****RESTRICTIONS:**

- Do not apply more than 0.5 pound active ingredient per acre per season.
- Do not apply within 14 days of harvest.
- Do not graze livestock in treated areas or cut treated crops for feed.
- Do not make more than 10 synthetic pyrethroid applications (of one product or combinations or products) to a cotton crop in one growing season.

**CUCURBITS**

Apply as directed using the rates in the table below.

| CROP   | PEST/RATE  | APPLICATION INSTRUCTIONS   |
|--|--|--|
| <p>Chayote (fruit)<br/>           Chinese waxgourd (Chinese preserving melon)<br/>           Citron melon<br/>           Cucumber<br/>           Gherkin<br/>           Gourd, edible (includes hyotan, cucuzza); (<i>Luffa</i> spp.) (includes hechima, Chinese okra), (<i>Momordica</i> spp.) (includes balsam apple, balsam pear, bitter melon, Chinese cucumber),<br/>           Muskmelon (hybrids and/or cultivars or <i>Cucumis melo</i>) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon),<br/>           Pumpkin (<i>Cucurbita</i> spp.), Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini).<br/>           Squash, winter (includes butternut squash, calabaza, hubbard squash (<i>C. mixta</i>; <i>C. pepo</i>) (includes acorn squash, spaghetti squash), Watermelon (includes hybrids and/or varieties of <i>Citrullus</i> spp.).</p> | <p><b>PESTS:</b><br/>           Aphids<br/>           Armyworms<br/>           Cabbage Looper<br/>           Corn Earworm<br/>           Cucumber Beetles<br/>           Cutworms<br/>           Grasshoppers<br/>           Leafhoppers<br/>           Melonworms<br/>           Pickleworms<br/>           Plant Bugs<br/>           Rindworms<br/>           Squash Bugs<br/>           Squash Vine Borer<br/>           Stink Bugs<br/>           Tobacco Budworm</p> <p><b>RATE:</b><br/>           2.6 to 6.4 fl. oz. (0.04 to 0.1 lb. ai) per acre</p> <p><b>PESTS:</b><br/>           Whitefly<br/>           Banks Grass Mite<br/>           Twospotted spider Mite<br/>           Carmine Mite<br/> <i>Lygus</i> spp.</p> <p><b>RATE:</b><br/>           5.12 to 6.4 fl. oz. (0.08 to 0.1 lb. ai) per acre</p> | <p>Apply in a minimum of 5 gallons of finished spray per acre by air or in a minimum of 20 gallons per acre with ground equipment.</p> <p>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.</p> <p>Thorough coverage is essential to achieve control.</p> |

**CUCURBITS cont'd****RESTRICTIONS:**

- Do not apply more than 0.3 lb. active ingredient (19.2 ounces formulated product) per acre per season.
- Do not make more than two applications after bloom.
- Do not make applications less than 7 days apart.
- Do not apply within 3 days of harvest.

**DRIED BEANS AND PEAS**

Apply as directed using the rates in the table below.

| CROP  | PEST   | RATE   | APPLICATION INSTRUCTIONS  |
|---|--|--|---|
| <b>Dried cultivars of Beans</b><br><i>(Lupinus )</i><br><b>Beans</b> ( <i>Phaseolus</i> )<br>Field bean<br>Kidney bean<br>Lima bean (dry)<br>Navy bean<br>Pinto bean<br>Tepary bean<br><b>Bean</b> ( <i>Vigna</i> )<br>Adzuk bean<br>Blackeyed pea<br>Catjang<br>Cowpea<br>Crowder pea<br>Moth bean<br>Mung bean<br>Rice bean<br>Southern pea<br>Urd bean<br>Broad bean (dry)<br>Chickpea<br>Guar<br>Lablab bean<br>Lentil<br><b>Peas</b> ( <i>Pisicum</i> )<br>Field pea<br>Pigeon pea | Aster Leafhopper<br>Flea Beetle<br>Grasshoppers<br>Leafhoppers   | 1.6 to 6.4 fl. oz. (0.025 to 0.10 lb. ai) per acre | Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.<br><br>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.<br><br>Thorough coverage is essential to achieve control. |
|   | Aphids<br>Beet Armyworm<br>Fall Armyworm<br>Southern Armyworm<br>Yellowstriped Armyworm<br>Bean Leaf Beetle<br>Cucumber Beetle<br>Japanese Beetle (Adult)<br>Sap Beetle<br>Plant Bugs<br>Stink Bugs<br>Tarnished Plant Bug<br>Alfalfa Caterpillar<br>Cloverworm<br>European Corn Borer<br>Cutworms<br>Western Bean Cutworm<br>Corn Earworm<br>Loopers<br>Corn Rootworm (Adult)<br>Thrips<br>Webworms<br>Pea Weevil<br>Pea Leaf Weevil<br>Whitefly<br>Imported cabbageworm<br>Saltmarsh caterpillar<br>Tobacco budworm<br>Leafminer | 2.1 to 6.4 fl. oz. (0.033 to 0.10 lb. ai) per acre |   |
|   | Banks Grass Mite<br>Twospotted Spider Mite<br>Carmine Mite<br><i>Lygus spp</i>   | 5.12 to 6.4 fl. oz. (0.08 to 0.10 lb. ai) per acre |   |

**DRIED BEANS AND PEAS *cont'd*****RESTRICTIONS:**

- Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated product) to peas, or 0.3 lb. active ingredient (19.2 ounces formulated product) to beans per acre per season.
- Do not apply within 14 days of harvest.
- Do not make applications less than 7 days apart.

**FRUITING VEGETABLES**

Apply as directed using the rates in the table below.

| CROP  | PEST  | RATE   | APPLICATION INSTRUCTIONS  |
|---|---|--|---|
| Eggplant<br>Pepper<br>(Bell & Non-Bell)<br>Groundcherry<br>Pepino | Armyworms including Beet<br>Armyworm, Fall Armyworm,<br>Southern Yellowstriped<br>Armyworm<br>Cabbage Looper<br>Colorado Potato Beetle<br>Corn Earworm<br>Cucumber Beetle<br>Cutworms<br>European Corn Borer<br>Flea Beetle<br>Leafminers<br>Loopers<br>Pepper weevil<br>Plant Bugs<br>Stink Bugs<br>Thrips<br>Tomato Hornworm<br>Tomato Pinworm<br>Vegetable Leafminer<br>Whitefly | 2.1 to 6.4 fl. oz.<br>(0.033 to 0.10 lb. ai)<br>per acre | Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.<br><br>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.<br><br>Thorough coverage is essential to achieve control. |
|   | Banks Grass Mite<br>Broad Mite<br>Carmine Mite<br>Lygus spp<br>Pacific Spider Mite<br>Twospotted Spider Mite  | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.10 lb. ai)<br>per acre |   |

**RESTRICTIONS:**

- Do not make applications less than 7 days apart.
- Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated product) per acre per season.
- Do not apply within 7 days of harvest.

**FRUITING VEGETABLES *cont'd***

| <b>CROP</b>         | <b>PEST</b>   | <b>RATE</b>  | <b>APPLICATION INSTRUCTIONS</b>  |
|---------------------|---|--|--|
| Tomato<br>Tomatillo | Aphids<br>Armyworms including Beet<br>Armyworm, Fall Armyworm,<br>Southern Armyworm,<br>Yellowstriped Armyworm<br>Bean Leaf Beetle<br>Cabbageworm<br>Carmine Mite<br>Cloverworm<br>Corn Earworm<br>Cucumber Beetle<br>Cutworms<br>Diamondback Moth<br>European Corn Borer<br>Flea Beetles<br>Flea Hopper<br>Grasshopper<br>Japanese Beetle (Adult)<br>Leafhoppers<br>Loopers<br>Lygus spp.<br>Melonworm<br>Pea Weevil<br>Pea Leaf Weevil<br>Pickleworm<br>Plant Bug<br>Rindworm<br>Salt Marsh Caterpillar<br>Sap Beetle<br>Seedpod Weevil<br>Squash Bugs<br>Stink Bug species<br>Tobacco Budworm<br>Tarnished Plant Bug<br>Thrips<br>Whitefly | 2.1 to 5.2 fl. oz.<br>(0.033 to 0.08 lb. ai)<br>per acre | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment.<br><br>Thorough coverage is essential to achieve control. |
|                     | Twospotted Spider Mite  | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.10 lb. ai)<br>per acre |  |

**FRUITING VEGETABLES cont'd****RESTRICTIONS:**

- A maximum of 4 applications may be applied per season.
- Do not apply within 1 day of harvest.

**GRAPES**

Apply as directed using the rates in the table below.

| PEST   | RATE  | APPLICATION INSTRUCTIONS   |
|--|---|--|
| Cutworms<br>Eastern Grape Leafhopper<br>Grape Berry Moth<br>Japanese Beetles Adults<br>Variegated Leafhopper<br>Western Grape Leafhopper                       | 3.2 to 6.4 fl. oz.<br>(0.05 to 0.10 lb. ai)<br>per acre | Apply in a minimum of 10 gallons of finished spray by air or in a minimum of 25 gallons of finished spray with ground equipment.<br><br>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray. |
| Black Vine Weevil<br>Glassywinged Sharpshooter<br>Twospotted Spider Mite   | 6.4 fl. oz.<br>(0.10 lb. ai)<br>per acre                | Thorough coverage is essential to achieve control.<br><br>When pest pressure is moderate to severe, use higher rate.   |
| <b>RESTRICTIONS:</b>   |   |  |
| <ul style="list-style-type: none"> <li>• Do not apply more than 0.10 lb. ai per acre per season.</li> <li>• Do not apply within 30 days of harvest.</li> </ul> |   |  |

**HOPS**

Apply as directed using the rates in the table below.

| PEST  | RATE   | APPLICATION INSTRUCTIONS   |
|---|--|--|
| Aphids<br>Armyworms<br>Cutworms<br>Leafrollers<br>Loopers | 3.8 to 6.4 fl. oz.<br>(0.06 to 0.1 lb. ai)<br>per acre | <b>Application by ground:</b> For best results, full coverage is essential. Early season - recommend 100-150 gallons of spray per acre. Late season -recommend 200-250 gallons of spray per acre.  |
| Root Weevils  | 3.2 to 6.4 fl. oz. (0.05 to 0.1 lb. ai) per acre       | For <b>Root Weevil</b> control, make a directed spray to the base of the plant. Spray up the vine 3 feet and the soil surface 1.5 to 2 feet on either side of the plant.   |
| Twospotted Spider Mite                                    | 6.4 fl. oz<br>(0.1 lb. ai) per acre                    | <b>Application by air for late season control of Twospotted Spider Mites:</b> Apply no less than 6.4 oz (0.1 lb ai) per application in a minimum of 10 gallons per acre.<br><br>Use of ultra low volume (ULV) application on hops is prohibited. |

**HOPS cont'd****RESTRICTIONS:**

- Do not exceed 0.1 lb. ai per acre per application.
- Do not exceed 0.3 lb. ai per acre per season.
- A spray interval of 21 days between applications must be maintained.
- A 14-day preharvest interval must be observed.

**LEAFY BRASSICAS\*, TURNIP GREENS\***

Apply as directed using the rates in the table below.

| CROP  | PEST  | RATE  | APPLICATION INSTRUCTIONS   |
|---|---|---|--|
| Broccoli Raab<br>Bok Choy<br>Collards<br>Kale<br>Mizuna<br>Mustard Greens<br>Mustard Spinach<br>Rape Greens | Cutworms<br>Corn Earworm<br>Tobacco Budworm<br>Saltmarsh Caterpillar<br>Leafhoppers<br>Flea Beetles<br>Imported Cabbage-worm<br>Cucumber Beetles<br>Aphids<br>Whitefly<br>Armyworms<br>Loopers<br>Stink Bugs<br>Crickets<br>Ground Beetles<br>Thrips<br>Wireworm (Adults)<br>Diamondback Moth<br>Japanese Beetles (Adult)<br>Grasshoppers | 2.1 to 6.4 fl. oz./acre<br>(0.033 to 0.1 lb. ai/acre) | Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.<br><br>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.<br><br>Thorough coverage is essential to achieve control.<br><br>*See resistance Management Statement under Directions For Use section |
|   | Banks Grass Mite<br>Twospotted Spider Mite<br>Carmine Mite<br>Pacific Spider Mite<br><i>Lygus</i> spp.  | 5.12 to 6.4 fl. oz./acre<br>(0.08 to 0.1 lb. ai/acre) |  |

**LEAFY BRASSICAS\*, TURNIP GREENS\* *cont'd*****RESTRICTIONS:**

- Do not apply more than 0.4 lb. active ingredient per acre per season.
- Do not make applications less than 7 days apart.
- Do not apply within 7 days of harvest.

**LETTUCE, HEAD**

Apply as directed using the rates in the table below.

| PEST  | RATE   | APPLICATION INSTRUCTIONS   |
|---|--|--|
| Aphids<br>Armyworms<br>Corn Earworm<br>Cucumber Beetles<br>Cutworms<br>Diamondback Moth<br>Flea Beetle<br>Imported Cabbageworm<br>Leafhoppers<br>Loopers<br>Salt marsh Caterpillar<br>Stink Bug spp.<br>Tobacco Budworm<br>Whitefly | 2.1 to 6.4 fl. oz.<br>(0.033 to 0.10 lb.<br>ai) per acre | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.<br><br>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.<br><br>Thorough coverage is essential to achieve control. |
| <i>Lygus</i> spp.<br>Carmine Mite<br>Twospotted Spider Mite   | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.10 lb.<br>ai) per acre |  |

**RESTRICTIONS:**

- Do not make applications less than 7 days apart.
- A maximum of 0.5 lb. active ingredient may be applied per acre per season.
- Do not apply within 7 days of harvest.

**MAYHAW**

Apply as directed using the rates in the table below.

| PEST          | RATE   | APPLICATION INSTRUCTIONS   |
|---------------|--|--|
| Plum Curculio | 5.12 to 6.4 fl. oz. (0.08 to 0.10 lb. ai) per acre | Apply as a foliar treatment using at least of 28 gallons per acre. |

**RESTRICTIONS:**

- Do not make applications less than 7 days apart.
- Do not apply more than 0.2 pound active ingredient per acre per season.
- Do not apply within 30 days of harvest.

**OKRA**

Apply as directed using the rates in the table below.

| PEST  | RATE  | APPLICATION INSTRUCTIONS   |
|---|---|--|
| Aphids<br>Armyworms<br>Corn earworm<br>Cucumber Beetles<br>Cutworms<br>European Corn Borer<br>Flea Beetles<br>Japanese Beetle<br>(Adult)<br>Leafminers<br>Loopers<br>Stink bugs<br>Thrips<br>Whitefly | 2.1 to 6.4 fl. oz. (0.033 to 0.1 lb. ai) per acre | Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft. |
| <i>Lygus</i> spp.<br>Broad Mite<br>Carmine Mite<br>Two Spotted Spider Mite  | 5.12 to 6.4 fl. oz. (0.08 to 0.1 lb. ai) per acre |  |

**RESTRICTIONS:**

- Do not make applications less than 7 days apart.
- Do not apply more than 0.2 pound active ingredient per acre per season.
- Do not apply within 7 days of harvest.

**PEANUT**

Apply as directed using the rates in the table below.

| PEST  | RATE  | APPLICATION INSTRUCTIONS   |
|---|---|--|
| Beet Armyworms<br>Corn earworm<br>Cucumber Beetles<br>Cutworm species<br>Fall Armyworm<br>Grasshoppers<br>Green cloverworm<br>Leafhoppers<br>Lesser Cornstalk Borer<br>Loopers<br>Rednecked Peanut Worm<br>Southern Armyworm  | 2.1 to 6.4 fl. oz.<br>(0.033 to 0.1 lb. ai)<br>per acre | Apply foliar treatments in at least 10 gallons per acre at the rate of 6.4 fl. oz. (0.1 lb. active ingredient) per acre at a minimum of 14 days intervals. |
| Southern Corn Rootworm<br>Stink Bugs<br>Threecornered Alfalfa Hopper<br>Velvetbean Caterpillar<br>Yellowstriped Armyworm  | 2.1 to 6.4 fl. oz.<br>(0.033 to 0.1 lb. ai)<br>per acre | Apply foliar treatments in at least 10 gallons per acre at the rate of 6.4 fl. oz. (0.1 lb. active ingredient) per acre at a minimum of 14 days intervals. |
| Aphids<br>Spider Mites<br>Thrips<br>Whitefly  | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.1 lb. ai)<br>per acre |  |
| <b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>• Do not feed green immature plants and peanut hay to livestock.</li> <li>• Do not apply more than 0.5 pound active ingredient per acre per season.</li> <li>• Do not apply within 14 days of harvest.</li> </ul> |   |  |

**PEARS**

Apply as directed using the rates in the table below.

| PEST  | RATE  | APPLICATION INSTRUCTIONS   |
|---|---|--|
| Aphids<br>Codling Moth<br>Cutworms<br>Green Fruitworm<br>Leafhopper<br>Leafminer<br>Leafroller<br>Lygus spp.<br>Plant Bugs<br>Plum Curculio<br>San Jose Scale<br>(Crawlers)<br>Stink Bugs<br>Tarnished Plant Bug  | 2.6 to 12.8 fl. oz. (0.04 to<br>0.2 lb. ai) per acre  | <p>Application by ground: Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.</p> <p>Application by air: Apply the specified dosage in a minimum of 10 gallons per acre by air.</p> <p>Apply as necessary to maintain control using a minimum of 30 day spray interval.</p> <p>Apply up to 14 days prior to harvest.</p> |
| Twospotted Spider Mite<br>Yellow Mite   | 3.8 to 12.8 fl. oz. (0.06 to<br>0.2 lb. ai) per acre  |  |
| European Red Mite   | 5.12 to 12.8 fl. oz. (0.08<br>to 0.2 lb. ai) per acre |  |
| <p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 0.5 pound active per acre per season with no more than 0.45 pound active per acre applied after petal fall.</li> <li>• Do not graze livestock in treated orchards or cut treated cover crops for feed.</li> </ul> |   |  |

**ROOT CROPS**

Apply as directed using the rates in the table below.

| CROP  | PEST  | RATE   | APPLICATION INSTRUCTIONS                                 |
|---|---|--|--|
| Burdock, edible<br>Carrot<br>Celeriac<br>Chervil, turnip rooted<br>Chicory<br>Ginseng<br>Horseradish<br>Parsley, turnip rooted<br>Parsnip<br>Radish<br>Radish, oriental<br>Rutabaga<br>Salsify<br>Salsify, black<br>Salsify, Spanish<br>Skirret<br>Turnip | Aphids<br>Beet Armyworm<br>Celery leaf-tier<br>Corn Earworm<br>Cross-Striped Cabbage-worm<br>Cutworms<br>Diamondback moth<br>European Corn Borer<br>Fall Armyworm<br>Fire Ants<br>Flea Beetles<br>Green Cloverworms<br>Hornworms<br>Imported Cabbageworm<br>Loopers<br>Southern Armyworm<br>Spider Mites<br>Tobacco Budworm<br>Velvetbean Caterpillar<br>Whitefly<br>Yellowstriped Armyworm | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.10 lb. ai)<br>per acre | Apply foliar treatments in at least 25 gallons per acre. |
| <b>RESTRICTIONS:</b>  |   |  |  |
| <ul style="list-style-type: none"> <li>• Do not make applications less than 7 days apart.</li> <li>• Do not apply more than 0.5 pound active per acre per season.</li> <li>• Do not apply within 21 days of harvest.</li> </ul>                           |   |  |  |

**ROOT CROPS *cont'd***

|  |   |  |  |
|--|---|--|--|
| Garden Beet  | Aphids<br>Fire Ants<br>Flea Beetles<br>Lepidopterous Larvae<br>Spider Mites<br>Whitefly | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.10 lb. ai)<br>per acre | Apply foliar treatments in at least 25 gallons per acre. |
| <b>RESTRICTIONS:</b>   |   |  |  |
| <ul style="list-style-type: none"> <li>• Do not make applications less than 7 days apart.</li> <li>• Do not apply more than 0.4 lb. active ingredient per acre per season.</li> <li>• Do not apply within 1 day of harvest.</li> </ul> |   |  |  |

**SOYBEAN**

Apply as directed using the rates in the table below.

| <b>PEST</b>   | <b>RATE</b>  | <b>APPLICATION INSTRUCTIONS</b>  |
|---|--|--|
| Alfalfa Caterpillar<br>Aphids<br>Aster Leafhopper<br>Bean Leaf Beetle<br>Beet Armyworm*<br>Cloverworm<br>Corn Earworm<br>Corn Rootworm Adult<br>Cucumber Beetles<br>Cutworms<br>European Corn Borer<br>Fall Armyworm<br>Flea Beetle<br>Grasshoppers<br>Imported cabbageworm<br>Japanese Beetle Adult<br>Leafhoppers<br>Leafminer<br>Loopers<br>Mexican Bean Beetle Adult<br>Pea Leaf Weevil<br>Pea Weevil<br>Plant Bug<br>Saltmarsh caterpillar<br>Sap Beetle | 2.1 to 6.4 fl. oz.<br>(0.033 to 0.1lb. ai)<br>per acre   | Apply as a foliar treatment using at least of 10 gallons per acre at the rate of 6.4 fl. oz. (0.1 lb.) per acre at a minimum of 30 day intervals.<br><br>*Pyrethroid resistance is common for Beet Armyworm and Tobacco Budworm. Consult your local or state agricultural authority to determine if resistant pest populations are in your area. If so refer to the Resistance Management statement in the Directions For Use section of this label. |
| Southern Armyworm<br>Stink Bugs<br>Tarnished Plant Bug<br>Thrips<br>Tobacco budworm1<br>Webworms<br>Western Bean Cutworm<br>Whitefly<br>Yellowstriped Armyworm  | 2.1 to 6.4 fl. oz.<br>(0.033 to 0.1lb. ai)<br>per acre   |  |
| <i>Lygus</i> Species<br>Whitefly<br>Two-Spotted Spider Mite   | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.10 lb. ai)<br>per acre |  |

**SOYBEAN cont'd****RESTRICTIONS:**

- Do not apply more than 0.3 pound active ingredient per acre per season.
- Do not apply within 18 days of harvest.

**SPINACH**

Apply as directed using the rates in the table below.

| PEST  | RATE   | APPLICATION INSTRUCTIONS   |
|---|--|--|
| Armyworms<br>Colorado Potato Beetle<br>Corn Earworm<br>Cucumber Beetles<br>Cutworms<br>European Corn Borer<br>Flea Beetles<br>Leafminers<br>Loopers<br>Pepper Weevil<br>Thrips<br>Tomato Pinworm<br>Tomato Hornworm<br>Whitefly | 2.1 to 6.4 fl. oz.<br>(0.033 to 0.10 lb. ai)<br>per acre | Apply the specified dosage in 5-50 gallons of finished spray per acre by air or 10-50 gallons of finished spray per acre by ground.<br><br>For control of <b>Whiteflies</b> , apply foliar treatments of REVEAL ENDURX by ground or air at rates of up to 0.4 pt. (0.1 lb active) per acre at minimum 7-day intervals up to a maximum of 4 applications.<br><br>For control of <b>Fire Ants</b> , apply REVEAL ENDURX to the soil (at planting) or as a foliar treatment by ground or air at rates up to 0.4 pt. (0.1 lb. active) per acre at minimum 7-day intervals up to a maximum of 4 applications. |
| Broad mite<br>Banks Grass Mite<br>Twospotted Spider Mite<br>Carmine Mite<br>Pacific Spider Mite<br><i>Lygus</i> spp.<br>Fire Ants   | 5.12 to 6.4 fl. oz.<br>(0.08 to 0.10 lb. ai)<br>per acre |  |

**RESTRICTIONS:**

- Do not make applications less than 7 days apart.
- Do not apply more than 0.4 pound active ingredient per acre per season.
- Do not apply within 40 days of harvest.

## SUCCULENT PEAS AND BEANS

Apply as directed at the table rates below.

| CROP  | PEST   | RATE  | APPLICATION INSTRUCTIONS   |
|---|--|---|--|
| <p><b>Peas</b> (<i>Pisum</i> spp.) including:<br/>Dwarf pea<br/>Edible-pod pea<br/>English pea<br/>Garden pea<br/>Green pea<br/>Snow pea<br/>Sugar snap pea<br/>Pigeon pea</p> <p><b>Bean</b> (<i>Phaseolus</i> spp.) including:<br/>Broadbean(succulent),<br/>Lima bean (green),<br/>Runner bean,<br/>Snap bean,<br/>Wax bean</p> <p><b>Bean</b> (<i>Vigna</i> spp.) including:<br/>Asparagus bean,<br/>Blackeyed pea,<br/>Chinese longbean,<br/>Cowpea,<br/>Moth bean,<br/>Southern pea,<br/>Yardlong bean,<br/>Jackbean, Soybean (immature seed),<br/>Sword bean</p> | <p>Flea Beetle<br/>Grasshoppers<br/>Aster Leafhopper<br/>Leafhoppers</p> <p>Aphids Beet Armyworm<br/>Fall Armyworm<br/>Southern Armyworm<br/>Yellowstriped Armyworm<br/>Bean Leaf Beetle<br/>Cucumber Beetle<br/>Japanese Beetle Adult<br/>Sap Beetle<br/>Plant Bugs<br/>Stink Bugs<br/>Tarnished Plant Bug<br/>Alfalfa Caterpillar<br/>Clover Worm<br/>European Corn Borer<br/>Cutworms<br/>Western Bean Cutworm<br/>Corn Earworm<br/>Loopers<br/>Corn Rootworm Adult<br/>Thrips<br/>Webworms<br/>Pea Weevil<br/>Pea Leaf Weevil<br/>Whitefly</p> | <p>1.6 to 6.4 fl. oz. (0.025 to 0.10 lb. ai) per acre</p> <p>2.1 to 6.4 fl. oz. (0.033 to 0.10 lb. ai) per acre</p> | <p>Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.</p> <p>When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.</p> <p>Thorough coverage is essential to achieve control.</p> |
|   | <p>Banks Grass Mite<br/>Twospotted Spider Mite<br/>Carmine Mite<br/><i>Lygus</i> spp.</p>  | <p>5.12 to 6.4 fl. oz. (0.08 to 0.10 lb. ai) per acre</p>   |  |

### RESTRICTIONS:

- Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated product) per acre per season.
- Do not apply within 3 days of harvest.

**TOBACCO**

Apply as directed using the rates in the table below.

| PEST  | RATE  | APPLICATION INSTRUCTIONS   |
|---|---|--|
| Cutworm spp.<br>Tobacco Flea Beetle (larvae)<br>White Grubs<br>Wireworms<br>Mole Crickets<br>Armyworm spp.<br>Stalkborers   | 4.0 to 6.4 fl. oz.<br>(0.0625 to 0.10 lb.<br>ai) per acre | <p><b>Pre-Transplant Soil Application:</b> Apply 0.0625 to 0.1 pounds active ingredient in a minimum of 10 gallons per acre to control soil pests. Use of suitable equipment to incorporate into top 4" of the soil is required to control below ground pests.</p> <p><b>At-Plant Water Treatment Application:</b> Apply 0.0625 to 0.1 pounds active ingredient in a water treatment application volume of 10 to 200 gallons per acre.</p> |
| Aphid spp.<br>Armyworm spp.<br>Flea Beetle (Adult)<br>Chinch bugs<br>Stink bugs<br>Japanese Beetles<br>Grasshoppers<br>Cutworm spp.<br>Tarnished Plant Bugs<br>Green bugs<br>Thrips<br>Whiteflies   | 2.56 to 6.4 fl. oz..<br>(0.04 to 0.10 lb. ai)<br>per acre | <p><b>Foliar Application:</b> Apply 0.04 to 0.1 pounds active ingredient per foliar application up to and including layby in a minimum of 10 gallons per acre.</p>   |
| Spider Mites<br><i>Lygus</i> spp.   | 6.4 fl. oz. ( 0.10 lb.<br>ai) per acre                    |  |
| <p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>• Do not apply later than layby</li> <li>• Do not apply more than 0.2 pound active ingredient per acre per season.</li> <li>• May be tank mixed with other herbicides labeled for tobacco use.</li> <li>• Do not make more than 2 foliar applications per season.</li> </ul> |   |  |

## TUBEROUS AND CORM VEGETABLES

Apply as directed using the rates in the table below.

| CROP   | PEST   | APPLICATION INSTRUCTIONS  |
|--|--|---|
| Potato<br>Sweet potato<br>Arracacha<br>Arrowroot<br>Chinese artichoke<br>Jerusalem artichoke<br>Edible canna<br>Cassava (bitter & sweet)<br>Chayote (root)<br>Chufa<br>Dasheen (taro)<br>Ginger<br>Leren<br>Tanier<br>Turmer<br>Yam bean<br>True yam   | Corn wireworm<br>Tobacco wireworm<br>Southern potato wireworm<br>Japanese beetle grubs<br>June beetle<br>Sweetpotato flea beetle<br>Cucumber beetle<br>Sweetpotato Weevil<br>Banded Cucumber Beetle<br>Black flea beetle<br>Whitefringed beetle<br>White grub<br>Sugarcane beetle<br>Rootworms | <b>In-Furrow At Planting Application:</b> Apply REVEAL ENDURX to control <b>wireworms, rootworms, and white grubs</b> . Apply at the rate of 0.3 pounds active ingredient (19.2 ounces formulated product) per acre as an in-furrow or T-band spray at planting time.<br><br><b>Lay-By Application:</b> Apply REVEAL ENDURX to control <b>wireworms, rootworms, and white grubs</b> . Apply to the drill area and cover with soil utilizing cultivation equipment set to throw soil to the drill area. Apply at the rate of 0.05 to 0.15 pounds active ingredient (3.2 to 9.6 ounces formulated product) in 10 gallons per acre of spray.<br><br><b>Foliar Application:</b> Apply REVEAL ENDURX to control the adult life stages of <b>flea beetles, click beetles (wireworms), cucumber beetles (rootworms), Whitefringed beetles and May/June beetles (White grubs)</b> . Apply at the rate of 0.1 lbs. active ingredient (6.4 ounce formulated product) per acre in 10 gallons of spray by ground equipment and 3 gallons of spray by air. |
| <b>RESTRICTIONS:</b>   |  |   |
| <ul style="list-style-type: none"> <li>Do not make more than 2 foliar applications per season no sooner than 21 days apart.</li> <li>Do not apply more than 0.5 lb. active ingredient per acre per season, including soil application.</li> <li>Do not apply within 21 days of harvest.</li> </ul> |  |   |

## APPLICATIONS INSTRUCTIONS – ORNAMENTALS

For use on plants intended for aesthetic purposes or climatic modifications and being grown in interior landscapes and on outdoor ornamentals, Christmas trees, nurseries, lawns, sod farms and golf courses.

## PRODUCT INSTRUCTIONS

REVEAL ENDURX mixes with water and other aqueous carriers to control listed insects and mites on trees, shrubs, foliage plants, non-bearing fruit and nut trees, and flowers in interiorscapes, including hotels, shopping malls, office buildings, etc. and outdoor landscapes such as, but not limited to, nurseries, residential dwellings, parks, institutional buildings, recreational areas, athletic fields, golf courses, sod farms, and home lawns. Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application.

REVEAL ENDURX may be tank-mixed with other products, including insect growth regulators. When tank mixing REVEAL ENDURX with other products observe all precautions and limitations on each separate product label. The addition of spreader stickers is not necessary. The physical compatibility of REVEAL ENDURX may vary with different sources of pesticide products, and local cultural practices. Any tank mixture which has not been previously tested should be prepared on a small scale (pint or quart jar), using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture.

The following procedure is recommended for preparation of a new tank mix, unless specified otherwise in label directions:

1. Add wettable powders to tank water
2. Agitate
3. Add fluids and flowables
4. Agitate
5. Add emulsifiable concentrates
6. Agitate

If a mixture is found to be incompatible following the order of addition, try reversing the order of addition, or increase the volume of water. **Note:** If the tank mixture is found to be compatible after increasing the amount of water then the sprayer will need to be recalibrated for a higher volume application. Do not allow tank mix to stand overnight. When using tank mixes, observe all restrictions and precautions which appear on the labels of these products. Provide constant agitation to keep the mixture in solution.

## APPLICATION RATES

### TRUNK SPRAYS TO ORNAMENTAL TREES (including Christmas trees)

#### For Control of Bark Beetles and Boring Beetles

Refer to the table below. Application rates and timing differ according to the target pest and other factors specific to each local situation. Consult your local State Extension specialist or other qualified expert for recommendations. Note: Do not apply more than 12.8 fl. oz. (0.2 lbs. AI) per acre of this product to trees. Repeat application may be necessary if reinfestation is likely.

| PEST   | RATE  | SPRAY VOLUME                                  | USE INSTRUCTIONS   |
|--|---|---|--|
| <b>Dendroctonus bark beetles</b> such as mountain pine beetle, southern pine beetle, western pine beetle, and black turpentine beetle. | 16 -32 fl. oz. per 100 gallons (0.25 – 0.5 lb. AI per 100 gallons)  | Use 1-4 gallons of finished spray per tree.   | Make applications to the trunk of the tree with a hydraulic sprayer in the early spring or prior to adult beetle flight and tree infestation.  |
| Engraver beetle ( <i>Ips</i> spp.)   | 16 – 32 fl. oz. per 100 gallons (0.25 – 0.5 lb. AI per 100 gallons) | Use 10-14 gallons of finished spray per tree. | Apply spray directly to the main trunk from the base of the tree to at least half-way into the live crown. Spray until the bark is thoroughly wet.   |
| <b>Other bark beetles</b> such as ambrosia beetles, elm bark beetles, and metallic wood borers such as emerald ash borer.              | 16 – 32 fl. oz. per 100 gallons (0.25 – 0.5 lb. AI per 100 gallons) | Use 2-5 gallons of finished spray per tree.   | Make applications of a spray mixture to the trunk, scaffolding and limbs of the tree with a hydraulic sprayer in the early spring or prior to adult beetle flight and tree infestations. Spray until the bark is thoroughly wet. |

## TRUNK SPRAYS TO ORNAMENTAL TREES (including Christmas trees) cont'd

| PEST   | RATE   | SPRAY VOLUME                                | USE INSTRUCTIONS   |
|--|--|---|--|
| <b>Clearwing moth borers</b><br>such as ash borer, banded ash clearwing, dogwood borer, lesser peachtree borer, lilac borer, oak borer, peachtree borer, rhododendron borer<br><b>Coleopteran borers</b> such as bronze birch borer, flatheaded apple tree borer | 6.4 – 12.8 fl. oz.<br>per 100 gallons<br>(0.1 – 0.2 lb.<br>AI per 100 gallons) | Use 1-4 gallons of finished spray per tree. | Apply to the branches and trunks prior to adult emergence. Spray until the bark is thoroughly wet. For maximum residual control, use highest recommended rate. |

### Treatment of Infested Trees to Control Emerging Brood

Make applications of a spray mixture containing 2.0 pints of REVEAL ENDURX per 100 gallons of water to trees that still have beetles in the bark. Apply spray directly to the main trunk from the base of the tree to at least half-way into the live crown. Spray until the bark is thoroughly wet (usually 1 to 4 gallons of spray per tree). Do not apply more than 0.2 lbs. AI (12.8 fl. oz.) of this product to trees per acre.

Trees on which all needles have turned brown generally have been vacated and should not be sprayed unless infestation is confirmed. To confirm an infestation, scrape off the outer bark to determine if trees are still infested. If live infestations remain in the trunks, fell the trees and cut into sections. Spray the trunk and large limbs and turn sections so that all of the surface area can be treated. Do not apply more than 0.2 lbs. AI (12.8 fl. oz.) of this product to trees per acre.

### FOLIAR SPRAYS TO ORNAMENTALS AND TREES

**(Including Field and Container Grown Nursery Stock, Christmas Trees, Interiorscapes and Plantscapes, Lawns, Trees and Shrubs, and on Golf Courses and Sod Farms**

For applications to ornamentals (including but not limited to trees, shrubs, ground covers, bedding plants and foliage plants, conifers (field and container grown), Christmas Trees and pine seed orchards) apply 0.04 to 0.32 fl. oz. REVEAL ENDURX per 1,000 sq. ft. or 1.8 to 14.4 fl. oz. per 100 gallons. REVEAL ENDURX may be diluted and applied in various volumes of water providing that the maximum label rate (0.32 fl. oz. per 1,000 sq. ft. or 14.4 fl. oz. per 100 gallons) is not exceeded. REVEAL ENDURX may be applied through low volume application equipment by dilution with water or other carriers and providing that the maximum label rate (0.32 fl. oz. per 1,000 sq. ft. or 14.4 fl. oz. per 100 gallons) is not exceeded.

### Calculating Dilution Rates Using the Ornamental Application Rates Table and the REVEAL ENDURX Dilution Chart

Use the following steps to determine the appropriate dilution of this product required to control the specific pests:

1. Find the least susceptible target pest (the pest that requires the highest application rate for control).
2. Select an application rate in terms of fluid ounces of this product.
3. Find your application volume and how much spray you want to prepare.
4. Use the **Ornamental Dilution Chart** to determine the appropriate volume of this product that must be mixed in your desired volume of water.

For example, to control black vine weevil adults on rhododendron, the **Ornamental Application Rates** table shows that 0.08 to 0.16 fl. oz. of this product should be applied per 1,000 sq. ft. You select an application rate of 0.16 fl. oz. per 1,000 sq. ft. because maximum residual control is desired. Your application volume is approximately 300 gallons per acre which is equivalent to 6.9 gallons per 1,000 sq. ft. Consulting the **Ornamental Dilution Chart** shows that you should dilute 0.24 fl. oz. of this product in 10 gallons of water.

| REVEAL ENDURX ORNAMENTAL DILUTION CHART |   |     |           |      |            |      |             |
|---|---|-----|-----------|------|------------|------|-------------|
| Application Rate                        | Fluid Ounces (mL) of REVEAL ENDURX diluted to the Volumes of Finished Spray |     |           |      |            |      |             |
|   | 1 Gallon  |     | 5 Gallons |      | 10 Gallons |      | 100 Gallons |
| Fl. oz./1,000 sq. ft.                   | Fl. oz.   | mL  | Fl. oz.   | mL   | Fl. oz.    | mL   | Fl. oz.     |
| 0.04                                    | 0.018   | 0.5 | 0.09      | 2.6  | 0.18       | 5.3  | 1.8         |
| 0.08                                    | 0.036   | 1.1 | 0.18      | 5.3  | 0.36       | 10.6 | 3.6         |
| 0.16                                    | 0.072   | 2.1 | 0.36      | 10.6 | 0.72       | 21.3 | 7.2         |
| 0.32                                    | 0.144   | 4.3 | 0.72      | 21.3 | 1.44       | 42.6 | 14.4        |

(25)(Fl. Oz. of REVEAL ENDURX added to tank = Percent Active Ingredient of Spray Mix  
(gallons of finished spray mix)(128)

## ORNAMENTAL AND TREE FOLIAR APPLICATION RATES

The application rates listed in the following table will provide excellent control of the noted pests under typical conditions. However, at the discretion of the applicator, this product may be applied at up to 0.32 fl. oz. per 1,000 sq. ft (14.4 f. oz. per 100 gallons) to control each of the pest listed in this table. The higher application rates should be used when maximum residual control is desired.

| PEST   | RATES  | USE INSTRUCTIONS   |
|--|--|--|
| Bagworms <sup>1</sup><br>Cutworms<br>Elm Leaf Beetles<br>Fall Webworms<br>Gypsy Moth Caterpillars<br>Lace Bugs<br>Leaf Feeding Caterpillars<br>Tent Caterpillars<br>Tussock moth   | 0.04 – 0.08 fl.<br>oz. per 1,000<br>sq. ft.<br><br>(1.8 – 3.8 fl. oz.<br>per 100 gallons)      | <sup>1</sup> Bagworms: For best results, apply when larvae begin to hatch and spray larvae directly. Applications when larvae are young will be most effective.<br><br><sup>2</sup> Beetles, Scale Crawlers, Twig Borers, and Weevils: May treat trunks, stems and twigs in addition to plant foliage.<br><br><sup>3</sup> Spider Mites: REVEAL ENDURX provides optimal twospotted spider mite control when applied during spring to mid-summer. Higher application rates and/or more frequent treatments may be required for acceptable twospotted spider mite control during mid- to late-summer. The addition of a surfactant or horticultural oil may increase the effectiveness of this product. Combinations of this product with other registered miticides have also proven effective. Alternately, REVEAL ENDURX applications may be rotated with those of other products that have different modes of action in control programs that are designed to manage resistance by twospotted spider mites. Consult your local Cooperative Extension Service for resistance management recommendations in your region. |
| Adelgids<br>Ants<br>Aphids<br>Bees<br>Beet Armyworm<br>Beetles <sup>2</sup><br>Black Vine Weevil (Adults)<br>Scales, such as<br>Brown Soft Scales<br>California Red Scale (Crawlers) <sup>2</sup><br>Elongated Hemlock Scale<br>Pine Needle Scales (crawlers) <sup>2</sup><br>San Jose Scales (Crawlers) <sup>2</sup><br>Broad Mites<br>Budworms<br>Cicadas<br>Citrus Thrips<br>Clover Mites<br>Crickets<br>Earwigs<br>European Red Mite<br>Flea Beetles<br>Fungus Gnats (Adults)<br>Glassywinged Sharpshooter<br>Grasshoppers | 0.08 – 0.16 fl.<br>oz. per 1,000<br>sq. ft.<br><br>0.16 – 0.32 fl.<br>oz. per 1,000<br>sq. ft. | (Continued from above)   |

**ORNAMENTAL AND TREE FOLIAR APPLICATION RATES *cont'd***

| PEST  | RATES  | USE INSTRUCTIONS   |
|---|--|--|
| Japanese Beetle (Adult)<br>Leafhoppers<br>Leafrollers<br>Mealybugs<br>Mites<br>Mosquitoes<br>Nantucket Pine Tip Moth<br>Pillbugs<br>Pine sawflies<br>Plant Bugs (including Lygus spp.)<br>Psyllids<br>Scorpions<br>Spider Mites <sup>3</sup><br>Spiders<br>Spittlebugs*<br>Thrips<br>Tip Moths<br>Treehoppers<br>Twig Borers <sup>2</sup><br>Wasps<br>Weevils <sup>2</sup> such as<br>White Pine Weevil<br>Pales Weevil<br>Diaprepes adults<br>Orchid Weevil<br>White flies<br>Zimmerman pine moths | 0.08 – 0.16 fl. oz. per 1,000 sq. ft.<br><br>0.16 – 0.32 fl. oz. per 1,000 sq. ft. | <sup>1</sup> Bagworms: For best results, apply when larvae begin to hatch and spray larvae directly. Applications when larvae are young will be most effective.<br><br><sup>2</sup> Beetles, Scale Crawlers, Twig Borers, and Weevils: May treat trunks, stems and twigs in addition to plant foliage.<br><br><sup>3</sup> Spider Mites: REVEAL ENDURX provides optimal twospotted spider mite control when applied during spring to mid-summer. Higher application rates and/or more frequent treatments may be required for acceptable twospotted spider mite control during mid- to late-summer. The addition of a surfactant or horticultural oil may increase the effectiveness of this product. Combinations of this product with other registered miticides have also proven effective. Alternately, REVEAL ENDURX applications may be rotated with those of other products that have different modes of action in control programs that are designed to manage resistance by twospotted spider mites. Consult your local Cooperative Extension Service for resistance management recommendations in your region. |
| Imported Fire Ants**<br>Leafminers<br>Pecan Leaf Scorch Mite<br>Pine Shoot Beetle (Adults)<br>Spider Mites <sup>3</sup>   | (3.6 – 7.2 fl. oz. per 100 gallons)<br>(7.2 – 14.4 fl. oz. per 100 gallons)        |  |

\*\*For foraging ants

**BROADCAST SPRAYS TO TURFGRASS (including lawns, golf courses, sod farms, parks, etc).**

Apply REVEAL ENDURX as a broadcast treatment. Use higher volumes up to 10 gallons of carrier per 1000 square feet to get uniform coverage when treating dense grass foliage.

For low water volume usage, less than 2 gallons/1000 square feet, add a non-ionic or silicone based surfactant (0.25% v/v). Irrigation to treated area within a few hours following application can improve efficacy to listed sub-surface pests.

**Restrictions:**

**In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).**

**In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.**

**Spray Drift Precautions (For Turf & Ornamental Uses)**

Do not apply when wind conditions favor downwind drift to nearby water bodies.

Do not apply when wind velocity exceeds 10 miles per hour. Avoid application when wind gusts approach 10 mph.

Apply using nozzles that provide the largest droplet size compatible with adequate coverage

**Turfgrass Application Rates**

The application rates listed in the following table will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, REVEAL ENDURX may be applied at up to 0.32 fl. oz. per 1000 square feet to control each of the pests listed in this table. The higher application rates should be used when maximum residual control is desired or heavy pest populations occur.

| PEST   | RATES                                  |
|--|--|
| Armyworms <sup>1</sup><br>Cutworms <sup>1</sup><br>Sod Webworm <sup>1</sup>  | 0.05 to 0.08 fl. oz. per 1,000 sq. ft. |
| Annual Bluegrass Weevil (Hyperodes) (Adult) <sup>2</sup><br>Banks Grass Mite <sup>9</sup><br>Billbugs (Adult) <sup>3</sup><br>Black Turfgrass Ataenius (Adult) <sup>4</sup><br>Crickets<br>Earwigs<br>Fleas (Adult)<br>Grasshoppers<br>Mealybugs<br>Mites <sup>6</sup> | 0.08 to 0.16 fl. oz. per 1,000 sq. ft. |
| Ants<br>Chinch Bugs <sup>5</sup><br>Fleas (Larvae) <sup>7</sup><br>Imported Fire Ants <sup>8</sup><br>Japanese Beetle (Adult)<br>Mole Cricket (Adult) <sup>9</sup><br>Mole Cricket (Nymph) <sup>10</sup><br>Ticks <sup>11</sup>  | 0.16 to 0.32 fl. oz. per 1,000 sq. ft. |

**1 Armyworms, Cutworms and Sod Webworms:** To ensure optimum control, delay watering (irrigation) or mowing for 24 hours after application. If the grass area is being maintained at a mowing height of greater than 1 inch, then higher application rates (up to 0.32 fluid oz. per 1000 square feet) may be required during periods of high pest pressure.

**2 Annual Bluegrass Weevil (*Hyperodes*) adults:** Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when Forsythia is in full bloom and concludes when flowering dogwood (*Cornus florida*) is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

**3 Billbug adults:** Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of overwintered chinch bugs.

**4 Black Turfgrass *Ataenius* adults:** Applications should be made during May and July to control the first and second generation of black turfgrass *ataenius* adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with this blooming of Rose of Sharon (*Hibiscus syriacus*).

**5 Chinch Bugs:** Chinch Bugs infest the base of grass plants and are often found in the thatch layer. Irrigation of the grass area before treatment will optimize the penetration of the insecticide to the area where the chinch bugs are located. Use higher volume applications if the thatch layer is excessive or if a relatively long mowing height is being maintained. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates (up to 0.32 fluid oz. per 1000 square feet) may be required to control populations that contain both nymphs and adults during the middle of the summer.

**6 Mites:** To ensure optimal control of eriophyid mites, apply in combination with the labeled application rate of a surfactant. A second application, five to seven days after the first, may be necessary to achieve acceptable control.

**7 Flea larvae:** Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Use a higher volume application when treating these areas to ensure penetration of the insecticide into the soil. Note: if the lawn area is being treated with this product at 0.08 fluid oz. per 1000 square feet for adult flea control, then the larval application rate may be achieved by increasing the application volume two- to four-fold.

**8 Imported Fire Ants:** Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound drenches that will control existing colonies. If the soil is not moist, then it is important to irrigate before application or use a high volume application. Broadcast treatments should apply 0.32 fluid oz. per 1,000 square feet. Treat mounds by diluting 0.05 fluid oz of REVEAL ENDURX per gallon of water and applying 1 to 2 gallons of finished spray per mound. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. A four foot diameter cede around the mound should also be frosted. For best results, apply in cool weather (65 - 80°F) or in early morning or late evening hours. Note: a spray rig that is calibrated to apply 0.32 fluid oz. per 1,000 square feet of this product in 5 gallons per 1,000 square feet contains the approximate dilution (0.05 fluid as per gallon) that is required for fire ant mound drenches in the spray tank.

**9 Mole Cricket adults:** Achieving acceptable control of adult mole crickets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized. Gross

areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

<sup>10</sup> **Mole Cricket nymphs:** Grass areas that received intense adult mole cricket pressure in the sprang should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher application rates and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

<sup>11</sup> **Ticks (Including ticks that may transmit Lyme Disease and Rocky Mountain Spotted fever):** Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher spray volumes when treating areas with dense ground cover or heavy leaf liner. Ticks may be reintroduced from surrounding areas on host animals. Retreatment may be necessary to achieve and/or maintain control during periods of high past pressure. Repeat application is necessary only if there are signs of renewed activity. Repeat application should be limited to no more than once per seven days.

**Deer ticks (bodes spp.)** have a complicated life cycle that ranges over a two year period and involves four life stages. Applications should be made in the late fall and/or early spring to control adult ticks that are usually located on brush or grass above the soil surface and in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter,

**American dog ticks** may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

**REVEAL ENDURX LAWN DILUTION CHART**

| Application Volume: Gallons/ 1000 sq. ft. | Application Rate: Fl. Oz./ 1000 sq. ft. | Fluid Ounces (mL) of REVEAL ENDURX diluted to the Volumes of Finished Spray |      |           |       |            |      |             |  |
|---|---|---|------|-----------|-------|------------|------|-------------|--|
|   |   | 1 Gallon  |      | 5 Gallons |       | 10 Gallons |      | 100 Gallons |  |
|   |   | Fl. oz.   | mL   | Fl. oz.   | mL    | Fl. oz.    | mL   | Fl. oz.     |  |
| 1   | 0.05                                    | 0.05  | 1.48 | 0.25      | 7.39  | 0.50       | 14.8 | 5.00        |  |
| 1   | 0.08                                    | 0.08  | 2.37 | 0.40      | 11.83 | 0.80       | 23.7 | 8.00        |  |
| 1   | 0.16                                    | 0.16  | 4.73 | 0.80      | 23.66 | 1.60       | 47.3 | 16.00       |  |
| 1   | 0.32                                    | 0.32  | 9.46 | 1.60      | 47.32 | 3.20       | 94.6 | 32.00       |  |
| 2   | 0.05                                    | 0.025   | 0.74 | 0.13      | 3.70  | 0.25       | 7.4  | 2.50        |  |
| 2   | 0.08                                    | 0.040   | 1.18 | 0.20      | 5.91  | 0.40       | 11.8 | 4.00        |  |
| 2   | 0.16                                    | 0.080   | 2.37 | 0.40      | 11.83 | 0.80       | 23.7 | 8.00        |  |
| 2   | 0.32                                    | 0.160   | 4.73 | 0.80      | 23.66 | 1.60       | 47.3 | 16.00       |  |
| 3   | 0.05                                    | 0.017   | 0.49 | 0.08      | 2.46  | 0.17       | 4.9  | 1.67        |  |
| 3   | 0.08                                    | 0.027   | 0.79 | 0.13      | 3.94  | 0.27       | 7.9  | 2.67        |  |
| 3   | 0.16                                    | 0.053   | 1.58 | 0.27      | 7.89  | 0.53       | 15.8 | 5.33        |  |
| 3   | 0.32                                    | 0.107   | 3.15 | 0.53      | 15.77 | 1.07       | 31.5 | 10.67       |  |
| 4   | 0.05                                    | 0.013   | 0.37 | 0.06      | 1.85  | 0.13       | 3.7  | 1.25        |  |
| 4   | 0.08                                    | 0.020   | 0.59 | 0.10      | 2.96  | 0.20       | 5.9  | 2.00        |  |
| 4   | 0.16                                    | 0.040   | 1.18 | 0.20      | 5.91  | 0.40       | 11.8 | 4.00        |  |
| 4   | 0.32                                    | 0.080   | 2.37 | 0.40      | 11.83 | 0.80       | 23.7 | 8.00        |  |
| 5   | 0.05                                    | 0.010   | 0.30 | 0.05      | 1.48  | 0.10       | 3.0  | 1.00        |  |
| 5   | 0.08                                    | 0.016   | 0.47 | 0.08      | 2.37  | 0.16       | 4.7  | 1.60        |  |
| 5   | 0.16                                    | 0.032   | 0.95 | 0.16      | 4.73  | 0.32       | 9.5  | 3.20        |  |
| 5   | 0.32                                    | 0.064   | 1.89 | 0.32      | 9.46  | 0.64       | 18.9 | 6.40        |  |
| 10  | 0.05                                    | 0.005   | 0.15 | 0.03      | 0.74  | 0.05       | 1.5  | 0.50        |  |
| 10  | 0.08                                    | 0.008   | 0.24 | 0.04      | 1.18  | 0.08       | 2.4  | 0.80        |  |
| 10  | 0.16                                    | 0.016   | 0.47 | 0.08      | 2.37  | 0.16       | 4.7  | 1.60        |  |
| 10  | 0.32                                    | 0.032   | 0.95 | 0.16      | 4.73  | 0.32       | 9.5  | 3.20        |  |

**Attention**

- Do not apply to pets, crops, or sources of electricity.
- Firewood is not to be treated.
- Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.
- Do not apply this pesticide in livestock buildings (barns).
- Keep children and pets off treated areas following application until the spray has dried.
- Do not apply by air.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE AND SPILL PROCEDURES:** Keep out of reach of children and animals. Store in original containers only, in a cool, dry place and avoid excess heat. Do not freeze. Do not store below 400 F. Carefully open containers.

If crystals are observed, warm material to above 600F by placing container in warm location. Shake or roll container periodically to redissolve solids.

After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills.

**To confine spill:** If liquid, dike surrounding area or absorb with sand, cat litter, or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### **DISPOSAL STATEMENTS:**

**Nonrefillable container:** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org).

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. If recycling is not available puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning. If burned stay out of smoke.

**For packages up to 5 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 56 gallons:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

*Cont'd next page*

## STORAGE AND DISPOSAL *cont'd*

**For refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

- Do not use in greenhouses.
- Do not apply this product through any type of irrigation system. Do not apply when a temperature inversion exists.
- Do not apply for surface feeding pests if rain is expected within 12 hours (or whatever time is necessary for the spray to dry).
- For turf treatment, apply with nozzles not more than 2 feet above the grass.
- Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not apply when grass areas are water logged or the soil is saturated with water (i.e., will not accept irrigation).
- Vinyl and Aluminum Siding: Do not spray directly onto vinyl or aluminum siding. If REVEAL ENDURX inadvertently contacts vinyl and aluminum siding (particularly lightly colored, aged, weathered or otherwise damaged), it may result in staining, bleaching or discoloration. Wash off thoroughly with detergent and water. Factors such as extreme heat and direct sunlight can promote damage when using emulsifiable concentrates. Avoid application to vinyl or aluminum siding while exposed to direct sunlight or during the heat of the day.

## CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of INNICTIS CROP CARE, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold INNICTIS CROP CARE, LLC and Seller harmless for any claims relating to such factors.

INNICTIS CROP CARE, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or INNICTIS CROP CARE, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, INNICTIS CROP CARE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

**To the extent consistent with applicable law, neither INNICTIS CROP CARE, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE**

**LIABILITY OF INVICTIS CROP CARE , LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF INVICTIS CROP CARE , LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

INVICTIS CROP CARE , LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of INVICTIS CROP CARE , LLC.

SPECIMEN

SPECIMEN

INTENTIONALLY LEFT BLANK