

STOLLER® SETT

Specimen Label

For Commercial Application Only
F1601

SETT



Manufactured by Stoller Enterprises, Inc.
4001 W. Sam Houston Parkway North, Suite 100
Houston Texas 77043 USA
Phone: 1-800-539-5283 or 713-461-1493 | Fax: 713-461-4467
E- Mail: stoller@stollerusa.com | Website:
www.StollerUSA.com

Z- SETT

Stoller's SETT

A Foliar Nutritional Spray

GUARANTEED ANALYSIS

Calcium(Ca):..... 8.0%

Boron (B)..... 0.5%

Derived from calcium chloride and sodium borate.

Net Weight: 11.45 lb./ gallon Chlorine (Cl)
maximum is less than 20%.

CAUTION: Keep Out Of Reach Of Children

GENERAL INFORMATION

SETT is designed to be sprayed on plant foliage to reduce localized ethylene production within the plant and guard against calcium replacement in cell walls if used at regular intervals.

Comments on Product Use:

SETT can be tank mixed and applied with most commonly used insecticides and fungicides. **DO NOT** tank mix with growth hormone products, plant growth regulators, or foliar fertilizers unless specifically recommended.

SETT is compatible with the following foliar fertilizers:

Any Stoller Enterprises, Inc. Foliar Fertilizer; Urea; Lignin Sulfonate Complexes; Phenolic Acid Chelates; Potassium Nitrate.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

WARNING

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat and/ or skin. Avoid contact with eyes, skin and clothing. **IF**

SWALLOWED: Induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention. **IF IN**

EYES: Wash eyes with plenty of clear water for at least 15 minutes. Get medical attention. Information regarding the contents and levels of metals in this product is available on the internet at [http:// www.aafpc.org/metals.htm](http://www.aafpc.org/metals.htm).

DIRECTIONS FOR USE

It is more effective to spray small amounts with greater frequency. For maximum yield results, use **SETT** in conjunction with any Stoller Enterprises foliar fertilizer. **DO NOT USE SETT WITH OTHER FOLIAR FERTILIZERS AS PRECIPITATION MAY OCCUR.**

Group I: Cotton, Soybeans, Beans, Tomatoes, Melons, Cucurbits, Peanuts, Peppers. Begin application when first flower appears. Spray enough water to achieve thorough coverage of the entire plant. Spray schedule may be discontinued when last fruit, boll or pods are set.

Spray Intervals	Rate per Acre	Rate per Hectare
7 days	2 quarts	5 liters
10 days	3 quarts	7 liters
14 days	4 quarts	10 liters

Group II: Celery, Cauliflower, Carrots, Beets, Cole crops, Potatoes, Sweet potatoes, Greens, Broccoli, Lettuce. Begin application when the crop goes into "rapid growth" stage or when storage tissue starts to develop. Spray enough water to achieve thorough coverage of the entire plant. Spray schedule may be discontinued when rapid growth ceases or when storage tissue is finished bulking.

Spray Intervals	Rate per Acre	Rate per Hectare
7 days	2 quarts	5 liters
10 days	4 quarts	10 liters

Group III Tree Crops: Citrus fruits, Nuts, Apples, Peaches, Pears, Cherries, Almonds, Tropical fruits. Begin application when buds first form. Spray enough water to achieve thorough coverage of the entire tree. Spray may be discontinued when fruit has sized.

Spray Intervals	Rate per Acre	Rate per Hectare
10-14 days	4 quarts	10 liters

If regular spray intervals have not been maintained, apply 10 quarts/ acre (24 liters/ hectare) at maximum fruit set or just before period when drop occurs.

Group IV: Grapes. Begin application when fruit first starts to form. Spray enough water to achieve thorough coverage of entire plant canopy. Spray may be discontinued when grapes are sized.

Spray Intervals	Rate per Acre	Rate per Hectare
7 days	2 quarts	5 liters
10 days	3 quarts	7 liters
14 days	4 quarts	10 liters

WARNING

This product contains Boron and the use of this product on any crop other than those recommended by this label may result in serious injury to the crop.