

QUALI-PRO[®]

Strobe[™] 2L

BROAD SPECTRUM FUNGICIDE
FOR CONTROL OF TURF AND ORNAMENTAL DISEASES

ACTIVE INGREDIENT:

Azoxystrobin: methyl
(E)-2-[2-[6-(2-cyanophenoxy)
pyrimidin-4-yloxy]Phenyl]-3-
methoxyacrylate***22.9%**

OTHER INGREDIENTS: **77.1%**

TOTAL: **100.0%**

*IUPAC

Contains 2.08 lb. of active ingredient per gallon
Suspended Concentrate

CONTENTS: 1 GALLON

GROUP 11 FUNGICIDE



**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See additional precautionary statements
and directions for use inside booklet.

Reformulation is prohibited. See individual
container labels for repackaging limitations.

EPA Reg. No. 42750-261-53883

EPA Est. No. 53883-TX-002



Manufactured for:
**Control
Solutions Inc.**
Pasadena, TX 77507



0 72693 00168 8

QUALI-PRO[®]

Strobe[™] 2L

BROAD SPECTRUM FUNGICIDE

FOR CONTROL OF TURF AND ORNAMENTAL DISEASES

ACTIVE INGREDIENT:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate*..... 22.9%

OTHER INGREDIENTS:..... 77.1%

TOTAL: 100.0%

*IUPAC

Contains 2.08 lb. of active ingredient per gallon
Suspended Concentrate

CONTENTS: 1 GALLON

GROUP 11 FUNGICIDE



**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See additional precautionary statements and
directions for use inside booklet.

Reformulation is prohibited. See individual container
labels for repackaging limitations.

EPA Reg. No. 42750-261-53883

EPA Est. No. 53883-TX-002

Manufactured by:



**Control
Solutions Inc.**

Pasadena, TX 77507

| FIRST AID | |
|--|---|
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International for emergency medical treatment at (866) 897-8050.</p> | |

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
3. Shoes plus socks

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Control Solutions immediately if you observe any adverse environmental effects due to use of this product.

Physical or Chemical Hazards

Do not mix or allow coming into contact with oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use of Strobe™ 2L through air blast application equipment on grapes is prohibited in the following townships and boroughs of Erie County, Pennsylvania: North East, Harborcreek, Lawrence Park, Erie, Presque Isle, Millcreek, Fairview, Girard and Springfield.

This prohibition is intended to help eliminate phytotoxicity problems with apples observed in this geographic location.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

1. Coveralls
2. Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

Do not treat areas while unprotected humans or domestic animals are present in the treatment areas. Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

Do not allow entry into treatment area until area that was treated with this product is dry.

PRODUCT INFORMATION

Strobe™ 2L is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. These additional benefits are due to positive effects on plant physiology. The effects may vary according to factors such as the crop, crop hybrid, or environment. Strobe™ 2L may be applied as a foliar spray in alternating spray programs or in tank-mixes with other registered crop protection products. All applications must be made according to the use directions that follow.

USE RESTRICTIONS

DO NOT spray Strobe™ 2L where spray drift may reach apple trees.

DO NOT use spray equipment which has been previously used to apply Strobe™ 2L to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

DO NOT graze or feed clippings from treated turf areas to animals.

DO NOT use in greenhouses.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply Strobe™ 2L to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

USE PRECAUTIONS

Strobe™ 2L is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Strobe™ 2L may demonstrate some phytotoxic effects when mixed with products that are formulated as ECs. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

PRODUCT USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Adjuvants: When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification is recommended.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of Strobe™ 2L has been used. If resistant isolates to Group 11 fungicides are present, efficacy can be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, with highly susceptible varieties, or when environmental conditions are conducive to disease.

INTEGRATED PEST (DISEASE) MANAGEMENT

Strobe™ 2L should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. This should include selection of varieties with disease tolerance, removal of plant debris in which inoculum overwinters, and proper timing and placement of irrigation. Consult your local agricultural authorities for additional IPM strategies established for your area. Strobe™ 2L may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

Crop Tolerance: Plant tolerance has been found to be acceptable for all crops on the label, however, not all possible tank-mix combinations have been tested under all conditions. When possible, it is recommended to test the combinations on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application. See Product Use Precautions for apple phytotoxicity information.

RESISTANCE MANAGEMENT

GROUP 11 FUNGICIDES

Strobe™ 2L (azoxystrobin) is a Group 11 fungicide. The mode of action for Strobe™ 2L is the inhibition of the Qol (quinone outside) site within the electron transport system [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label.

Resistance management strategies may include alternating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. Control Solutions encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Follow the crop specific resistance management recommendations in the directions for use.

If no resistance recommendation on number of applications is specified in the directions for use, follow the recommendations in the table below.

| | | | | | | | | | | | | |
|--|---|---|---|---|---|---|---|---|---|----|----|----|
| If planned total number of fungicide applications per crop is: | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| Recommended Solo Qol fungicide sprays | 1 | 1 | 2 | 2 | 2 | 2 | 2 | 3 | 3 | 3 | 3 | 4 |
| Recommended Qol fungicide sprays in mixture (tank-mix or formulated) | 1 | 2 | 2 | 2 | 2 | 3 | 3 | 4 | 4 | 5 | 5 | 6 |

In situations requiring multiple sprays, develop season-long spray programs for Group 11 (Qol) fungicides. In crops where two sequential Group 11 fungicide applications are made, they should be alternated with two or more applications of a fungicide that is not in Group 11. If more than 12 applications are made, observe the following guidelines:

- When using a Qol fungicide as a solo product, the number of applications must be no more than 1/3 (33%) of the total number of fungicide applications per season.
- For Qol mixes in programs in which tank-mixes or pre mixes of Qol with mixing partners of a different mode of action are utilized, the number of Qol containing applications must be no more than 1/2 (50%) of the total number of fungicide applications per season.
- In programs in which applications of Qol are made with both solo products and mixtures, the number of Qol containing applications must be no more than 1/2 (50%) of the total number of fungicide applications per season.

If a Group 11 fungicide is applied to the seed or soil, do not make another application with a Group 11 fungicide for at least 3 weeks.

SPRAY DRIFT MANAGEMENT

To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

MIXING AND APPLICATION METHODS

Spray Equipment

Strobe™ 2L may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on the suction side of the pump should be 16-mesh or coarser.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

Pump

- Use a pump with capacity to:
 1. Maintain 35-40 psi at nozzles
 2. Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

Mixing Instructions

- Strobe™ 2L is a suspension concentrate (SC) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Strobe™ 2L Alone (No Tank-mix)

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add Strobe™ 2L to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after Strobe™ 2L has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

Strobe™ 2L + Tank-Mixtures: Strobe™ 2L is usually compatible with all tank-mix partners listed on this label. To determine the physical compatibility of Strobe™ 2L with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Strobe™ 2L has demonstrated some phytotoxic effects when mixed with products that are formulated as emulsifiable concentrates (EC). These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

Mixing in the Spray Tank

- Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Strobe™ 2L to the spray tank.
- Allow Strobe™ 2L to completely disperse.
- Spray the mixture with the agitator running.

APPLICATION INSTRUCTIONS THROUGH IRRIGATION SYSTEMS (CHEMIGATION)**Application Through Irrigation Systems (Chemigation)**

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.1-0.25 inches/acre. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Spray Preparation: Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.

Drip Irrigation: Strobe™ 2L may be applied through drip irrigation systems for soil borne disease control. The soil should have adequate moisture capacity prior to drip application.

Terminate drip irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, subsequent irrigation (water only) should be delayed for at least 24 hours following drip application.

Sprinkler Irrigation

- Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems.
- Do not apply this product through any other type of irrigation system except as specified on this label.
- Apply with center pivot or continuous-move equipment distributing ½ acre-inch or less during treatment.
- In general, use the least amount of water required for proper distribution and coverage.
- If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, this product should be injected into no more than the last 20-30 minutes of the set.
- Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Plant injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water.
- Thorough coverage of foliage is required for good control.
- Good agitation should be maintained during the entire application period.

If you have questions about calibration you should contact State Extension Service specialist, equipment manufacturers or other experts.

Operating Instructions

1. Do not apply when wind speed favors drift beyond the area intended for treatment.
2. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
9. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Center Pivot Irrigation Equipment

Notes: (1) Use only with drive systems which provide uniform water distribution. (2) Do not use end guns when chemigating Strobe™ 2L through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply 1/8 to 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as specified by the equipment manufacturer.
- When applying Strobe™ 2L through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of Strobe™ 2L required to treat the area covered by the irrigation system.

- Add the required amount of Strobe™ 2L and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the Strobe™ 2L solution.
- Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the Strobe™ 2L solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying Strobe™ 2L through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of Strobe™ 2L required to treat the area covered by the irrigation system.
- Add the required amount of Strobe™ 2L into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the Strobe™ 2L solution has cleared the last sprinkler head.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Strobe™ 2L Rate Conversion Chart

| fl. oz. product/A | lb ai/A | treated acres/gal. product |
|-------------------|---------|----------------------------|
| 4.0 | 0.07 | 32.0 |
| 5.0 | 0.08 | 25.6 |
| 5.5 | 0.09 | 23.2 |
| 6.0 | 0.10 | 21.3 |
| 6.2 | 0.10 | 21.3 |
| 7.0 | 0.11 | 18.3 |
| 8.5 | 0.14 | 15.4 |
| 9.0 | 0.15 | 14.2 |
| 9.2 | 0.15 | 14.2 |
| 10.0 | 0.16 | 13.0 |
| 11.0 | 0.18 | 11.6 |
| 12.0 | 0.20 | 10.4 |
| 12.3 | 0.20 | 10.4 |
| 13.0 | 0.21 | 9.8 |
| 14.0 | 0.23 | 9.1 |
| 15.4 | 0.25 | 8.3 |
| 15.5 | 0.25 | 8.3 |
| 18.3 | 0.30 | 6.9 |
| 18.5 | 0.30 | 6.9 |
| 20.0 | 0.33 | 6.4 |
| 20.3 | 0.33 | 6.4 |
| 24.5 | 0.40 | 5.2 |

TURF

Golf course turf (not for use in California). **Commercial turf farms** (not for use in California).

Strobe™ 2L is recommended for control of anthracnose, brown patch, cool weather brown patch (yellow patch), Fusarium patch, gray leaf spot, gray snow mold (Typhula blight), leaf spot, melting out, necrotic ring spot, pink patch, pink snow mold, Pythium blight, Pythium root rot, red thread, Rhizoctonia large patch, southern blight, spring dead spot, summer patch, take-all patch, and Zoysia patch on **golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.**

Integrated Pest (Disease) Management:

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

Resistance Management:

Some turf disease pathogens are known to have developed resistance to products used repeatedly for their control. Strobe™ 2L should be applied in a tank-mix or alternation program with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Do not apply more than two sequential Strobe™ 2L applications for *Pythium* spp. control. For all other diseases when *Pythium* spp. is not present, do not apply more than three sequential applications of Strobe™ 2L.

Application Directions:

Strobe™ 2L should be applied prior to disease development. Mix Strobe™ 2L with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1000 square feet (87-174 gallons per acre). Repeat applications at specified intervals for as long as required. For spot treatments, use 0.4 fl. oz. Strobe™ 2L per 1 to 2 gallons of water. Do not apply more than 9.6 quarts product/acre/year (7.1 fl. oz. product/1000 square feet/year). Apply by ground only.

Rate Ranges:

Use the shortest specified application interval and/or use the higher specified rate when prolonged favorable disease conditions exist.

Dollar Spot:

Strobe™ 2L does not control dollar spot. Strobe™ 2L is compatible in tank-mixes with many other fungicides that control dollar spot. Always tank-mix Strobe™ 2L with another fungicide that controls dollar spot when this disease is present.

Follow directions under **Strobe™ 2L + Tank-Mixtures** above.

DIRECTIONS FOR APPLICATION FOR TURF DISEASES

| Target Diseases | Use Rate (fl. oz. product per 1000 sq. ft.) | Application Interval (days) | Remarks* |
|--|---|-----------------------------------|---|
| Anthraxnose (Colletotrichum graminicola) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Brown patch (Rhizoctonia solani) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Cool weather brown patch Yellow patch (Rhizoctonia cerealis) | 0.38-0.77 | 28 | Make one or two applications in fall or when conditions are favorable for disease development. |
| Fusarium patch (Microdochium nivale) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Gray leaf spot (Pyricularia grisea) | 0.38-0.77 | 14-28 | Begin applications before disease is present and continue applications while conditions are favorable for disease development. |
| Gray snow mold Typhula blight (Typhula incarnata, T. ishikariensis) | 0.77-1.35 | Single application 14 | Make a single application of 1.35 fl. oz. or two applications of 0.77 spaced 14 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide may enhance control under severe disease pressure. |
| Leaf spot (Bipolaris sorokiniana) | 0.38-0.77 | 14-21 | Apply when conditions are favorable for disease development. |

(continued)

| Target Diseases | Use Rate (fl. oz. product per 1000 sq. ft.) | Application Interval (days) | Remarks* |
|---|---|-----------------------------------|---|
| Melting out (Drechslera poae) | 0.38-0.77 | 14-21 | Apply when conditions are favorable for disease development. |
| Necrotic ring spot (Leptosphaeria korrae) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Pink patch (Limonomyses roseipellis) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Pink snow mold (Microdochium nivale) | 0.77-1.35 | Single application 14 | Make a single application of 1.35 fl. oz. or two applications of 0.77 spaced 14 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide may enhance control under severe disease pressure. |
| Pythium blight Pythium root rot (Pythium aphanidermatum, Pythium spp.) | 0.38-0.77 | 10-14 | Begin applications before disease is present. During periods of prolonged favorable conditions, treat on the 10 day application interval. For use on newly seeded as well as established turf. |
| Red thread (Laetisaria fuciformis) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Rhizoctonia large patch (Rhizoctonia solani) | 0.38-0.77 | 28 | Make one or two applications in fall or when conditions are favorable for disease development. |
| Southern blight (Sclerotium rolfsii) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Spring dead spot (Leptosphaeria korrae) or (Gaeumannomyces graminis var. graminis) or (Ophiosphaerella herpotricha) | 0.38-0.77 | 28 | Make one or two applications in fall or when conditions are favorable for disease development. |

(continued)

| Target Diseases | Use Rate (fl. oz. product per 1000 sq. ft.) | Application Interval (days) | Remarks* |
|---|---|-----------------------------------|--|
| Summer patch (Magnaporthe poae) | 0.38-0.77 | 14-28 | Apply when conditions are favorable for disease development. |
| Take-all patch (Gaeumannomyces graminis var. avenae) | 0.38-0.77 | 28 | Make two applications 28 days apart in the spring and two applications 28 days apart in the fall. |
| Zoysia patch (Rhizoctonia solani and/or Gaeumannomyces in crustana) | 0.38-0.77 | 28 | Make one or two applications in late fall before snow cover or when conditions are favorable for disease development. Do not apply on top of snow. |

* Do not apply more than two sequential applications of Strobe™ 2L for control of *Pythium* spp. For all other diseases, do not apply more than three sequential applications of Strobe™ 2L.

Strobe™ 2L Rate Conversion Chart for Turf

| Fluid Ounces Product Per 1000 Sq. Ft. | Ounces AI Per 1000 Sq. Ft. | Fluid Ounces Product Per Acre | Pints of Product Per Acre |
|---|-------------------------------|-------------------------------------|---------------------------------|
| 0.4 | 0.104 | 17.4 | 1.1 |
| 0.5 | 0.130 | 21.8 | 1.4 |
| 0.6 | 0.156 | 26.1 | 1.6 |
| 0.7 | 0.182 | 30.5 | 1.9 |
| 0.77 | 0.200 | 33.5 | 2.1 |
| 1.35 | 0.350 | 58.8 | 3.7 |

Amount of Strobe™ 2L to Mix 100 Gallons for Turf Applications

| Spray Volume (gallons/1000 square feet) | | | |
|---|------------------------|------------------------|------------------------|
| Strobe™ 2L Use Rate (fl. oz.) | 2.0 gals. (fl. oz.) | 3.0 gals. (fl. oz.) | 4.0 gals. (fl. oz.) |
| 0.4 | 20 | 13 | 10 |
| 0.5 | 25 | 17 | 13 |
| 0.6 | 30 | 20 | 15 |
| 0.7 | 35 | 23 | 18 |
| 0.77 | 38.5 | 25.7 | 19.3 |
| 1.35 | 67.5 | 45 | 33.75 |

ORNAMENTALS

Strobe™ 2L is recommended for control of certain pathogens causing foliar, aerial, and root diseases, including leaf, tip, and flower blights, leaf spots, downy mildew, powdery mildew, anthracnose, and rusts of ornamental plants. **Strobe™ 2L may be used to control certain diseases of container, bench, flat, plug, bed or field-grown ornamentals in greenhouses, shade-houses, outdoor nurseries, retail nurseries, and other landscape areas.**

INTEGRATED PEST (DISEASE) MANAGEMENT: Strobe™ 2L should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation. Immunoassay detection kits and diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

RESISTANCE MANAGEMENT: Some ornamental disease pathogens are known to have developed resistance to fungicides used repeatedly for their control. Strobe™ 2L should be applied in an alternation or tank-mix program with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Do not make more than three (3) sequential applications of Strobe™ 2L before alternating with a fungicide of a different mode of action. A sound resistance management program would include blocks of three Strobe™ 2L applications separated by blocks of two alternate fungicide applications. Do not alternate Strobe™ 2L with other strobilurin fungicides.

APPLICATION DIRECTIONS: Apply Strobe™ 2L as a broadcast or banded spray targeted at the foliage or crown of the plant. Apply to runoff in sufficient water to ensure complete coverage of the target plant. Good coverage and wetting of foliage is necessary for best control. Refer to the label for specific use directions for control of certain diseases. Repeat applications at specified intervals (plus alternations for resistance management) for as long as required. Applications may be made by ground only.

Strobe™ 2L applications should begin prior to disease development and continue throughout the season at specified intervals following resistance management guidelines. Strobe™ 2L works best when used as part of a preventative disease management program.

Use only surfactants approved for ornamental plants in combination with Strobe™ 2L. Do not use silicone based products with Strobe™ 2L due to possible phytotoxicity. Always test tank-mixes on a small group of representative plants prior to broadscale use.

Apply Strobe™ 2L at use rates of 1.9 - 7.7 fl. oz./100 gallons (0.95 - 3.85 fl. oz./50 gallons) and every 7-28 days (or as otherwise specified for a specific plant or disease). The addition of a non-silicone based wetter-sticker at the recommended use rate may enhance coverage on hard-to-wet plant foliage.

Under most conditions and for most diseases, apply 3.85 - 7.7 fl. oz./100 gallons (1.9 - 3.85 fl. oz./50 gallons) on a 7-14 day interval.

Under light to moderate disease pressure, use the lower rates (1.9 - 3.85 fl. oz./100 gallons, or 0.95 - 1.9 fl. oz./50 gallons) on a 7-14 day interval or the higher rates (5.75 - 7.7 fl. oz./100 gallons or 2.85 - 3.85 fl. oz./50 gallons) on a 14-28 day interval.

Under environmental conditions which promote severe disease development, use the higher rates (5.75 - 7.7 fl. oz./100 gallons or 2.85 - 3.85 fl. oz./50 gallons) on a 7-14 day interval.

Use of Strobe™ 2L as a "rescue" (late curative or eradicator) treatment may not always result in satisfactory disease control.

Do not exceed 2.4 gallons of product/crop acre/year or 8 applications/crop/year.

Do not exceed 600 gallons spray volume per acre for foliar applications. For drench and crown applications, do not exceed 2 pints volume per square foot.

In addition, do not tank-mix Strobe™ 2L with other fungicides, insecticides, herbicides, fertilizers, adjuvants, etc., unless local experience indicates that the tank-mix is safe to ornamental plants.

DRENCH APPLICATION: Strobe™ 2L may be applied to control soil-borne, seedling, and crown diseases of production ornamentals (greenhouse, shadehouse, and container grown) as a preventative, drench treatment prior to infection. Good coverage of the pre-infection area (root zone, root ball, crown, etc.) is necessary for satisfactory control. Strobe™ 2L may be drench applied to container grown ornamentals using 0.38 - 1.75 fl. oz./100 gallons of water. Apply 1-2 pints of the solution per square foot surface area on a 7-28 day interval. Apply drench prior to infection as healthy roots are necessary to optimize product uptake, systemic translocation and disease protection.

For resistance management do not make more than three sequential drench applications of Strobe™ 2L before alternating with a fungicide of a different mode of action.

Caution should be taken before making application of Strobe™ 2L as a drench to small bedding plants in the seedling/plug stage due to possible phytotoxicity. A limited quantity of plants should be tested prior to full-scale application.

DRIP IRRIGATION: Strobe™ 2L may be applied through drip irrigation systems to potted ornamentals or to bedded, field-grown ornamentals for soil-borne disease control. Apply 3.85 - 30.75 fl. oz. Strobe™ 2L per acre as a preventative disease application. The soil or potting media should have adequate moisture capacity prior to drip application.

Terminate drip irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, subsequent irrigation (water only) should be delayed for at least for 24 hours following drip application.

GENERAL ORNAMENTAL USE PRECAUTIONS

Do not apply Strobe™ 2L to apple or cherry trees (Flowering, Yoshina variety) due to possible phytotoxicity. Further, do not use spray equipment that has applied Strobe™ 2L for use in these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.

Strobe™ 2L may be applied to certain varieties of crabapple for control of apple scab. Strobe™ 2L has been shown to be safer when applied to the species and varieties listed in Table 4. However, due to the large number of genera, species, and varieties of crabapple, it is impossible to test every one for tolerance to Strobe™ 2L. The professional user should conduct small scale testing to insure plant safety prior to broad-scale commercial use on plant genera and species not listed on this label.

TABLE 1: DISEASES CONTROLLED: When used in accordance with the label directions, Strobe™ 2L will provide control of the following diseases of ornamental plants:

| DISEASE (Pathogen) | Use Rates and Remarks | |
|--|--|--|
| | 8 oz. and larger containers (fl. oz. product per 100 gallons) | 4 oz. containers (fl. oz. product per 50 gallons) |
| 1. CONIFER BLIGHTS | | |
| a. Phomopsis blight (<i>Phomopsis juniperovora</i>) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| b. Tip blight (<i>Sirococcus strobilinus</i>) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| 2. LEAF BLIGHTS/LEAF SPOTS | | |
| a. Alternaria leaf spot (<i>Alternaria</i> spp.) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| b. Anthracnose (<i>Colletotrichum</i> spp., <i>Elsinoë</i> spp.) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| c. Downy mildew of rose (<i>Peronospora sparsa</i>) | Apply 3.85 - 7.7 fl. oz. every 7-21 days during periods of active plant growth and prior to dormancy or severe infection. | Apply 1.9 - 3.85 fl. oz. every 7-21 days during periods of active plant growth and prior to dormancy or severe infection. |
| d. Entomosporium leaf spot (<i>Entomosporium mespili</i>) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| e. Iris leaf spot (<i>Mycosphaerella macrospora</i>) | Apply 3.85 - 7.7 fl. oz. every 7-21 days. | Apply 1.9 - 3.85 fl. oz. every 7-21 days. |
| f. Leaf spot (<i>Cladosporium echinulatum</i>) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| g. Rose black spot (<i>Diplocarpon rosae</i>) | Apply 7.7 - 15.4 fl. oz. every 7-14 days. Apply Strobe™ 2L on a 7 day interval unless disease pressure is light. Under severe disease conditions or if disease is already present, Strobe™ 2L may be tank-mixed with another rose black spot fungicide. Do not exceed 46 fl. oz./acre/application. | Apply 3.85 - 7.7 fl. oz. every 7-14 days. Apply Strobe™ 2L on a 7 day interval unless disease pressure is light. Under severe disease conditions or if disease is already present, Strobe™ 2L may be tank-mixed with another rose black spot fungicide. Do not exceed 46 fl. oz./acre/application. |

| DISEASE (Pathogen) | Use Rates and Remarks | |
|---|--|---|
| | 8 oz. and larger containers (fl. oz. product per 100 gallons) | 4 oz. containers (fl. oz. product per 50 gallons) |
| h. Myrothecium leaf spot (<i>Myrothecium</i> spp.) | Apply 3.85 - 7.7 fl. oz. every 7-21 days. | Apply 1.9 - 3.85 fl. oz. every 7-21 days. |
| i. Downy mildew of bedding plants (<i>Peronospora</i> spp.) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| j. Scab (<i>Venturia inaequalis</i>) | Apply 1.9 - 7.7 fl. oz. every 10-28 days. Do not apply to apple trees. For crabapples only, see Table 4 for tolerant species. | Apply 0.95 - 3.85 fl. oz. every 10-28 days. Do not apply to apple trees. For crabapples only, see Table 4 for tolerant species. |
| k. Marssonina leaf spot (<i>Marssonina</i> spp.) | Apply 1.9 - 7.7 fl. oz./100 gal every 14-28 days. | Apply 0.95 - 3.85 fl. oz. every 14-28 days. |
| l. Cercospora leaf spot | Apply 1.9 - 7.7 fl. oz./100 gal every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| 3. POWDERY MILDEW | Preventative applications only. Do not make more than 2 sequential applications before rotating to another class of fungicide. | Preventative applications only. Do not make more than 2 sequential applications before rotating to another class of fungicide. |
| a. <i>Erysiphe pannosa</i> , <i>Erysiphe</i> spp. | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| b. <i>Microsphaera azaleae</i> | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| c. <i>Sphaerotheca pannosa</i> | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| 4. RUSTS | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| a. Needle rust (<i>Melampsora occidentalis</i>) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| b. <i>Phragmidium</i> spp. | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| c. <i>Puccinia</i> spp. | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| d. <i>Gymnosporangium</i> spp. | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |
| 5. FLOWER BLIGHTS | | |
| a. Anthracnose (<i>Collectotrichum</i> spp., <i>Elsinoë</i> spp.) | Apply 1.9 - 7.7 fl. oz. every 7-28 days. | Apply 0.95 - 3.85 fl. oz. every 7-28 days. |

(continued)

| DISEASE (Pathogen) | Use Rates and Remarks | |
|---|---|---|
| | 8 oz. and larger containers (fl. oz. product per 100 gallons) | 4 oz. containers (fl. oz. product per 50 gallons) |
| b. <i>Botrytis slight (Botrytis cinerea)</i> | Apply 7.7 - 15.4 fl. oz. every 7-21 days. For suppression only. Do not exceed 46 fl. oz./acre. | Apply 3.85 - 7.7 fl. oz. every 7-21 days. For suppression only. Do not exceed 46 fl. oz./acre. |
| 6. SHOOT/STEM DISEASES | | |
| a. Aerial/shoot blight (<i>Phytophthora</i> spp.) | Apply 1.9 - 3.85 fl. oz. every 7-28 days. | Apply 0.95 - 1.9 fl. oz. every 7-28 days. |
| 7. SOILBORNE DISEASES (Directed Spray) | | |
| | For directed spray applications utilize the following rates below. | For directed spray applications utilize the following rates below. |
| a. <i>Rhizoctonia solani</i> | Apply 1.9 - 7.7 fl. oz. every 7-21 days. | Apply 0.95 - 3.85 fl. oz. every 7-21 days. |
| b. <i>Sclerotium rolfsii</i> | Apply 1.9 - 7.7 fl. oz. every 7-21 days. | Apply 0.95 - 3.85 fl. oz. every 7-21 days. |
| c. <i>Fusarium</i> spp. | Apply 1.9 - 7.7 fl. oz. every 7-21 days. | Apply 0.95 - 3.85 fl. oz. every 7-21 days. |
| 8. SOILBORNE DISEASES (Drench) | | |
| | See Ornamentals Section for additional drench directions. | See Ornamentals Section for additional drench directions. |
| a. <i>Rhizoctonia solani</i> | Apply 0.35 - 1.75 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days. | Apply 0.19 - 0.95 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days. |
| b. <i>Sclerotium rolfsii</i> | Apply 0.35 - 1.75 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days. | Apply 0.19 - 0.95 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days. |

PLANT SAFETY: Strobe™ 2L has been shown to be safe when applied to the ornamental plants listed in Tables 2, 3, and 4. However, due to the large number of genera, species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to Strobe™ 2L. Neither the manufacturer nor the seller has determined whether or not Strobe™ 2L can be used safely on genera, species, or varieties of ornamental and nursery plants not specified on this label. The professional user should conduct small scale testing to insure plant safety prior to broadscale commercial use on plant genera and species not listed in this label. In addition, do not tank-mix Strobe™ 2L with other fungicides, insecticides, herbicides, fertilizer, adjuvants, etc., unless local experience indicates that the tank-mix is safe to ornamental plants.

Do not apply Strobe™ 2L to certain apple, crabapple or cherry trees due to possible phytotoxicity. Further, do not use spray equipment that has applied Strobe™ 2L for use in these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.

Tolerant Ornamental Plants: Strobe™ 2L has been found to be safe when applied to the plants listed in Tables 2, 3 and 4 when applied according to recommended application methods, rates, and timings:

TABLE 2: Tolerant Plants Listed by Botanical Name:

| BOTANICAL NAME | COMMON NAME | DISEASES |
|---|----------------------------------|----------|
| <i>Abelia</i> spp. | Abelia | 2 |
| <i>Abies fraseri</i> | Fraser Fir | 1, 4 |
| <i>Abies procera</i> | Noble Fir | 1, 4 |
| <i>Acer palmatum</i> | Japanese Maple | 2 |
| <i>Acer saccharum</i> | Sugar Maple | 2 |
| <i>Ageratum</i> spp. | Floss-Flower | 3, 4 |
| <i>Ageratum</i> spp. | Pussy's-Foot | 3, 4 |
| <i>Aglaonema</i> spp. | Chinese-Evergreen | 2, 4 |
| <i>Ajuga reptans</i> | Bugle, Bugleweed | 3 |
| <i>Antirrhinum</i> spp. | Snapdragon | 2i, 3, 4 |
| <i>Aphelandra</i> spp. | Zebra Plant | 2 |
| <i>Artemisia</i> spp. | Mugwort-Sagebrush | 2 |
| <i>Artemisia</i> spp. | Wormwood | 2 |
| <i>Aster</i> spp. | Aster, Starwort | 4 |
| <i>Aucuba japonica</i> | Japanese aucuba, Japanese laurel | 7 |
| <i>Begonia</i> spp. (except Rieger begonia) | Begonia | 2, 3 |
| <i>Berberis thunbergii</i> | Barberry | 3, 4 |
| <i>Betula nigra</i> | River Birch | 3, 4 |
| <i>Bougainvillea</i> spp. | Bougainvillea | 2 |
| <i>Brassaia actinophylla</i> | Rubber-Tree, Umbrella-Tree | 2, 7 |
| <i>Buddleia davidii</i> | Buddleia, Butterfly-Bush | 2 |
| <i>Buxus sempervirens</i> | Boxwood | 2, 7a |

| BOTANICAL NAME | COMMON NAME | DISEASES |
|---------------------------------|--|----------|
| <i>Caladium</i> spp. | Caladium | 7 |
| <i>Camelia japonica</i> | Camelia | 2 |
| <i>Caryota urens</i> | Sago Palm | 2,7 |
| <i>Catharanthus roseus</i> | Vinca | 2 |
| <i>Ceanoihus sanguineus</i> | Wild Lilac | 3 |
| <i>Ceanothus</i> spp. | Ceanothus, California Lilac, Snowball | 3 |
| <i>Cedrus atlantica</i> | Atlas Cedar | 2, 4 |
| <i>Cedrus</i> spp. | White Cedar | 2, 4 |
| <i>Cercis occidentalis</i> | Western Redbud | 2 |
| <i>Chamaecyparis</i> spp. | Cypress, Leyland Cypress | 1 |
| <i>Chamaecyparis pisifera</i> | Sawara Cypress | 1 |
| <i>Chamaedorea elegans</i> | Parlor Palm | 7 |
| <i>Chrysanthemum</i> spp. | Chrysanthemums | 2, 7c |
| <i>Clethra alnifolia</i> | Clethra, White Alder | 2 |
| <i>Cornus</i> spp. | Dogwood, Pink Dogwood, Flowering Dogwood | 2b, 3 |
| <i>Cornus florida</i> | Dogwood | 2b, 3 |
| <i>Cortaderia selloana</i> | Pampas Grass | 3 |
| <i>Cotoneaster adpressus</i> | Creeping Cotoneaster | 7 |
| <i>Cotoneaster horizontalis</i> | Cotoneaster-Variiegated Rockspray | 7 |
| <i>Cyclamen</i> spp. | Cyclamen | 7c |
| <i>Cyperus</i> spp. | Cyperus | 1 |
| <i>Delphinium</i> spp. | Larkspur | 2 |
| <i>Dianthus caryophyllus</i> | Carnation | 3, 4 |
| <i>Dianthus</i> spp. | Pink | 3, 4 |
| <i>Dieffenbachia</i> spp. | Dumb Cane | 2 |

(continued)

| BOTANICAL NAME | COMMON NAME | DISEASES |
|-------------------------------|-------------------------------|----------|
| <i>Diets iridiodes</i> | African Iris, Butterfly Iris | 4c |
| <i>Digitalis</i> spp. | Foxglove | 2, 3 |
| <i>Epipremnum</i> spp. | Pothos | 2 |
| <i>Erica dareyensis</i> | Heather | 2 |
| <i>Euonymus alata</i> | Dwarf Winged Euonymus | 2 |
| <i>Euonymus alatus</i> | Burning Bush | 2 |
| <i>Euonymus japonicus</i> | Evergreen Euonymus | 2 |
| <i>Euphorbia</i> spp. | Poinsettia | 2a |
| <i>Fatsia japonica</i> | Japanese Ratsia, Paper-Plant | 2 |
| <i>Ficus</i> spp. | Fig | 2 |
| <i>Forsythia viridissima</i> | Forsythia | 2 |
| <i>Gaillardia</i> spp. | Blanket-Flower | 2 |
| <i>Gardenia jasminoides</i> | Gardenia | 3 |
| <i>Geranium</i> spp. | Cranesbill | 5b |
| <i>Gerbera jamesonii</i> | Gerber Daisy, Transvaal Daisy | 3 |
| <i>Hedera algeriensis</i> | Algerian Ivy | 2 |
| <i>Hedera helix</i> | English Ivy | 2 |
| <i>Hibiscus moscheutos</i> | Hibiscus | 2, 3 |
| <i>Hibiscus rosa-sinensis</i> | Hibiscus | 2, 3 |
| <i>Hibiscus syriacus</i> | Rose of Sharon | 2, 3 |
| <i>Hosta</i> spp. | Hosta | 2 |
| <i>Hydrangea macrophylla</i> | French Hydrangea | 2, 3 |
| <i>Hydrangea</i> spp. | Hydrangea | 2, 3 |
| <i>Ilex</i> spp. | Holly, Winterberry, Yaupon | 3 |
| <i>Impatiens</i> spp.* | Balsam, Impatiens* | 2a, 7a |

(continued)

| BOTANICAL NAME | COMMON NAME | DISEASES |
|--------------------------------|--|-----------------|
| <i>Iris xiphium</i> | Iris (Bulbous, Spanish, Dutch) | 2e |
| <i>Itea virginica</i> | Virginia Willow | 3,4 |
| <i>Juniperus procumbens</i> | Juniper | 1a, 4 |
| <i>Juniperus scopulorum</i> | Juniper | 1a, 4 |
| <i>Juniperus</i> spp. | Juniper | 1a, 4 |
| <i>Juniperus virginiana</i> | Red Cedar | 1a, 4 |
| <i>Lagerstroemia indica</i> | Crapemyrtle | 2, 3 |
| <i>Laurus nobilis</i> | Laurel | 3 |
| <i>Lilium</i> spp. | Asiatic Lily | 2 |
| <i>Liriope muscari</i> | Lilyturf | 2 |
| <i>Lobularia maritima</i> | Sweet Alyssum | 7 |
| <i>Magnolia grandiflora</i> | Southern Magnolia | 2 |
| <i>Magnolia soulangiana</i> | Saucer Magnolia | 2 |
| <i>Magnolia</i> spp. | Magnolia | 2 |
| <i>Malus</i> spp. | Crabapple (See Table 4 for variety list) | 2i |
| <i>Nandina domestica</i> | Nandina | 2 |
| <i>Nerium oleander</i> | Oleander, Rosebay | 2 |
| <i>Pelargonium</i> spp. | Geranium | 3, 4, 5b |
| <i>Perisetum alopecuroides</i> | Grass | 2 |
| <i>Peperomia</i> spp. | Baby Rubber-Plant | 2, 7 |
| <i>Petunia</i> spp. | Petunia | 6a |
| <i>Phalaris</i> spp. | Dwarf Pampas Grass | 3 |
| <i>Philodendron</i> spp. | Philodendron | 2j |
| <i>Phlox</i> spp. | Phlox | 3 |
| <i>Phoenix dactylifera</i> | Date Palm | 2, 7 |

(continued)

| BOTANICAL NAME | COMMON NAME | DISEASES |
|----------------------------|----------------------------------|----------|
| <i>Phoenix roebelenii</i> | Roebelin's Palm | 2, 7 |
| <i>Photinia glabra</i> | Red-Tip Photinia | 2, 3, 4 |
| <i>Picea abies</i> | Norway Spruce | 1 |
| <i>Picea glauca</i> | White Spruce | 1 |
| <i>Picea purgens</i> | Blue Spruce | 1 |
| <i>Pieris japonica</i> | Japanese Andromeda | 2, 7 |
| <i>Pinus muhgo</i> | Muhgo Pine | 1b, 4 |
| <i>Pinus nigra</i> | Black Pine | 1b, 4 |
| <i>Pinus silvestris</i> | Scotch Pine | 1, 4 |
| <i>Pinus</i> spp. | Pine | 1b, 4 |
| <i>Pinus strobus</i> | Eastern White Pine | 1b, 4 |
| <i>Pittosporum</i> spp. | Australian Laurel | 3, 4 |
| <i>Pittosporum tobira</i> | Mock-Orange | 3, 4 |
| <i>Plectranthus</i> spp. | Swedish Ivy, Coleus | 2 |
| <i>Populus trichocarpa</i> | Poplar | 4 |
| <i>Populus</i> spp. | Aspen Trees | 2 |
| <i>Potentilla</i> spp. | Cinquefoil | 2 |
| <i>Primula</i> spp. | Primrose | 2 |
| <i>Prunus pumila</i> | Cherry | 2, 5 |
| <i>Prunus</i> spp. | Flowering Plum, Purple-Leaf Plum | 2, 5 |
| <i>Pseudotsuga</i> spp. | Douglas Fir | 1, 4 |
| <i>Pyrus calleryana</i> | Bradford's Pear | 3 |
| <i>Quercus falcata</i> | Red Oak | 2, 3 |
| <i>Quercus palustris</i> | Pin Oak | 2, 3 |
| <i>Rhaphiolepis indica</i> | Indian Hawthorn | 2, 3, 4 |

(continued)

| BOTANICAL NAME | COMMON NAME | DISEASES |
|-----------------------------------|--------------------------|-----------------|
| <i>Rhododendron</i> spp. | Azaleas, Rhododendron | 2b, 3, 6, 7 |
| <i>Rhododendron</i> spp. | Glacier Azalea | 2b, 3, 6, 7 |
| <i>Rosa</i> spp. | Rose | 2a, 2c, 3c, 4b |
| <i>Rosmarinus</i> spp. | Rosemary (prostrate) | 2 |
| <i>Rudbeckia hirta</i> | Black-Eyed-Susan | 2j |
| <i>Salvia</i> spp. | Sage | 3 |
| <i>Schlumbergera</i> | Holiday Cactus | 2, 7 |
| <i>Sedum</i> spp. | Orpine, Stonecrop | 2 |
| <i>Sempervivum</i> spp. | Live-forever, House-Leek | 2 |
| <i>Setaria</i> spp. | Ribbon Grass | 2, 3 |
| <i>Spathiphyllum floribundium</i> | Peace Lily | 2, 7 |
| <i>Spirea budalda</i> | Spirea | 3 |
| <i>Spirea japonica</i> | Spirea | 3 |
| <i>Syagrus romanzoffianum</i> | Queen Palm | 2 |
| <i>Tagetes</i> spp. | Marigold | 2a |
| <i>Taxus baccata</i> | Spreading Yew | 7 |
| <i>Thuja plicata</i> | Western Red Cedar | 4 |
| <i>Thuja</i> spp. | Arborvitae | 2 |
| <i>Thymus serpyllum</i> | Creeping Thyme | 2 |
| <i>Tsuga heterophylla</i> | Western Hemlock | 4 |
| <i>Tsuga</i> spp. | Hemlock | 4 |
| <i>Verbenas</i> spp. | Verbena, Vervain | 3 |
| <i>Viburnum</i> spp. | Viburnum | 2,3,4 |
| <i>Vinca</i> spp. | Periwinkle | 2, 6a |
| <i>Viola</i> spp.* | Viola, Pansy* | 2 |

(continued)

| BOTANICAL NAME | COMMON NAME | DISEASES |
|------------------------|--------------|----------|
| <i>Weigela florida</i> | Pink Weigela | 2 |
| <i>Yucca</i> spp. | Yucca | 7 |
| <i>Zinnia</i> spp. | Zinnia | 2a, 3 |

* Do not exceed 3.85 fl. oz./100 gallons on these species

TABLE 3 Tolerant Plants Listed by Common Name

| COMMON NAME | BOTANICAL NAME |
|---------------------------------|----------------------------|
| Abelia | <i>Abelia</i> spp. |
| Andromeda, Japanese | <i>Pieris japonica</i> |
| Arborvitae | <i>Thuja</i> spp. |
| Aspen Trees | <i>Populus</i> spp. |
| Aster | <i>Aster</i> spp. |
| Aucuba, Japanese | <i>Aucuba japonica</i> |
| Azalea, Glacier | <i>Rhododendron</i> spp. |
| Azaleas | <i>Rhododendron</i> spp. |
| Balsam | <i>Impatiens</i> spp. |
| Barberry | <i>Berberis thunbergii</i> |
| Begonia (except Rieger begonia) | <i>Begonia</i> spp. |
| Birch, River | <i>Betula nigra</i> |
| Black-Eyed-Susan | <i>Rudbeckia hirta</i> |
| Blanket-Flower | <i>Gaillardia</i> spp. |
| Bougainvillea | <i>Bougainvillea</i> spp. |
| Boxwood | <i>Buxus sempervirens</i> |
| Buddleia | <i>Buddleia davidii</i> |
| Bugle | <i>Ajuga reptans</i> |

(continued)

| COMMON NAME | BOTANICAL NAME |
|--|---|
| Bugleweed | <i>Ajuga reptans</i> |
| Burning Bush | <i>Euonymus alatus</i> |
| Butterfly Bush | <i>Buddleia davidii</i> |
| Cactus, Holiday | <i>Schlumbergera</i> |
| Caladium | <i>Caladium</i> spp. |
| Camellia | <i>Camellia japonica</i> |
| Carnation | <i>Dianthus caryophyllus</i> |
| Ceanothus | <i>Ceanothus</i> spp. |
| Cedar, Atlas | <i>Cedrus atlantica</i> |
| Cedar, Red | <i>Juniperus virginiana</i> |
| Cedar, Western Red | <i>Thuja plicata</i> |
| Cedar, White | <i>Cedrus</i> spp. |
| Cherry | <i>Prunus pumila</i> |
| Christmas Tree | See Fraser fir, Scotch pine and Douglas fir |
| Chrysanthemum | <i>Chrysanthemum</i> spp. |
| Cinquefoil | <i>Potentilla</i> spp. |
| Clethra | <i>Clethra ainifolia</i> |
| Coleus | <i>Plectranthus</i> spp. |
| Cotoneaster, Creeping | <i>Cotoneaster adpressus</i> |
| Cotoneaster, Variegated Rockspray | <i>Cotoneaster horizontalis</i> |
| Cranapple (See Table 4 for variety list) | <i>Malus</i> spp. |
| Cranesbill | <i>Geranium</i> spp. |
| Crapemyrtle | <i>Lagerstroemia indica</i> |
| Cyclamen | <i>Cyclamen</i> spp. |
| Cyperus | <i>Cyperas</i> spp. |

(continued)

| COMMON NAME | BOTANICAL NAME |
|------------------------|---------------------------------|
| Cypress, Sawara | <i>Chamaecyparis pisifera</i> |
| Cypress, Leyland | <i>Chamaecypans</i> spp. |
| Daisy, Gerber | <i>Gerbera jamesonii</i> |
| Daisy, Transvaal | <i>Gerbera jamesonii</i> |
| Dogwood | <i>Cornus</i> spp. |
| Dogwood | <i>Cornus florida</i> |
| Dogwood, Pink | <i>Cornus</i> spp. |
| Dumb-Cane | <i>Dieffenbachia</i> spp. |
| Euonymus, Dwarf Winged | <i>Euonymus alatus</i> |
| Euonymus, Evergreen | <i>Euonymus japonicus</i> |
| Evergreen, Chinese | <i>Aglaotiema</i> spp. |
| Fatsia, Japanese | <i>Fatsia japonica</i> |
| Fig | <i>Ficus</i> spp. |
| Fir, Douglas | <i>Pseudotsuga</i> spp. |
| Fir, Fraser | <i>Abies fraseri</i> |
| Fir, Noble | <i>Abies procera</i> |
| Floss-Flower | <i>Ageratum</i> spp. |
| Forsythia | <i>Forsythia viridissima</i> |
| Foxglove | <i>Digitalis</i> spp. |
| Gardenia | <i>Gardenia jasminoides</i> |
| Geranium | <i>Pelargonium</i> spp. |
| Grass | <i>Pennisetum alopecuroides</i> |
| Grass, Dwarf Pampas | <i>Phalaris</i> spp. |
| Grass, Pampas | <i>Cortaderia selloana</i> |
| Hawthorn, Indian | <i>Rhaphiolepis indica</i> |

(continued)

| COMMON NAME | BOTANICAL NAME |
|--------------------------------|-------------------------------|
| Heather | <i>Erica dareyensis</i> |
| Hemlock | <i>Tsuga</i> spp. |
| Hemlock, Western | <i>Tsuga heterophylla</i> |
| Hibiscus | <i>Hibiscus moscheutos</i> |
| Hibiscus | <i>Hibiscus rosa-sinensis</i> |
| Holly | <i>Hex</i> spp. |
| Hosta | <i>Hosta</i> spp. |
| House-Leek | <i>Sempervivum</i> spp. |
| Hydrangea | <i>Hydrangea</i> spp. |
| Hydrangea, French | <i>Hydrangea macrophylla</i> |
| Impatiens* | <i>Impatiens</i> spp.* |
| Iris (Bulbous, Spanish, Dutch) | <i>Iris xiphium</i> |
| Iris, African | <i>Diets irioides</i> |
| Iris, Butterfly | <i>Diets irioides</i> |
| Ivy, Algerian | <i>Hedera aigeriensis</i> |
| Ivy, English | <i>Hedera helix</i> |
| Ivy, Swedish | <i>Plectranthus</i> spp. |
| Juniper | <i>Juniperus procumbens</i> |
| Juniper | <i>Juniperus scopulorum</i> |
| Juniper | <i>Juniperus</i> spp. |
| Larkspur | <i>Delphinium</i> spp. |
| Laurel | <i>Laurus nobilis</i> |
| Laurel, Australian | <i>Pittosporum</i> spp. |
| Laurel, Japanese | <i>Aucuba japonica</i> |
| Lilac, California | <i>Ceanothus</i> spp. |

(continued)

| COMMON NAME | BOTANICAL NAME |
|--------------------|-----------------------------------|
| Lilac, Wild | <i>Ceanothus sanguineus</i> |
| Lily, Asiatic | <i>Lilium</i> spp. |
| Lily, Peace | <i>Spathiphyllum floribundium</i> |
| Lilyturf | <i>Uriope muscari</i> |
| Live-Forever | <i>Sempervivum</i> spp. |
| Magnolia | <i>Magnolia</i> spp. |
| Magnolia, Saucer | <i>Magnolia soutangiana</i> |
| Magnolia, Southern | <i>Magnolia grandiflora</i> |
| Maple, Japanese | <i>Acer palmatum</i> |
| Maple Sugar | <i>Acer saccharum</i> |
| Marigold | <i>Tagetes</i> spp. |
| Mock-Orange | <i>Pittosporum tobira</i> |
| Mugwort | <i>Artemisia</i> app. |
| Nandina | <i>Nandina domestica</i> |
| Oak, Pin | <i>Quercus patustris</i> |
| Oak, Red | <i>Quercus falcata</i> |
| Oleander | <i>Nerium oleander</i> |
| Orpine | <i>Sedum</i> spp. |
| Palm, Date | <i>Phoenix dactylifera</i> |
| Palm, Parlor | <i>Chamaedorea elegans</i> |
| Palm, Queen | <i>Syagris romanzoffianum</i> |
| Palm, Roebelin's | <i>Phoenix roebelenii</i> |
| Palm, Sago | <i>Cycas urens</i> |
| Pansy* | <i>Viola</i> spp.* |
| Paper Plant | <i>Fatsia japonica</i> |

(continued)

| COMMON NAME | BOTANICAL NAME |
|---------------------|----------------------------|
| Pear Bradford's | <i>Pyrus calleryana</i> |
| Periwinkle | <i>Vinca</i> spp. |
| Petunia | <i>Petunia</i> spp. |
| Philodendron | <i>Philodendron</i> spp. |
| Phlox | <i>Phlox</i> spp. |
| Photinia, Red-Tip | <i>Photinia glabra</i> |
| Pine | <i>Pinus</i> spp. |
| Pine, Black | <i>Pinus nigra</i> |
| Pine, Eastern White | <i>Pinus strobus</i> |
| Pine, Muhgo | <i>Pinus muhgo</i> |
| Pine, Scotch | <i>Pinus sylvestris</i> |
| Pink | <i>Dianthus</i> spp. |
| Plum, Flowering | <i>Prunus</i> spp. |
| Plum, Purple-Leaf | <i>Prunus</i> spp. |
| Poinsettia | <i>Euphorbia</i> spp. |
| Poplar | <i>Populus trichocarpa</i> |
| Pothos | <i>Epipremnum</i> spp. |
| Primrose | <i>Primula</i> spp. |
| Pussy's-Foot | <i>Ageratum</i> spp. |
| Redbud, Western | <i>Cercis occidentalis</i> |
| Rhododendron | <i>Rhododendron</i> spp. |
| Ribbon-Grass | <i>Setaria</i> spp. |
| Rose of Sharon | <i>Hibiscus syriacus</i> |
| Rose | <i>Rosa</i> spp. |
| Rosebay | <i>Nerium oleander</i> |

(continued)

| COMMON NAME | BOTANICAL NAME |
|----------------------|------------------------------|
| Rosemary (Prostrate) | <i>Rosmarinus</i> spp. |
| Rubber-Plant, Baby | <i>Peperomia</i> spp. |
| Rubber Tree | <i>Brassaia actinophylla</i> |
| Sage | <i>Salvia</i> spp. |
| Sagebrush | <i>Artemisia</i> spp. |
| Snapdragon | <i>Antirrhinum</i> spp. |
| Snowball | <i>Ceanothus</i> spp. |
| Spirea | <i>Spirea budaida</i> |
| Spirea | <i>Spirea japonica</i> |
| Spruce, Blue | <i>Picea pungens</i> |
| Spruce, Norway | <i>Picea abies</i> |
| Spruce, White | <i>Picea glauca</i> |
| Starwort | <i>Aster</i> spp. |
| Stonecrop | <i>Sedum</i> spp. |
| Sweet Alyssum | <i>Lobularia maritima</i> |
| Thymes, Creeping | <i>Thymus serpyllum</i> |
| Umbrella-Tree | <i>Brassamactinophylla</i> |
| Verbena | <i>Verbena</i> spp. |
| Vervain | <i>Verbena</i> spp. |
| Viburnum | <i>Viburnum</i> spp. |
| Vinca | <i>Catharanthus roseus</i> |
| Viola | <i>Viola</i> spp. |
| White Alder | <i>Ciethora</i> spp. |
| Weigela, Pink | <i>Weigeia florida</i> |
| Willow, Virginia | <i>Itea virginica</i> |

(continued)

| COMMON NAME | BOTANICAL NAME |
|----------------|------------------------|
| Winterberry | <i>Ilex</i> spp. |
| Wormwood | <i>Artemisia</i> spp. |
| Yaupon | <i>Ilex</i> spp. |
| Yew, Spreading | <i>Taxus baccata</i> |
| Yucca | <i>Yucca</i> spp. |
| Zebra-Plant | <i>Aphelandra</i> spp. |
| Zinnia | <i>Zinnia</i> spp. |

* Do Not Exceed 3.85 fl. oz./100 gallons on these species.

TABLE 4. Tolerant Varieties of Crabapple Species (Genus Malus) Tolerant Varieties of Malus

| | | | |
|--|-------------------|------------------|-----------------------|
| Arkansas Black | Eleyi | Mary Potter | <i>seiboldii</i> |
| <i>atrosanguinea</i> | Enterprise | Molten Lava | Selkirk |
| <i>baccata</i> | Evereste | New Centennial | Sentinel |
| <i>baccata</i> var. <i>jackii</i> | Eyeiynn | Ormiston Roy | Silver Moon |
| <i>baccata</i> var. <i>mandshurica</i> | <i>floribunda</i> | Pink Satin | Silverdrift |
| Callaway | Gloriosa | Prairie Maid | Sinai Fire |
| Candy mint Sargent | Golden Delicious | Prairifire | <i>spectabilis</i> |
| Christmas Holly | Golden Raindrops | Profusion | Sugar Tyme |
| <i>coronaria</i> | Hopa | <i>pumila</i> | Van Eseltine |
| David | Indian Magic | Ralph Shay | White Angel |
| Dolgo | Island | Red Jade | Williams Pride |
| Donald Wyman | Katherine | Red Baron | Winter Gold |
| Dorothea | Lancelot | Sargent | Yellow Delicious |
| Doubloons | Louisa | <i>sargentii</i> | <i>zumi Calocarpa</i> |

TABLE 5. Intolerant Plants (Do not apply Strobe™ 2L to these species or varieties)

| COMMON NAME | BOTANICAL NAME |
|--|--|
| Apple | <i>Malus domestica</i> |
| Crabapple - Flame variety | <i>Malus</i> spp. |
| Crabapple - Brandywine variety | <i>Malus</i> spp. |
| Crabapple - Novamac variety | <i>Malus</i> spp. |
| Cherry, Flowering - Yoshina variety | <i>Prunus yedoensis</i> |
| Leatherleaf Fern and Other Ferns for cut foliage | <i>Rumohra adiantiformis</i> and other species for cut foliage |
| Privet | <i>Ligustrum</i> spp. |

CONIFERS INCLUDING CHRISTMAS TREES, COMMERCIAL PRODUCTION ROSES

Strobe™ 2L may be used to control certain diseases on conifers in production (indoor and outdoor) and landscape situations.

Please see the Ornamental Section above for more detailed directions for use in landscape situations.

| Crop | Target Diseases | Use Rate fl. oz. product/Acre (lb ai/A) | Remarks |
|------------------------------------|---|--|---|
| Conifers including Christmas Trees | Diplodia tip blight (<i>Diplodia pinea</i>) Lophodermium needlecast (<i>Lophodermium pinastri</i>) Swiss needlecast (<i>Phaeocryptopus gaumannii</i>) | 6.1 - 15.3 (0.10 - 0.25) | Integrated Pest (Disease) Management: Strobe™ 2L should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance and removal of plant debris in which inoculum may overwinter. Resistance Management: Do not apply more than four sequential applications of Strobe™ 2L before alternating with a fungicide that is not in Group 11. Do not make more than eight applications of Strobe™ 2L per acre per year. Application Directions: Strobe™ 2L applications should begin prior to disease development and continue throughout the season at 7-21 day intervals following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at recommended rates. |

(continued)

| Crop | Target Diseases | Use Rate fl. oz. product/Acre (lb ai/A) | Remarks |
|---|--|--|--|
| Roses (Commercial Rose Production) | Downy mildew (<i>Peronospora sparsa</i>) Powdery mildew (<i>Spherotheca pannosa</i>) Rust (<i>Phragmidium mucronatum</i> , <i>P. tuberculatum</i> , and other <i>Phragmidium</i> spp.) Septoria leaf spot (<i>Septoria rosae</i>) Alternaria leaf spot (<i>Alternaria alternata</i>) | 3.0 - 15.3 (0.05 - 0.25) | Integrated Pest (Disease) Management: Strobe™ 2L should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation. Resistance Management: Do not make more than four sequential applications of Strobe™ 2L before alternating with a fungicide that is not in Group 11. Do not make more than eight applications per acre per year. Application Directions: Strobe™ 2L application should begin prior to disease development and continue throughout the season on 7-21 day intervals following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at recommended rates. Plant Safety: Strobe™ 2L has been shown to be safe when applied to roses. However, all varieties of roses have not been evaluated for safety. Small scale variety safety testing must be conducted to insure plant safety prior to large scale application. In addition, do not tank-mix Strobe™ 2L with other fungicides, insecticides, herbicides, fertilizer, etc. unless local experience indicates that the tank-mix is safe to roses. |

Specific Use Restrictions: Do not apply more than 123 fluid ounces of product/acre/season (2.0 lb ai/A).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of CONTROL SOLUTIONS INC. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold CONTROL SOLUTIONS INC. and Seller harmless for any claims relating to such factors.

CONTROL SOLUTIONS INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or CONTROL SOLUTIONS INC., and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, CONTROL SOLUTIONS INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall CONTROL SOLUTIONS INC. be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

CONTROL SOLUTIONS INC. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of CONTROL SOLUTIONS INC.

NOTES

NOTES

Manufactured by:
Control Solutions Inc.
Pasadena, Texas 77507

4/27/2015

QUALI-PRO

Strobe™ 2L

BROAD SPECTRUM FUNGICIDE

FOR CONTROL OF TURF AND ORNAMENTAL DISEASES

ACTIVE INGREDIENT:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yl]oxyphenyl]-3-methoxyacrylate*.....

22.9%

OTHER INGREDIENTS:.....

77.1%

TOTAL: 100.0%

*IUPAC

Contains 2.08 lb. of active ingredient per gallon
Suspended Concentrate



**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See additional precautionary statements and
directions for use inside booklet.

Reformulation is prohibited. See individual container
labels for repackaging limitations.

EPA Reg. No. 42750-261-53883

EPA Est. No. 53883-TX-002

Manufactured by:



**Control
Solutions Inc.**

Pasadena, TX 77507

CONTENTS: 1 GALLON

GROUP 11 FUNGICIDE