

Supplemental Labeling



EPA Reg. No. 62719-442-5905

Control of Lepidoptera Larvae Infesting Sorghum (Grain and Sweet) (Not for Use and Distribution in New York State)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Troubadour™ 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Troubadour 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Troubadour 2F.

Directions for Use

Refer to product label for General Information, Mixing and Application directions.

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
southwestern corn borer sugarcane borer	8 – 10 (0.12 – 0.16lb ai/acre)	Apply at first sign of egg hatch or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities. Apply as broadcast or multinozzle over the row application to mid- and late-season infestations.	<ul style="list-style-type: none"> • Do not apply more than 12 fl oz per acre per application or 48 fl oz of Troubadour 2F (0.75 lb ai) per acre per season. • Preharvest Interval: Do not apply within 21 days of grain or stover harvest, or within 3 days of forage or sweet sorghum stalk harvest.
beet armyworm fall armyworm		Apply at first sign of egg hatch, feeding damage, or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities. Under heavy infestations, continuous moth flights, or rapid crop growth and development, reapply at 10-day re-treatment intervals.	<ul style="list-style-type: none"> • See Rotational Crop Restrictions on the label affixed to the container.

Ground Application: Apply in a minimum of 15 gallons per acre by conventional ground equipment. Use a spray volume that assures uniform coverage of the infested portions of the treated crop. Calibrate equipment to the desired spray volume.

Aerial Application: Apply in a minimum of 10 gallons per acre.

Spray Adjuvants: The addition of agricultural adjuvants to sprays of Troubadour 2F may improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Helena Chemical Company recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

™ Trademark of Helena Holding Company

**Manufactured for
Helena Chemical Company
225 Schilling Boulevard, Suite 300
Collierville, TN 38017**