



Specimen Label

X-CYTE™

A Plant Growth Regulator and Yield Stimulant

ACTIVE INGREDIENTS

Cytokinin, as kinetin, based on biological activity

0.04%

INERT INGREDIENTS

99.96%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

FIRST AID


If on skin or clothing	Take off clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Do not induce vomiting unless told to do so by the poison control center or doctor. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

EPA Reg. No. 57538-15
EPA Est. No. 57538- TX-2

EPA Reg. No. 57538-15
EPA Est. No. 57538- TX-2

NET CONTENTS

(8.5 lb/ gallon or 1.02 kg/ liter)



Manufactured by



Manufactured for Stoller Enterprises, Inc.
4001 W. Sam Houston Parkway North, Suite 100
Houston Texas 77043 USA
Phone: 1-800-539-5283 or 713-461-1493 | Fax: 713-461-4467
E- Mail: stoller@stollerusa.com | Website:
www.StollerUSA.com

STOLLER® X-Cyte™

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Exposed treated seed may be hazardous to birds and other wildlife. Treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of all excess treated seed and seed packaging by burial away from streams and bodies of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms and in forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted- entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, wear: coveralls over long- sleeved shirt and pants, waterproof gloves and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool place and out of direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION

Apply this product only through the following types of irrigation systems. Do not apply through any other types of irrigation systems. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non- uniform distribution of treated water. If you have questions about calibration, you should contact State Experiment Station specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label- prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

A. Center Pivot, Traveler, Big Gun, Motorized Lateral Move, End Tow, and Side (Wheel) Roll Irrigation Equipment: Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank of injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and acreage contained in circle or run. Mix recommended amount of product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution of run, but continue to operate irrigation system until product has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

B. Solid Set and Hand Move Irrigation Equipment: Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty – five minute period. Mix desired amount of product for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to insure that product will remain in suspension during the injection cycle. Product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until pesticide is cleared from last sprinkler head.

CROP USAGE – ALL CROPS LISTED FOR STRESS RELIEF

Use 1 pint X- CYTE per acre (1.2 liters per hectare) any time a crop is prematurely dying down (loss of color) due to stress caused by one or more of the following conditions: weather (frost, drought, excessive moisture), insect infestation, fungus attack, and/ or herbicide burn.

SAFETY DEVICES

(1) The systems designated above must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. (2) All pesticide injection pipelines must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. (3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid- operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. (5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

SYSTEMS CONNECTED TO PUBLIC WATER SOURCES

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year. Chemigation systems connected to public water systems must contain a functional, reduced- pressure, zone, backflow presenter (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

For additional instructions on safety precautions, refer to statements (2), (3), (4), (6) and (7) in the section SAFETY DEVICES.

GENERAL USE INSTRUCTIONS

For best results, **X- CYTE** should be applied before noon or after 4 p.m. a spreader- sticker (surfactant) cleared for application to growing crops should be used with the product. Store **X- CYTE** in a cool place and out of direct sunlight. Any spigot or pump put into an **X- CYTE** drum should be cleaned with disinfectant (bleach and water, etc.) before using, unless the whole drum is to be used at one time. Mix **X- CYTE** with enough water to get thorough coverage of plant surfaces. **X- CYTE** is compatible with most other spray materials.

CUCURBITS - 0.5 to 1 pint/ acre (0.6 to 1.2 lt/ ha) each application (Cucumbers, Muskmelon, Cantaloupe, Watermelon, Honey Dew, Squash, etc.):

- 1st application: 4 to 8- inch stage.
- 2nd application: at early bloom.
- 3rd application: start of fruiting.

GRAPES - 1 pint/ acre (1.2 lt/ ha) each application:

- 1st application: between leafout and prebloom.
- 2nd application: at petal fall.
- 3rd application: after harvest.

ORANGES - 1 pint/ acre (1.2 lt/ ha) each application:

- 1st application: at prebloom.
- 2nd application: at calyx (petal fall).
- 3rd application: 3 weeks after 2nd spraying.
- 4th application: 4 weeks after 3rd spraying.

CROP USAGE – ALL CROPS LISTED

FOR TRANSPLANTING AND SEED BED TREATMENT

Use 2 pints **X- CYTE** per acre (2.4 lt/ ha) or 1 part **X- CYTE** to 1000 parts water (approximately 1 tablespoon **X- CYTE** to 1 gallon or 4 liters of water) as a root dip and watering solution when transplanting.

Use 2 pints **X- CYTE** per acre (2.4 lt/ ha) applied to the seed bed at time of seeding or up to 20 days thereafter.

ALFALFA – 1 pint/ acre (1.2 lt/ ha) each application.

- 1st application: after cutting, with repeat sprays at 14 to 21 day intervals.

APPLES – 1 pint/ acre (1.2 lt/ ha) each application.

- 1st application: at full pink.
- 2nd application: at calix (petal fall).
- 3rd application: 3 weeks after 2nd spraying.
- 4th application: 4 weeks after 3rd spraying.

ASPARAGUS – 1 to 2 pints/ acre (1.2 to 2.4 lt/ ha) each application.

- 1st application: spray crowns when growth begins.
- 2nd application: spray crowns after each cutting.

BANANAS – 1 to 2 pints per 100 gallons of water (1.2 to 2.4 lt in 400 lt of water)

- 1st application: apply just prior to flower bud formation.
- Alternatively: make one application one week following transplant or at the start of new sucker growth.
- Repeat at intervals of 1-2 months until harvest.

BEANS – 0.5 to 1 pint/ acre (0.6 to 1.2 lt/ ha) each application

- 1st application: 4-5 inch stage.
- 2nd application: at early bloom.
- 3rd application: at early pod set.

CARROTS – 1 pint/ acre (1.2 lt/ ha) each application

- 1st application: at tuber initiation.
- 2nd application: 2-3 weeks after first spraying.

CELERY

- 1st application: use 2 pints/ acre (2.4 lt/ ha) applied to the seed bed at time of seeding or up to 20 days thereafter.
- 2nd application: use 2 pints/ acre (2.4 lt/ ha) at the time seedlings are transplanted, see transplanting instructions above.
- 3rd application: use 1 pint/ acre (1.2 lt/ ha) 2-3 weeks after transplanting.

CORN - 1 pint/ acre (1.2 lt/ ha) each application

- 1st application: at the 1 to 1.5 foot stage.
- 2nd application: at tassel time.

COTTON - 1 pint/ acre (1.2 lt/ ha) each application

- 1st application: at pinhead square with repeat applications at 14 to 21 day intervals.

CRUCIFEROUS CROPS - 0.5 to 1 pint/ acre (0.6 to 1.2 lt/ ha) each application.

- (Cabbage, Broccoli, Cauliflower, Brussels Sprouts)
- 1st application: 3 to 4- inch stage. Repeat at 10 to 14 day intervals.

SUGAR BEETS - 1 pint/ acre (1.2 lt/ ha) each application

- 1st application: at tuber initiation.
- 2nd application: 2-3 weeks after 1st spraying.

PEACHES - 1 pint/ acre (1.2 lt/ ha) each application:

1st application: at prebloom.

2nd application: at calyx (petal fall).

3rd application: 3 weeks after 2nd spraying.

4th application: 4 weeks after 3rd spraying.

PEANUTS - 1 pint/ acre (1.2 lt/ ha) each application:

1st application: at pegging.

2nd application: 2-3 weeks after 1st spraying.

PEAS - 0.5 to 1 pint/ acre (0.6 to 1.2 lt/ ha) each application:

1st application: 3 to 4- inch stage.

2nd application: Prebloom.

3rd application: at early pod set.

PEPPERS AND EGGPLANT - 0.5 to 1 pint/ acre (0.6 to 1.2 lt/ ha) each application:

1st application: just prior to 1st bloom.

2nd application: 10 days after 1st spraying

3rd application: 10 days after 2nd spraying.

POTATOES - 1 pint/ acre (1.2 lt/ ha) each application:

1st application: at tuber set. The time of application is determined by pulling an average size plant in the field 4 weeks (and every 7 days thereafter if necessary) after planting. Observe the roots to see if tubers are forming. Anytime you see the small tubers forming, it is time for the

1st application. Usually tubers start to set 5 to 6 weeks after planting.

2nd application: at full blossom. Russet Burbanks, which do not show full blossom, should be sprayed 2-3 weeks after 1st spray.

RICE - 1 pint/ acre (1.2 lt/ ha) each application

1st application: at 2 to 5 leaf stage with repeat application 14 to 21 days after.

SOYBEANS - 1 pint/ acre (1.2 lt/ ha) each application

Application: at first bud formation.

SPINACH AND LETTUCE - 0.5 to 1 pint/ acre (0.6 to 1.2 lt/ ha) each application.

Application: 3 to 4- inch stage.

STRAWBERRIES

1st application: use 2 pints **X- CYTE** per acre (2.4 lt/ ha) as a transplant solution. See "Transplanting Instructions" above.

2nd application: use 1 pint **X- CYTE** per acre (1.2 lt/ ha) at prebloom.

3rd application: use 1 pint **X- CYTE** per acre (1.2 lt/ ha) at petal fall.

4th application: use 1 pint **X- CYTE** per acre (1.2 lt/ ha) after harvest.

TOMATOES

1st application: use 2 pints **X- CYTE** per acre (2.4 lt/ ha) applied to the seed bed at time of seeding or up to 20 days thereafter.

2nd application: use 2 pints **X- CYTE** per acre (2.4 lt/ ha) at the time seedlings are transplanted. See "Transplanting Instructions".

3rd application: use 1 pint **X- CYTE** per acre (1.2 lt/ ha) 2 to 3 weeks after 1st bloom.

WHEAT - 1 pint/ acre (1.2 lt/ ha) each application

Application: 1-2 weeks before boot stage.

ORNAMENTAL TREES AND HERBACEOUS PLANTS

Apply 2 pints per acre (2.4 lt/ ha) in transplant water.

Apply 1 pint per acre (1.2 lt/ ha) as a foliar spray when growth begins in the early spring.

Apply 1 pint per acre (1.2 lt/ ha) at bud burst.

Apply 1 pint per acre (1.2 lt/ ha) at bud set.

Apply 1 pint per acre (1.2 lt/ ha) at the end of summer to maintain color through autumn.

SEED TREATMENT

Use only on seeds for crops listed elsewhere on this label. Do not use treated seed for food, feed or oil purposes. Commercially treated seed must be labeled in accordance with the requirements of the Federal Seed Act and applicable State laws. An approved dye must be added to distinguish treated seed and prevent inadvertent use for food, feed, or oil purposes.

Per hundredweight (cwt) of seed, dilute 2 fl. oz. **X- CYTE** (65 ml per 50 kg of seed) in equal amounts of water and mist spray on seed. **X- CYTE** can be poured/ mixed on seed in hopper at planting.

CONDITIONS OF SALE: recommendations for use of this product are based upon tests believed to be reliable. Because time, place, rate of application and other conditions of use are beyond the control of the manufacturer and/ or responsibility resulting from its misuse, Seller liability from handling, storage and use of this product is limited to replacement of product or reimbursement of purchase price.